

**Supplementary Table 1.** Pearson correlation coefficients (*P* values) between standardized biomarkers (CA125, HE4, mesothelin, B7-H4, DcR3, and spondin-2) assessed at baseline\*

Biomarker	Pearson correlation coefficients ( <i>P</i> values)					
	Ca125	HE4	Mesothelin	B7-H4	DcR3	Spondin-2
<b>All control subjects (n = 70)</b>						
CA125	1.00					
HE4	-.08 (.49)	1.00				
Mesothelin	-.08 (.51)	.20 (.10)	1.00			
B7-H4	.20 (.10)	-.17 (.38)	.14 (.25)	1.00		
DcR3	.14 (.26)	-.06 (.63)	.25 (.04)	.30 (.04)	1.00	
Spondin 2	.37 (.002)	.18 (.15)	.01 (.93)	.41 (.003)	.44 (<.001)	1.00
<b>All cancer patients (n = 34)</b>						
CA125	1.00					
HE4	.04 (.82)	1.00				
Mesothelin	.26 (.14)	.07 (.68)	1.00			
B7-H4	.16 (.36)	-.05 (.77)	.29 (.10)	1.00		
DcR3	-.10 (.59)	.24 (.17)	.14 (.42)	.08 (.66)	1.00	
Spondin-2	.23 (.19)	.25 (.16)	.26 (.14)	.40 (.02)	.59 (<.001)	1.00

\**P* values are based on *t* tests. All statistical tests were two-sided. HE4 = human epididymis protein 4; DcR3 = decoy receptor 3.