

Supplemental Table 1: Effects of mutations on salivary gland nucleoli.

Locus	Encoded Protein/Molecular Function	Alleles Tested	Refs.	# Nucleoli		
				Mean ± SD	Range	^c P values
Wild type ^a				1±0	1	NA
<i>Su(var)3-9</i>	H3K9 methyltransferase	1699 (missense), 6 and 17 (nulls)	1,2	2.7± 1.4 N = 54	1 to 9 (null); 1 to 12 (1699)	<0.001
<i>Su(var)2-5 / HP1</i>	chromodomain/ binds H3K9me	transheterozygous 1009/1209 (nulls)	1	2.8± 0.83 N = 21	1 to 4	<0.001
<i>Su(var)2-10 / dPIAS</i>	Protein Inhibitor of Activated STAT / SUMOylation	transheterozygous 02/Pex14A (nulls)	3	multiple ^b	N.D. ^b	
<i>Su(TDA-PEV) 1650</i>	??	1650	1	3.6± 1.6 N = 35	1 to 8	<0.001
<i>Su(var)3-7</i>	zinc Finger/ DNA binding	234	1	1 ^b		
<i>I(3)73Ah</i>	ubiquitin ligase	1044	1	1 ^b		
<i>Su(TDA-PEV) 1025</i>	??	1025	1	1 ^b		
<i>Su(TDA-PEV) 1260</i>	??	1260	1	1 ^b		
<i>Su(TDA-PEV) 1128</i>	??	1128	1	1 ^b		
<i>Su(var)4-20</i>	H4K20 methyltransferase	BG00814 and EY07422 P insertions (hypomorphic)	4	1 ^b		
<i>dSIR2</i>	NAD-dependent histone deacetylase	17 (null)	5	1 ^b		
<i>Pc</i>	PcG complex	1 and 7 (nulls)	6,7	1 ^b		
<i>Ph</i>	PcG complex	410 (null)	8	1 ^b		
<i>Lig4</i>	DNA ligase	29 and 57 (nulls)	9	1 ^b		
<i>Lig4; Su(var)3-9</i>	N.A.	double transheterozygous mutant		1.7± 0.8 N = 83	1 to 4	<0.001 ^d
<i>Ago2</i>	siRNA loading	51B (null)	10	2.5± 1.3 N = 79	1 to 5	0.001
<i>Aub</i>	RNAi complex	QC42 (?)	11	1.1± 0.4 N = 85	1 to 3	0.004
<i>hls / Spn-E</i>	RNA helicase (of RNAi complex)	Δ215 (null) and 1 (hypomorph)	12,13	1.1± 0.3 N = 40 and 1.6± 0.8 N = 78	1 to 4	0.083 and <0.001

SUPPLEMENTARY INFORMATION

<i>piwi</i>	mRNA binding (of RNAi complex)	06843 (null)	¹⁴	1.8± 1.0 N = 102	1 to 5	<0.001
<i>dcr-2</i>	RNase III-like endonuclease and helicase for siRNA production	<i>L811fsx</i> (hypomorph) and P insertion (null)	¹⁵	3.4± 1.9 N = 60	1 to 6	<0.001

^a heterozygous for *Su(var)*

^b evaluated qualitatively from multiple images

^c p values reflect comparisons of the mean # nucleoli in wild type versus mutant

^d p values reflect comparisons of the mean # nucleoli in the double mutant versus wild type, and separately versus *Su(var)3-9^{null}* mutant

1. Donaldson, K.M., Lui, A. & Karpen, G.H. Modifiers of terminal deficiency-associated position effect variegation in Drosophila. *Genetics* **160**, 995-1009 (2002).
2. Schotta, G. et al. Central role of Drosophila SU(VAR)3-9 in histone H3-K9 methylation and heterochromatic gene silencing. *Embo J* **21**, 1121-31 (2002).
3. Hari, K.L., Cook, K.R. & Karpen, G.H. The Drosophila Su(var)2-10 locus regulates chromosome structure and function and encodes a member of the PIAS protein family. *Genes Dev* **15**, 1334-48 (2001).
4. Schotta, G. et al. A silencing pathway to induce H3-K9 and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev* **18**, 1251-62 (2004).
5. Astrom, S.U., Cline, T.W. & Rine, J. The Drosophila melanogaster sir2+ gene is nonessential and has only minor effects on position-effect variegation. *Genetics* **163**, 931-7 (2003).
6. Gindhart-Jr, J.G. & Kaufman, T.C. Identification of Polycomb and trithorax Group Responsive Elements in the Regulatory Region of the Drosophila Homeotic Gene Sex combs reduced. *Genetics* **139**, 797-814 (1995).
7. Tearle, R.G. & Nusslein-Volhard, C. *Tubingen mutants and stock list*, 209--269 (1987).
8. Hodgson, J.W. et al. The polyhomeotic locus of Drosophila melanogaster is transcriptionally and post-transcriptionally regulated during embryogenesis. *Mech Dev* **66**, 69-81 (1997).
9. Romeijn, R. & Ferro, W. ligase4 mutations. (2004).
10. Xu, K. et al. The fragile X-related Gene Affects the Crawling Behavior of Drosophila Larvae by Regulating the mRNA Level of the DEG/ENaC Protein Pickpocket1. *Current Biology* **14**, 1025-1034 (2004).
11. Wilson, J.E., Connell, J.E., Schlenker, J.D. & Macdonald, P.M. Novel genetic screen for genes involved in posterior body patterning in Drosophila. *Dev Genet* **19**, 199-209 (1996).
12. Gillespie, D.E. & Berg, C.A. Homeless is required for RNA localization in Drosophila oogenesis and encodes a new member of the DE-H family of RNA-dependent ATPases. *Genes Dev* **9**, 2495-508 (1995).
13. Gonzalez-Reyes, A., Elliott, H. & St Johnston, D. Oocyte determination and the origin of polarity in Drosophila: the role of the spindle genes. *Development* **124**, 4927-4937 (1997).
14. Lin, H. & Spradling, A.C. A novel group of pumilio mutations affects the asymmetric division of germline stem cells in the Drosophila ovary. *Development* **124**, 2463-2476 (1997).
15. Lee, Y.S. et al. Distinct Roles for Drosophila Dicer-1 and Dicer-2 in the siRNA/miRNA Silencing Pathways. *Cell* **117**, 69 (2004).