

PARTITIONING OF PERFLUOROCTANOATE INTO PHOSPHATIDYLCHOLINE BILAYERS IS CHAIN LENGTH-INDEPENDENT

Wei Xie^a, Geoffrey D. Bothun^b, Hans-Joachim Lehmler^{a*}

^a Department of Occupational and Environmental Health, University of Iowa, College of Public Health, Iowa City, IA 52242, USA; ^b Department of Chemical Engineering, University of Rhode Island, Kingston, RI, 02881, USA.

Figure S1. Partial phase diagrams of the DMPC-PFOA mixture obtained using TMA-DPH as fluorescence probe.

Figure S2. Partial phase diagrams of the DSPC-PFOA mixture obtained using TMA-DPH as fluorescence probe.

* Corresponding Author:
Dr. H.-J. Lehmler
The University of Iowa
Department of Occupational and Environmental Health
100 Oakdale Campus #221 IREH
Iowa City, IA 52242-5000
Phone: (319) 335-4414
Fax: (319) 335-4290
e-mail: hans-joachim-lehmler@uiowa.edu




Figure S1. Partial phase diagrams of the DMPC-PFOA mixture obtained using TMA-DPH as fluorescence probe. As indicated by the arrow, all samples were cooled at a rate of 0.2°/minute. The data points represent averages of at least three experiments \pm one standard deviation. ■ = onset of main phase transition; ▲ = maximum of the main phase transition; ◆ = offset of the main phase transition.




Figure S2. Partial phase diagrams of the DSPC-PFOA mixture obtained using TMA-DPH as fluorescence probe. As indicated by the arrow, all samples were cooled at a rate of $0.2^\circ/\text{minute}$. The data points represent averages of at least three experiments \pm one standard deviation. ■ = onset of main phase transition; ▲ = maximum of the main phase transition; ● = offset of the main phase transition.