

Supplementary Table II: Event analysis by NPC1L1 genotypes at baseline

	Genotype	History of Claudication, N(%)	Adjusted ^b OR ^a	P	History of MI, N(%)	Adjusted ^b OR ^a	P	History of Vascular disease, N(%)	Adjusted ^b OR ^a	P
-133 A>G	AA	175 (7.3)	1	0.17	329 (13.6)	1	0.45	1081 (44.8)	1	0.30
	AG + GG	191 (6.4)	0.86 (0.70-1.07)		397 (13.3)	0.94 (0.80-1.11)		1312 (44.0)	0.94 (0.83-1.06)	
-18 A>C	AA	275 (7.07)	1	0.21	542 (13.9)	1	0.13	1755 (45.1)	1	0.13
	AC + CC	91 (6.0)	0.85 (0.67-1.09)		185 (12.3)	0.87 (0.72-1.04)		641 (42.5)	0.90 (0.79-1.03)	
L272L	CC	134 (6.4)	1	0.38	458 (13.9)	1	0.26	1481 (44.9)	1	0.34
	CG + GG	232 (7.0)	0.91 (0.72-1.13)		269 (12.8)	0.91 (0.72-1.13)		916 (43.6)	0.94 (0.83-1.06)	
V1296V	CC	253 (7.4)	1	0.02	479 (14.0)	1	0.23	1536 (44.8)	1	0.54
	CT + TT	113 (5.8)	0.76 (0.61-0.97)		248 (12.6)	0.90 (0.76-1.07)		858 (43.7)	0.96 (0.85-1.09)	
U3 A>G	AA	252 (7.4)	1	0.02	475 (13.9)	1	0.25	1527 (44.7)	1	0.54
	AG + GG	113 (5.7)	0.76 (0.60-0.96)		250 (12.6)	0.90 (0.76-1.07)		865 (43.7)	0.96 (0.85-1.09)	