

Supplementary Material

Table A (Supplement to Table 3). Summary of *F* ratios from repeated measures ANOVA performed on CSD-PCA factors at selected sites after restricting participants' age to 20-40 years (40 controls, 47 patients)

	Factor (Sites)									
	442					723				
	P3 source					CP source				
	(AFz, Fz, FCz, Cz)	(POz)	(AF7/8, F7/8, F5/6, FC5/6)	(P7/8, P9/10, PO7/8)		(CP3/4, CP5/6, P3/4, P5/6, P7/8, PO7/8)	(P9/10, TP9/10)			
C	132.4 ****	36.6 ****	49.5 ****	23.5 ****		89.8 ****	72.8 ****			
G	6.00 *			11.9 ***		4.98 *	2.82			
C × G						6.35 *	15.0 ***			
T	28.8 ****		20.5 ****	23.2 ****		37.5 ****	3.55			
T × G	6.38 *	7.85 **		4.99 *						
T × C	9.03 **	16.1 ****	13.6 ***			24.8 ****	5.67 *			
T × C × G						5.63 *				
H	-	-	4.16 *	16.8 ****		52.2 ****	31.1 ****			
H × C	-	-	7.40 **	15.5 ***		63.9 ****	2.98			
H × C × G	-	-				3.87 *				
H × T	-	-					37.3 ****			
H × T × C	-	-	17.2 ****							

Note. G = Group (patients, controls); T = task (word, face); C = condition (new, old); H = hemisphere (left, right). Only *F* ratios with $p < .10$ are reported for effects pooled over gender and site (subsets as indicated; for all tabled effects, $df = 1, 83$).

- Effect not applicable.

* $p \leq .05$. ** $p \leq .01$. *** $p \leq .001$, **** $p \leq .0001$.

Table B (Supplement to Table 3). Summary of *F* ratios from repeated measures ANOVA performed on CSD-PCA factors at selected sites after excluding participants learning English as a secondary language (43 controls, 48 patients)

	Factor (Sites)									
	442					723				
	P3 source					CP source				
	(AFz, Fz, FCz, Cz)	(POz)	(AF7/8, F7/8, F5/6, FC5/6)	(P7/8, P9/10, PO7/8)	(CP3/4, CP5/6, P3/4, P5/6, P7/8, PO7/8)	(P9/10, TP9/10)				
C	143.3 ****	46.6 ****	47.7 ****	28.7 ****	85.7 ****	64.7 ****				
G	4.15 *			4.16 *	5.02 *					
C × G			4.84 *		8.05 **	14.5 ***				
T	31.0 ****		20.1 ****	22.4 ****	21.9 ****	6.24 *				
T × G	3.13	5.22 *		3.38						
T × C	13.3 ***	12.1 ***	12.0 ***		28.2 ****	6.10 *				
T × C × G					5.67 *					
H	-	-		13.5 ***	48.7 ****	21.0 ****				
H × C	-	-	6.95 **	20.8 ****	71.5 ****	7.00 **				
H × C × G	-	-			5.48 *					
H × T	-	-		2.96		32.7 ****				
H × T × C	-	-	26.4 ****							

Note. G = Group (patients, controls); T = task (word, face); C = condition (new, old); H = hemisphere (left, right). Only *F* ratios with $p < .10$ are reported for effects pooled over gender and site (subsets as indicated; for all tabled effects, $df = 1, 87$).

- Effect not applicable.

* $p \leq .05$. ** $p \leq .01$. *** $p \leq .001$, **** $p \leq .0001$.