

Supplementary materials:

Figure S1. Expression of *PTTG1* in normal fibroblasts. (A) The recombinant 6xHis-*PTTG1* protein was expressed in *E. coli* BL21/plysS cells harboring plasmid pET6H-*PTTG1* and then isolated using Ni-NTA agarose beads (Qiagen). Two µg of the purified *PTTG1* was analyzed by 10% SDS-PAGE and stained with Coomassie blue. (B) Preparation of polyclonal antibodies against *PTTG1*. The purified recombinant *PTTG1* was used to generate antibodies from rabbit. Ten µg of H1299 cell extracts were separated by 10% SDS-PAGE and then analyzed by immunoblotting using rabbit polyclonal antibodies against *PTTG1* generated from recombinant *PTTG1*. (C) IMR90 cells were infected with an adenovirus carrying the *PTTG1* gene using indicated the M.O.I. values for 2 days (upper panel). Total cell extracts were prepared, separated by 10% SDS-PAGE and then subjected to immunoblotting analysis using polyclonal antibodies against *PTTG1*. Adenovirus-carrying *PTTG1* was infected with IMR90 cells at an M.O.I. equivalent to 20. Cell extracts were prepared at the indicated time points and then analyzed (lower panel). Mock indicates experiments without virus infections.

