

Incorporation of Specific Fatty Acid Precursors During Spore Germination and Outgrowth in *Bacillus thuringiensis*

KENNETH W. NICKERSON¹ AND LEE A. BULLA, JR.^{2*}

School of Life Sciences, University of Nebraska, Lincoln, Nebraska 68588,¹ and U.S. Grain Marketing Research Laboratory, Science and Education Administration, U.S. Department of Agriculture, Manhattan, Kansas 66502²

The selective incorporation of precursors specific for individual fatty acids in germinating and outgrowing spores of *Bacillus thuringiensis* is described. The specific precursors utilized were [¹⁴C]butyrate, -isobutyrate, -valerate, and -isovalerate, which were incorporated into even-numbered normal-chain isomers, even-numbered iso-isomers, odd-numbered normal-chain acids, and odd-numbered isohomologs, respectively. This preferential incorporation by *B. thuringiensis* allows the terminal carbons of specific normal and branched-chain fatty acids, contained within the cytoplasmic membrane, to be labeled with ¹⁴C and, potentially, ¹³C.

Microbial systems have been very popular for the study of membrane structure and function (5, 6, 9). Interestingly, many bacteria have fatty acids characterized by the presence of a methyl branch at the ultimate or penultimate carbon. The gram-positive bacilli are especially prominent in this regard, and, commonly 85 to 90% of their overall fatty acids are branched (9, 10). The branched fatty acids are formed in these bacilli because branched primers are used to initiate fatty acid synthesis; the enzyme acyl coenzyme A (acyl-CoA)-acyl carrier protein transacylase has a strong preference for branched CoA esters of 4-6 carbon atoms instead of acetyl-CoA (3, 9). The methyl branches usually are derived from the three branched amino acids valine, leucine, and isoleucine via transamination to their α -keto acids and oxidative decarboxylation to the branched acyl-CoA primers (9). Consequently, there are three families of branched fatty acids, iso-even, iso-odd, and anteiso-odd, common to gram-positive bacilli, depending on the amino acid from which the branched acyl-CoA primer was derived.

A key question pertains to the physiological function of these branched fatty acids. They are essential for growth. Wegner and Foster (15) demonstrated that many rumen bacteria require isobutyric acid as a growth factor and fatty acid primer. Also, Willecke and Pardee (16) isolated a mutant of *Bacillus subtilis* with a defective α -keto acid dehydrogenase that required branched fatty acids or their precursors for growth. There was little specificity regarding which branched-chain fatty acid was required. Indeed, branched fatty acids from each of the three common classes were acceptable, as were

numerous unnatural branched substitutes. Straight-chain fatty acids definitely were not suitable. Kaneda (8, 9) has concluded that the ratios among the three classes of branched fatty acids found in the cytoplasmic membrane of gram-positive bacilli are determined by precursor availability.

A plausible hypothesis to explain the required presence of branched fatty acids is that they serve the same function as unsaturated fatty acids in organisms which do not synthesize branched fatty acids (16), i.e., they determine membrane fluidity. However, there is still only meager information available on how the branched fatty acids influence membrane properties. One of the more promising methods of defining the lipid-lipid and lipid-protein interactions in membranes is ¹³C nuclear magnetic resonance (1, 13). To accomplish such goals for the branched fatty acids requires a technique to incorporate ¹³C-enriched precursors specifically into the branched portion of the membrane fatty acids. Accordingly, we have extended our previous studies on fatty acid and membrane synthesis during spore germination and outgrowth (10, 11) to include the incorporation of four- and five-carbon ¹⁴C-precursors as primers for specific fatty acids. We used ¹⁴C, rather than ¹³C, in these studies because it is relatively inexpensive and readily available and because we wanted to determine whether germinating spores of *Bacillus thuringiensis* would selectively incorporate specific primers into individual fatty acids before we utilized the more expensive ¹³C-precursors.

Spores of *B. thuringiensis* were germinated as previously described (10, 11). Four individual pulses of 30 min were employed for each of the

specific precursors. Because the radioactive precursors were added in low concentration without accompanying carrier, the overall fatty acid composition of the cells remained unchanged. Subsequently, the fatty acids were extracted and converted to their methyl esters, and the extent of precursor incorporation into specific fatty acids was determined on a gas chromatograph interfaced to a liquid scintillation counter (10, 14). Efficiency of recovery of radioactively labeled fatty acids was about 75% using this technique.

The results obtained, although previously observed in other species, had not been obtained under conditions of differentiation before. The data in Table 1 indicate that isobutyrate is incorporated exclusively into the iso-even fatty acids, isovalerate is incorporated into the iso-odds, butyrate is incorporated into the normal-evens, and valerate is incorporated into the atypical normal-odds. As indicated by the different pulse times, incorporation of these precursors differs during germination and during outgrowth. Of course, the normal-odd fatty acids are not synthesized unless precursors such as propionate or valerate are added exogenously (9, 10). The relative proportions within each family of fatty acids, e.g., i-C₁₃, i-C₁₅, i-C₁₇, were very similar to those we observed previously (10) with radioactive acetate. Also, we very recently observed efficient incorporation of ¹⁴C-isoleucine into anteiso-isomers (data not shown).

Thus, we have demonstrated that during spore germination, the C₄ and C₅ precursors used

in these experiments are incorporated into specific fatty acids, as expected, and therefore this procedure has potential for labeling the terminal carbons of the fatty acids present in membrane lipids. Recently, Ingram et al. (7) suggested the use of exogenous [¹³C]propionate to label the terminal carbons of the fatty acids in *Escherichia coli*. Likewise, we have shown that a gram-positive bacillus such as *B. thuringiensis* can be similarly employed and we believe that the use of gram-positive bacilli should offer several significant advantages over gram-negative bacteria: (i) approximately 95% of the total fatty acids are localized in the cytoplasmic membrane of gram-positive bacilli (4); (ii) it is easier to form protoplasts; (iii) propionate added to *E. coli* competes with acetate to function as a primer (2, 7) and, consequently, is diluted in the process, whereas the addition of a ¹⁴C-branched amino acid or [¹⁴C]propionate to *B. thuringiensis* can result in the almost exclusive production of a single branched fatty acid (L. A. Bulla, Jr., unpublished data); (iv) also, because the gram-positive bacilli do not utilize acetyl-CoA as a primer for normal-fatty acids (9), the addition of exogenous butyrate or valerate leads to the undiluted formation of the normal fatty acids; (v) because the gram-positive bacilli are relatively unspecific in their branched fatty acid requirements (12, 16), a great many atypical branched primers also can be employed, e.g. 2-methyl valerate, 2-ethyl butyrate, or trimethyl acetate (8, 16); and (vi) the membrane interactions operative during vegetative growth of gram-posi-

TABLE 1. Specific incorporation of radioactive precursors into individual fatty acids during germination and outgrowth of *B. thuringiensis*^a

Pre-cursor	Pulse time (min)	¹⁴ C-precursor incorporation (cpm) ^b												
		n-C ₁₁	i-C ₁₂	n-C ₁₂	i-C ₁₃	n-C ₁₃	i-C ₁₄	n-C ₁₄	i-C ₁₅	n-C ₁₅	i-C ₁₆	n-C ₁₆	i-C ₁₇	n-C ₁₇
Butyrate	0-30			1,800				27,000				55,000		
	30-60			2,000				67,000				137,000		
	60-90			12,000				125,000				179,000		
	90-120			25,000				221,000				365,000		
Isobutyrate	0-30		1,500				23,000				25,000			
	30-60		23,000				87,000				95,000			
	60-90		35,000				138,000				153,000			
	90-120		7,200				214,000				432,000			
Valerate	0-30					18,000				82,000				
	30-60	1,100				27,000				191,000				400
	60-90	3,100				45,000				317,000				900
	90-120	5,000				76,000				482,000				1,200
Isovalerate	0-30				35,000					30,000			23,000	
	30-60				65,000					127,000			23,000	
	60-90				76,000					176,000			37,000	
	90-120				131,000					316,000			75,000	

^a Sixty milligrams of spores per 23 ml.

^b Sodium [1-¹⁴C]butyrate, sodium [1-¹⁴C]isobutyrate, and sodium [1-¹⁴C]valerate = 25 μCi/23 ml; sodium [1-¹⁴C]isovalerate = 10 μCi/23 ml.

tive bacilli can be contrasted to those participating in the developmental phenomena of sporulation and spore germination.

LITERATURE CITED

- Birdsall, N. J. M., D. J. Ellar, A. G. Lee, J. C. Metcalfe, and G. B. Warren. 1975. ^{13}C -enriched phosphatidyl ethanolamines from *Escherichia coli*. *Biochim. Biophys. Acta* **380**:344-354.
- Bloch, K. 1975. Fatty acid synthase from *Mycobacterium phlei*. *Methods Enzymol.* **35**:84-90.
- Butterworth, P. H. W., and K. Bloch. 1970. Comparative aspects of fatty acid synthesis in *Bacillus subtilis* and *Escherichia coli*. *Eur. J. Biochem.* **12**:496-501.
- Daniels, M. J. 1969. Lipid synthesis in relation to the cell cycle of *Bacillus megaterium* KM and *Escherichia coli*. *Biochem. J.* **115**:697-701.
- Daron, H. H. 1973. Nutritional alteration of the fatty acid composition of a thermophilic *Bacillus* species. *J. Bacteriol.* **116**:1096-1099.
- Dawes, I. W., and H. O. Halvorson. 1972. Membrane synthesis during outgrowth of bacterial spores, p. 449-455. In H. O. Halvorson, R. Hanson, and L. L. Campbell (ed.), *Spores V*. American Society for Microbiology, Washington, D.C.
- Ingram, L. O., L. S. Chevalier, E. J. Gabbay, K. D. Ley, and K. Winters. 1977. Propionate-induced synthesis of odd-chain-length fatty acids by *Escherichia coli*. *J. Bacteriol.* **131**:1023-1025.
- Kaneda, T. 1963. Biosynthesis of branched chain fatty acids. II. Microbial synthesis of branched long chain fatty acids from certain short chain fatty acid substrates. *J. Biol. Chem.* **238**:1229-1235.
- Kaneda, T. 1977. Fatty acids of the genus *Bacillus*: an example of branched-chain preference. *Bacteriol. Rev.* **41**:391-418.
- Nickerson, K. W., L. A. Bulla, and T. L. Mounts. 1975. Lipid metabolism during bacterial growth, sporulation, and germination: differential synthesis of individual branched- and normal-chain fatty acids during spore germination and outgrowth of *Bacillus thuringiensis*. *J. Bacteriol.* **124**:1256-1262.
- Nickerson, K. W., J. DePinto, and L. A. Bulla. 1975. Lipid metabolism during bacterial growth, sporulation, and germination: kinetics of fatty acid and macromolecular synthesis during spore germination and outgrowth of *Bacillus thuringiensis*. *J. Bacteriol.* **121**:227-233.
- Scandella, C. J., and A. Kornberg. 1969. Biochemical studies in bacterial sporulation and germination. XV. Fatty acids in growth, sporulation, and germination of *Bacillus megaterium*. *J. Bacteriol.* **98**:82-86.
- Stoffel, W., K. Bister, C. Schreiber, and B. Tunggal. 1976. ^{13}C -NMR studies of membrane structure of enveloped virions (vesicular stomatitis virus). *Hoppe-Seyler's Z. Physiol. Chem.* **357**:905-915.
- Thomas, P. J., and H. J. Dutton. 1969. Improved resolution in high-sensitivity dual-label gas radiochromatography. *Anal. Chem.* **41**:657-660.
- Wegner, G. H., and E. M. Foster. 1963. Incorporation of isobutyrate and valerate into cellular plasmalogen by *Bacteroides succinogenes*. *J. Bacteriol.* **85**:53-61.
- Willecke, K., and A. B. Pardee. 1971. Fatty acid-requiring mutant of *Bacillus subtilis* defective in branched chain α -keto acid dehydrogenase. *J. Biol. Chem.* **246**:5264-5272.