

Supplementary Information

For

Enzymatic Route to Preparative-scale Synthesis of UDP-GlcNAc/GalNAc, their analogues and GDP-fucose

Guohui Zhao^{a,c}, Wanyi Guan^{b,a}, Li Cai^a & Peng George Wang^{a*}


a Departments of Chemistry and Biochemistry, The Ohio State University, 484 W. 12th Ave., Columbus,
OH 43210, USA.

b National Glycoengineering Research Center and The State Key Laboratory of Microbial Technology,
Shandong University, Jinan, Shandong 250100, People's Republic of China.


c College of Pharmacy, Nankai University, Tianjin 300071, China

Correspondence should be addressed to P.G.W. (E-mail:wang.892@osu.edu)


NMR, MS data and spectra for all new compounds




2-benzamido-2-deoxy- α -D-glucosyl diammonium Phosphate: ^1H NMR (400 MHz, D_2O) δ 7.88-7.90 (m, 2H), 7.56-7.69 (m, 3H), 5.58 (dd, $J = 7.4, 3.3$ Hz, 1H), 4.23-4.26 (m, 1H), 3.94-4.06 (m, 3H), 3.86 (dd, $J = 12.3, 4.7$ Hz, 1H), 3.63 (app t, $J = 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, D_2O) δ 171.6, 133.6, 132.2, 128.7, 127.5, 93.2 (d), 72.5, 71.1, 70.0, 60.6, 54.6 (d); ^{31}P NMR (162 MHz, D_2O) δ 0.71; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_9\text{P} (\text{M}-\text{H})^-$ 362.0646, found 362.0643 m/z .




2-Acetamido-6-azido-2,6-dideoxy- α -D-glucosyl diammonium Phosphate: ^1H NMR (400 MHz, D_2O) δ 5.42 (dd, $J = 7.2, 3.3$ Hz, 1H), 4.05-4.09 (m, 1H), 3.97-4.01 (m, 1H), 3.76-3.85 (m, 2H), 3.67 (dd, $J = 13.6, 4.2$ Hz, 1H), 3.59 (app t, $J = 9.3$ Hz, 1H), 2.10 (s, 3H); ^{13}C NMR (100 MHz, D_2O) δ 174.7, 93.1 (d), 71.0, 70.9, 70.4, 54.0 (d), 50.8, 22.0; ^{31}P NMR (162 MHz, D_2O) δ 0.60; HRMS (ESI) calcd for $\text{C}_8\text{H}_{14}\text{N}_4\text{O}_8\text{P} (\text{M}-\text{H})^-$ 325.0555, found 325.0559 m/z .




2-Acetamido-2-deoxy- α -D-allosyl diammonium Phosphate: ^1H NMR (400 MHz, D_2O) δ 5.39 (dd, $J = 7.7, 3.6$ Hz, 1H), 4.02-4.13 (m, 3H), 3.94 (dd, $J = 12.3, 2.3$ Hz, 1H), 3.79 (dd, $J = 12.3, 5.3$ Hz, 1H), 3.70 (dd, $J = 10.5, 3.4$ Hz, 1H), 2.09 (s, 3H); ^{13}C NMR (100 MHz, D_2O) δ 174.1, 92.3 (d), 69.9, 67.3, 66.1, 60.9, 50.3 (d), 22.0; ^{31}P NMR (162 MHz, D_2O) δ 2.17; HRMS (ESI) calcd for $\text{C}_8\text{H}_{15}\text{NO}_9\text{P} (\text{M}-\text{H})^-$ 300.0490, found 300.0481 m/z .




2-benzamido-2-deoxy- α -D-galactopyranosyl diammonium Phosphate: ^1H NMR (400 MHz, D_2O): δ = 7.83-7.86 (m, 2H), 7.61-7.66 (m, 1H), 7.52-7.56 (m, 2H), 5.59 (dd, $J = 7.5, 3.5$ Hz, 1H), 4.45-4.49 (m, 1H), 4.23-4.26 (m, 1H), 4.09-4.15 (m, 2H), 3.77-3.83 (m, 2H); ^{13}C NMR (125 MHz, D_2O): δ = 171.8, 133.6, 132.2, 128.7, 127.5, 93.6 (d), 71.7, 68.7, 67.7, 61.3, 50.7 (d); ^{31}P NMR (162 MHz, D_2O): δ = -0.10; HRMS (ESI) calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_9\text{P} (\text{M}-\text{H})^-$ 362.0646, found 362.0660 m/z .




2-Azidoacetamido-2-deoxy- α -D-galactopyranosyl diammonium Phosphate: ¹H NMR (400 MHz, D₂O): δ = 5.30 (dd, J = 7.3, 3.3 Hz, 1H), 4.06-4.10 (m, 1H), 3.86-4.00 (m, 4H), 3.81 (dd, J = 11.1, 3.0 Hz, 1H), 3.53-3.61 (m, 2H); ¹³C NMR (100 MHz, D₂O): δ = 171.1, 93.4 (d), 71.7, 68.4, 67.3, 61.1, 51.7, 50.1 (d); ³¹P NMR (162 MHz, D₂O): δ = -0.74; HRMS (ESI) calcd for C₈H₁₄N₄O₉P (M-H)⁻ 341.0504, found 341.0498 m/z.




2-Acetamido-6-azido-2,6-dideoxy- α -D-galactopyranosyl diammonium Phosphate: ¹H NMR (400 MHz, D₂O): δ = 5.42 (dd, J = 7.3, 3.2 Hz, 1H), 4.25 (app t, J = 6.6, 1H), 4.18-4.21 (m, 1H), 3.95-4.03 (m, 2H), 3.58-3.63 (m, 1H), 3.50-3.54 (m, 1H), 2.07 (s, 3H); ¹³C NMR (100 MHz, D₂O): δ = 174.9, 93.3 (d), 69.5, 68.6, 67.8, 50.5, 50.0 (d), 22.1; ³¹P NMR (162 MHz, D₂O): δ = 0.58; HRMS (ESI) calcd for C₈H₁₄N₄O₈P (M-H)⁻ 325.0555, found 325.0566 m/z.



2-Acetamido-4-azido-2,4-dideoxy- α -D-galactopyranosyl diammonium Phosphate: ¹H NMR (400 MHz, D₂O): δ = 5.43 (dd, J = 7.4, 3.0 Hz, 1H), 4.22-4.27 (m, 3H), 4.15-4.16 (m, 1H), 3.75-3.78 (m, 2H), 2.06 (s, 3H); ¹³C NMR (100 MHz, D₂O): δ = 174.8, 93.4 (d), 70.1, 67.8, 62.8, 61.1, 50.2 (d), 22.0; ³¹P NMR (162 MHz, D₂O): δ = -0.95; HRMS (ESI) calcd for C₈H₁₄N₄O₈P (M-H)⁻ 325.0555, found 325.0542 m/z.




Uridine 5'-diphospho-2-acetamido-6-azido-2,6-dideoxy- α -D-glucopyranose diammonium salt: ¹H NMR (400 MHz, D₂O): δ = 7.98 (d, J = 8.1 Hz, 1H), 5.97-5.99 (m, 2H), 5.51 (dd, J = 7.2, 3.3 Hz, 1H), 4.36-4.40 (m, 2H), 4.20-4.30 (m, 3H), 4.00-4.09 (m, 2H), 3.81 (t, J = 9.2 Hz, 1H), 3.74 (dd, J = 13.5, 2.5 Hz, 1H), 3.57-3.65 (m, 2H), 2.08 (s, 3H); ³¹P NMR (162 MHz, D₂O): δ = -11.3 (d, J = 21.4 Hz), -13.2 (d, J = 21.4 Hz); HRMS (ESI) calcd for C₁₇H₂₅N₆O₁₆P₂ (M-H)⁻ 631.0808, found 631.0806 m/z.




Uridine 5'-diphospho-2-acetamido-2-deoxy- α -D-allopyranose diammonium salt: ^1H NMR (400 MHz, D_2O): $\delta = 7.96$ (d, $J = 8.1$ Hz, 1H), 5.96-6.00 (m, 2H), 5.53 (dd, $J = 7.1, 3.8$ Hz, 1H), 4.36-4.40 (m, 2H), 4.21-4.30 (m, 3H), 4.08-4.15 (m, 3H), 3.92 (dd, $J = 12.6, 2.3$ Hz, 1H), 3.82 (dd, $J = 12.6, 4.5$ Hz, 1H), 3.77 (dd, $J = 10.5, 3.2$ Hz, 1H), 2.10 (s, 3H); ^{31}P NMR (162


MHz, D_2O): $\delta = -11.0$ (d, $J = 19.9$ Hz), -13.0 (d, $J = 21.7$ Hz); HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{26}\text{N}_3\text{O}_{17}\text{P}_2 (\text{M}-\text{H})^-$ 606.0743, found 606.0730 m/z .





Uridine 5'-diphospho-2-acetamido-6-azido-2,6-dideoxy- α -D-galactopyranose diammonium salt: ^1H NMR (400 MHz, D_2O): $\delta = 7.98$ (d, $J = 8.2$ Hz, 1H), 5.97-5.99 (m, 2H), 5.54 (dd, $J = 7.2, 3.4$ Hz, 1H), 4.35-4.39 (m, 2H), 4.19-4.30 (m, 5H), 4.02 (m, 1H), 3.98 (dd, $J = 10.8, 3.2$ Hz, 1H), 3.60 (dd, $J = 12.8, 7.2$ Hz, 1H), 3.50 (dd, $J = 12.8, 6.0$ Hz, 1H), 2.09 (s, 3H); ^{31}P


NMR (162 MHz, D_2O): $\delta = -11.2$ (d, $J = 21.5$ Hz), -13.0 (d, $J = 21.5$ Hz); HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{25}\text{N}_6\text{O}_{16}\text{P}_2 (\text{M}-\text{H})^-$ 631.0808, found 631.0801 m/z .








0.7087








0.0009







— 174.0574

92.3157
92.2636

69.8925
67.3361
66.0626
60.9207



50.3316
50.2666


21.9680




S5

2.1664








S5



-0.0965



S5





171.0594

83.4701
83.4125

71.7064
68.4417
67.3160


61.1472

51.7004
50.1121
50.0332




S5

-0.7352



S5



S5



— 174.8994

— 93.3398
93.2839

— 69.5111
68.6074
67.7520



— 50.5309
50.0567
49.9813

— 22.1210




S5


0.5813



S5





S5




S5


-0.9491




S5




S5




S5





S5




S5

 -11.2077
-11.3393

 -13.1037
-13.2364





S5




S5

-10.9861
-11.1089

-12.9464
-13.0802








S5

-11.1203
-11.2519


-12.9302
-13.0615



S5



Note: NMR regions (red box) showing typical common impurities after the first time silica gel purification of sugar-1-P, second silica gel chromatography is needed to remove the impurities for NMR purposes.





S5

C 8 H 14 N 4 O 8 P 1 ,325.05


$\times 10^4$
HRMS (negative mode)

325.0555




S5


HRMS (negative mode)




S5




S5




S5




S5




S5



S5



S5



S5