

Table S3. Summary of reported enzyme kinetic parameters of histidine kinases.

Enzyme	Autophosphorylation Activity			
	k_{cat} (s ⁻¹)	K_M ATP (μM)	k_{cat}/K_M (M ⁻¹ s ⁻¹)	References
DesKc	2.8×10^{-3}	64	44	This work
CheA	0.117	274	427	[1]
CheA	0.026	300	87	[2]
NRII	0.032	18	1805	[3]
NarX	5×10^{-5}	2.4	21	[4]
NarQ	2.2×10^{-4}	22.8	9.65	
KinA	1.9×10^{-3}	74	25.7	[5]
EnvZ115	8.1×10^{-5}	218	0.37	[6]
PhoQc	-	20.1	-	[7]
WalK(Spn) (VicK)	1.7×10^{-3}	37	46.0	[8]

1. Surette, M.G., et al., *Dimerization is required for the activity of the protein histidine kinase CheA that mediates signal transduction in bacterial chemotaxis*. J Biol Chem, 1996. **271**(2): p. 939-45.
2. Tawa, P. and R.C. Stewart, *Mutational activation of CheA, the protein kinase in the chemotaxis system of Escherichia coli*. J Bacteriol, 1994. **176**(14): p. 4210-8.
3. Jiang, P. and A.J. Ninfa, *Regulation of autophosphorylation of Escherichia coli nitrogen regulator II by the PII signal transduction protein*. J Bacteriol, 1999. **181**(6): p. 1906-11.
4. Noriega, C.E., et al., *Autophosphorylation and dephosphorylation by soluble forms of the nitrate-responsive sensors NarX and NarQ from Escherichia coli K-12*. J Bacteriol, 2008. **190**(11): p. 3869-76.
5. Grimshaw, C.E., et al., *Synergistic kinetic interactions between components of the phosphorelay controlling sporulation in Bacillus subtilis*. Biochemistry, 1998. **37**(5): p. 1365-75.
6. Kenney, L.J., *Kinase activity of EnvZ, an osmoregulatory signal transducing protein of Escherichia coli*. Arch Biochem Biophys, 1997. **346**(2): p. 303-11.
7. Marina, A., et al., *Structural and mutational analysis of the PhoQ histidine kinase catalytic domain. Insight into the reaction mechanism*. J Biol Chem, 2001. **276**(44): p. 41182-90.
8. Gutu, A.D., et al., *Kinetic characterization of the WalRKSpn (VicRK) two-component system of Streptococcus pneumoniae: dependence of WalKSpn (VicK) phosphatase activity on its PAS domain*. J Bacteriol, 2010. **192**(9): p. 2346-58.