

Table S9. Primers used for PCR amplification

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
<i>Gapdh</i>	CAAGAACGCTGACATCCTGA	CAGAAGTCACAAACGCCTCA
<i>Act5C</i>	GTGCCCATCTACGAGGGTTA	ACATCTGCTGGAAGGTGGAC
<i>Gβ76C</i>	TATCGCTGGCTTATCGCTTT	ACAGGCTGATGCAAGTGATG
<i>CG8889</i>	CTACATGGCCCAGTCCAGTT	AAGTCGCTGATGATGGCTCT
<i>TotB</i>	CGCACTGCTACTGATTGGAA	TGCACCGAGTTCGGTAAAG
<i>ninaC</i>	CTCGCTCCAGGACTTTGTC	TTTGCTGATAGTTGCGTTGC
<i>CG5172</i>	GTGGTGGTGGTGGTCAGAG	CAAGGACTTGCTTGCATGG
<i>Arr2</i>	CCAAGAAGTCCGTGAAGAGC	GGCGAAGTCCTCGAATACAA
<i>Osi18</i>	GTCGAATCCGCTTACTCAGG	TAACCAAGGCTCCAGACAGC
<i>CG31313</i>	GCAGTCCGTGAAAGTGATCTT	GCCTGTCCCAGATCTTTACG
<i>nimB2</i>	AGCGAAATCCGCACATCTAC	CCTGCAGATGCAGAAGTTTG
<i>PIP82</i>	TCGGATTGGTTCCTCTATGG	GCACCAACGACTGGAATTTT
<i>trp</i>	GGAGGAAGATCATCGAGCAG	CTTGCTAGCTTCGGTCTTGG
<i>CG10953</i>	ATTCCTGGTTGTTGCCTTTG	GGGGGAATGTAGGTGTTCACT
<i>Rh6</i>	CAAGGACTGGTGGAACAGGT	CATGGCATCTTCTCCCGTAG
<i>Rh2</i>	TGAGGATTGCGACAAAAGTG	CCGTATTGCCAAACACACAC
<i>Rh3</i>	TCTGACATCCTGCACCTTTG	GTGGTCTGCTGTGGTTCCTG
<i>Rh4</i>	GGAACCCCAATTGTCGTCTA	CCCCGATCAGCGACATTAC
<i>Rh5</i>	GAAATGGACTGGTCATCTGGA	GAGACGAAGGGCGTGATTAG
<i>Rh1</i>	CTGGGCAAGATTGCCTACAT	CCCCAAATGGTATTCAAGTGG