

Interleukin 1 α Causes Rapid Activation of Cytosolic Phospholipase A₂ by Phosphorylation in Rat Mesangial Cells

Joseph Gronich, Martha Konieczkowski, Michael H. Gelb,* Raphael A. Nemenoff,† and John R. Sedor
Departments of Medicine, and Physiology and Biophysics, School of Medicine, Case Western Reserve University, Cleveland, Ohio 44109; *Departments of Chemistry and Biochemistry, University of Washington, Seattle, Washington 98195; and ‡Department of Medicine, University of Colorado Health Sciences Center, Denver, Colorado 80262

Abstract

We have shown previously that interleukin 1 (IL-1) stimulates eicosanoid production in glomerular mesangial cells (MC) by de novo synthesis of a 14-kD, group II phospholipase A₂ (PLA₂). IL-1-stimulated prostaglandin E₂ synthesis precedes expression of this enzyme, suggesting that another PLA₂ isoform must be more rapidly activated. In the presence but not absence of calcium ionophore, [³H]arachidonate release is increased significantly as early as 5 min after addition of IL-1, and IL-1 concurrently stimulates a Ca²⁺-dependent phospholipase activity, which was characterized as the cytosolic form of PLA₂ (cPLA₂). IL-1 does not alter either cPLA₂ mRNA expression or mass in serum-stimulated MC, suggesting that cPLA₂ activity is increased by a posttranslational modification. IL-1 treatment for 30 min doubles ³²P incorporation into immunoprecipitable cPLA₂ protein, concordant with the increase in enzyme activity. Immunoblot analysis of extracts derived from IL-1-treated (30 min) cells demonstrates a decreased mobility of cPLA₂, and treatment of MC lysates with acid phosphatase significantly reduces cytokine-activated cPLA₂ activity, further indicating that IL-1 stimulates phosphorylation of the enzyme. IL-1 treatment (24 h) of serum-deprived MC doubled cPLA₂ mRNA, protein, and activity. In summary, IL-1 increases cPLA₂ activity in a biphasic, time-dependent manner both by posttranslational modification and de novo synthesis. We consider cPLA₂ activation a key step in IL-1-stimulated synthesis of pro-inflammatory, lipid mediators, and an integral event in the phenotypic responses induced in target cells by this cytokine. (*J. Clin. Invest.* 1994. 93:1224–1233.) Key words: arachidonic acid • cellular signaling • cytokine • eicosanoid metabolism • inflammation

Introduction

IL-1 is a potent pro-inflammatory cytokine that activates functional responses and biochemical processes in target cells (1). The release of arachidonic acid-derived lipid mediators is a

This work was presented in part at a FASEB Summer Conference, Renal Hemodynamics: Epithelial and Endothelial Interactions, (Saxton River, VT, June 1992) and the Annual Meeting of the American Society of Nephrology, Baltimore, MD, November 1992.

Address correspondence to Dr. John R. Sedor, Department of Medicine, MetroHealth Medical Center, 2500 MetroHealth Drive/Office A1112, Cleveland, OH 44109-1998.

Received for publication 2 February 1993 and in revised form 12 October 1993.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/94/03/1224/10 \$2.00

Volume 93, March 1994, 1224–1233

characteristic of chronic and acute inflammation, and IL-1 stimulates eicosanoid synthesis in vitro (1). The molecular mechanisms of IL-1-stimulated arachidonate metabolism have not been defined completely, but receptor-activated, phospholipase A₂ (PLA₂)¹-mediated release of arachidonic acid from the sn-2 position of membrane phospholipids is rate-limiting in eicosanoid biosynthesis (2).

Mammalian PLA₂s, until recently, have been classified into two groups (I and II) based on position of cysteine pairs and sequence similarities to families of 14-kD PLA₂ isozymes secreted in snake venoms (3). Mammalian pancreatic PLA₂ is a group I enzyme. Mammalian group II enzymes are synthesized in many tissues and abundantly secreted into inflammatory exudates. Both groups I and II secretory PLA₂'s (sPLA₂) require millimolar calcium concentrations for half-maximal enzyme activity and do not demonstrate preference for the fatty acid present in the sn-2 position of phospholipids. Recently a 97-kD cytosolic form of PLA₂ (cPLA₂) has been identified in glomerular mesangial cells and the human monocyte cell line U937 (4–6). This enzyme translocates to membrane vesicles in response to nanomolar changes in free Ca²⁺ concentration and selectively hydrolyzes arachidonyl-phospholipids in membrane vesicles (7). The primary structure of cPLA₂ lacks any identity with either cloned sPLA₂, and its biochemical characteristics suggest that this form of PLA₂ regulates intracellular arachidonic acid release (4–7).

Enhanced prostaglandin synthesis characterizes glomerular inflammation, and recent evidence suggests that IL-1 is important in the pathogenesis of glomerular injury (8, 9). Mesangial cells are the most abundant source of prostaglandins within the glomerulus, and in culture, express a phenotype that mimics their characteristics within an inflamed glomerulus. We believe IL-1-stimulated mesangial cells to be an appropriate in vitro model system to define the molecular basis for the enhanced arachidonate availability that characterizes glomerular injury.

We have demonstrated previously that IL-1 specifically stimulates mesangial cells to synthesize a group II (nonpancreatic type) sPLA₂ (10). IL-1-dependent induction of type II sPLA₂ mRNA expression increased coordinately with PLA₂ activity and PGE₂ synthesis. However, three observations reported in this study suggested that a PLA₂ distinct from this sPLA₂ isoform also must be activated by IL-1 stimulation. First, PGE₂ synthesis occurred before 6–10 h of IL-1 stimulation, when sPLA₂ mRNA is first detected. Second, IL-1 did not

1. Abbreviations used in this paper: AA-COCF₃, the trifluoromethyl ketone analogue of arachidonic acid which inhibits cPLA₂ activity; AA-COCH₃, an arachidonate analogue devoid of inhibitory activity for cPLA₂; MAP, mitogen-activated protein; PLA₂, phospholipase A₂; similarly, cPLA₂ and sPLA₂, cytosolic form of and 14-kD secretory PLA₂, respectively; PLAP, PLA₂-activating protein.

increase basal [^3H]arachidonic acid release but did not potentiate vasopressin-stimulated arachidonate release. Because cPLA₂ has been shown to be regulated by vasopressin (4), we reasoned that this PLA₂ may be primed by IL-1 for enhanced, agonist-stimulated arachidonic acid release. Third, although phosphatidylcholine is hydrolyzed poorly by group II sPLA₂, IL-1 increased hydrolysis of arachidonic acid from phosphatidylcholine, a finding consistent with enhanced cPLA₂ activity. Taken together, these data suggest that IL-1 directly or indirectly activates cPLA₂. We now report that IL-1 rapidly increases cPLA₂ activity by changing the phosphorylation state of the enzyme, and in longer incubations, can increase catalytic activity by inducing cPLA₂. cPLA₂ activation and enhanced arachidonate availability is integral to the glomerular synthesis of pro-inflammatory eicosanoids stimulated by IL-1.

Methods

Materials. Recombinant human IL-1 α (IL-1, 3×10^8 U/mg) was kindly provided by Dr. Peter Lomedico (Hoffmann-LaRoche, Nutley NJ). The endotoxin content and specific activity have been previously published (10). L- α -1-palmitoyl, 2-arachidonyl, [1- ^{14}C]arachidonyl-phosphatidylethanolamine (60 Ci/mmol), and arachidonic acid (5, 6, 8, 9, 11, 12, 14, 15, ^3H (N), 100 Ci/mmol) were purchased from DuPont New England Nuclear (Boston, MA). L- α -1-stearoyl, 2-arachidonyl, [1- ^{14}C]arachidonyl-phosphatidylcholine (56 Ci/mmol) was purchased from Amersham Corp. (Arlington Heights, IL). ^{32}P -orthophosphate (carrier free), *trans* ^{35}S -label (L-methionine, [^{35}S], 1,163 Ci/mmol), phosphate-free RPMI 1640 and methionine, cysteine-free RPMI 1640 medium were obtained from ICN Biomedicals, Inc. (Irvine, CA).

Cell culture. Well-characterized mesangial cells were grown from collagenase-treated rat glomeruli in RPMI 1640 medium containing 8.5% FBS and 8.5% calf serum, 15 mM Hepes, 100 U/ml penicillin, 100 $\mu\text{g}/\text{ml}$ streptomycin and fungizone, 5 $\mu\text{g}/\text{ml}$ of insulin and transferrin, and 5 ng/ml of sodium selenite as previously described (10).

The cells were used in passages 3–8, and as indicated in the specific methods or in the results were either maintained in 17% serum, acutely serum-deprived (2 h) or were held for 24–48 h in 0.5% FBS.

[^3H]arachidonate release. Basal and stimulated arachidonate release was determined as previously described (10), with some modifications. Cells in six-well clusters were labeled for 20 h with 0.5 $\mu\text{Ci}/\text{ml}$ [^3H]arachidonic acid in RPMI 1640 medium with 17% serum. The labeling media was aspirated and the cells washed and serum-deprived for 2 h at 37°C in Hepes (10 mM)-buffered RPMI (Hepes-RPMI) containing 0.1% BSA. The experiment was initiated by the addition of vehicle or IL-1 for 1, 4, or 20 min before the addition of the Ca²⁺ ionophore, A23187 (0.25 μM) (Calbiochem Corp., San Diego, CA). This A23187 concentration was chosen to cause only a minimal change in [^3H]arachidonate release (Fig. 1) and to mimic a receptor-mediated increase in cytosolic Ca²⁺ (M. B. Ganz, personal communication). Ionophore was added for an additional 1 min after the cells were incubated with IL-1 for 1 and 4 min, and for 10 min after the cells had been exposed to IL-1 for 20 min for total incubation times of 2, 5, and 30 min, respectively. As positive controls, cells were also incubated with PMA for 10 min followed by an additional 10 min in the presence of ionophore. In some experiments, the cells were preincubated for 2 h with a control or cPLA₂-inhibitory arachidonate acid analogues before stimulation with IL-1 or PMA (11). All reactions were terminated by removal of the incubation media and the addition of 0.4 ml of cold (4°C) methanol containing 10 μg unlabeled arachidonic acid to the monolayer. The cells were scraped, and this suspension along with an additional 0.4 ml methanol rinse was combined with the incubation supernatants and acidified with formic acid (0.2%, vol/vol). Chloroform/methanol (1:1.2, vol/vol, 0.6 ml) was added to the extraction tube followed by the addition of 0.6 ml of chloroform. After separating the organic and aqueous phases by centrifugation, the organic phase was dried under N₂ and resolubilized in ethanol. Free arachidonate was separated from cellular phospholipids using TLC. [^3H]arachidonic acid release was expressed as a percentage of total [^3H]arachidonate incorporated into the mesangial cell monolayer (determined by total TLC plate radioactivity).

Cell-free extracts and PLA₂ assay. At the time of the experiment, subconfluent cells were serum-restricted by replacing growth medium




Figure 1. IL-1 rapidly stimulates [^3H]arachidonate release. Subconfluent rat mesangial cells were labeled with [^3H]arachidonic acid (0.5 $\mu\text{Ci}/\text{well}$) for 20 h, washed with RPMI media containing 0.1% BSA, and incubated in this media for an additional 2 h. The cells were then stimulated for the indicated times with vehicle alone (control), IL-1 (50 ng/ml), or PMA (300 nM). Cells were exposed to IL-1 for 1 and 4 min before the addition of A23187 (0.25 μM) for an additional 1 min. In the longest incubations, mesangial cells were incubated for 10 and 20 min with PMA and IL-1, respectively. A23187 was subsequently added for 10 min. The times in the figure indicate the total exposure time to IL-1 or PMA in the absence and the presence of ionophore. [^3H]arachidonate in cells and supernatants was quantified by TLC after chloroform-methanol extraction. Results are the mean \pm SEM for triplicate

experiments of triplicate wells (n = number of separate experiments; * P < 0.05, ** P < 0.025 compared to the control value). In the absence and presence of IL-1, ionophore-stimulated [^3H]arachidonate release at 2 min was $4.4 \pm 0.6\%$ and $5.1 \pm 0.8\%$ release (n = 4, ionophore- vs. ionophore- and IL-1-stimulated, respectively, P = NS); at 5 min was $4.5 \pm 0.4\%$ vs. $6.9 \pm 0.9\%$ release (n = 3, ionophore- vs. ionophore- and IL-1-stimulated, respectively, P < 0.01); at 30 min was $5.3 \pm 0.2\%$ and $8.8 \pm 0.7\%$ release (n = 3, ionophore- vs. ionophore- and IL-1-stimulated, respectively, P < 0.01). After 10 min, ionophore-stimulated [^3H]arachidonate release was $5.3 \pm 0.2\%$ vs. $11.0 \pm 2.9\%$ in the absence and presence of PMA, respectively (n = 3, P < 0.005).

with Hepes-RPMI for 2 h before incubation with IL-1, 12-tetradecanoyl phorbol myristate 13-acetate (Calbiochem Corp.) or vehicle-control at the indicated doses. These stimuli were added directly to the preincubation medium, and the cells were incubated at 37°C in the presence of 5% CO₂, 95% air for the indicated times. In some experiments, cells were treated with heated IL-1 (90°C, 60 min) or native IL-1 in the presence of polymyxin B (10 µg/ml, Sigma Chemical Co., St. Louis, MO) to determine if LPS contaminating the reagents accounted for the effects of the cytokine. Treatment of cells with polymyxin B alone did not alter cPLA₂ activity in cell-free extracts (not shown). Three 100-cm dishes of cells were used for each condition to assay PLA₂ activity in 200,000-g supernatants of cell lysates as previously described (4, 10). Briefly after the incubation period, the media was removed. The cells were washed and scraped at 4°C into a buffer containing 250 mM sucrose, 50 mM Hepes (pH 7.5), 1 mM EDTA, 1 mM EGTA, and the protease inhibitors, pepstatin A (1 µg/ml), leupeptin (1 µg/ml), and PMSF (0.1 mM). The cells were homogenized using a Dounce homogenizer (25 strokes) and spun at 500 g for 5 min to remove the nuclei. The homogenate was centrifuged at 200,000 g for 1 h at 4°C using a model Ti-70.1 rotor (Beckman Instruments, Inc., Fullerton, CA). Protein content in the supernatant was determined (protein assay, Bio-Rad Laboratories, Richmond, CA). Samples from each condition were matched for protein concentration by appropriate dilution and assayed for PLA₂ activity. Unless otherwise indicated, [¹⁴C]arachidonyl-phosphatidylcholine was used as substrate (final concentration, 15 µM). The assay was initiated by adding mesangial cell extract containing 1.5–7 µg of protein to an assay buffer containing 5 mM CaCl₂ and 50 mM Hepes, pH 7.5 (final concentrations). In some experiments, control and cPLA₂-inhibitory arachidonic acid analogues (11), in the indicated concentrations, were added for 5 min at 37°C to the cellular lysates in assay buffer. The assay then was initiated by the addition of [¹⁴C]phosphatidylcholine. Reactions proceeded for 30 min at 37°C and were terminated by addition of 40 µl of 2% acetic acid in ethanol containing 10 µg of unlabeled arachidonic acid. The reaction mixtures and appropriate standards were spotted onto heat-activated LK 5 DF silica gel TLC plates (Whatman Inc., Clifton, NJ) and developed with ethyl acetate/iso-octane/acetic acid/water (55:75:8:100). Lipids were visualized by I₂ staining. Free arachidonate was scraped and quantified by liquid scintillation counting. Results were expressed as picomoles of arachidonate released per minute per milligram of protein.

Extract fractionation. As described (4, 12), protein-matched extracts from control, IL-1- and PMA-stimulated cells were applied to a prepacked 5 × 50-mm Mono-Q HR 5/5 anion exchange column (Pharmacia LKB Biotechnology Inc., Piscataway, NJ) preequilibrated with 150 mM NaCl, 1 mM EGTA, 1 mM EDTA, and 50 mM Hepes, pH 7.5. The column was washed with equilibration buffer. PLA₂ activity was eluted with a linear salt gradient (0.15–1.0 M NaCl). 0.5-ml fractions were collected and assayed for PLA₂ activity.

Measurement of cPLA₂ mRNA levels. IL-1-induced changes in cPLA₂ mRNA levels were assessed by Northern analysis of total cellular RNA (20 µg) as described (13) with the following modifications. A gel-purified cDNA fragment which encoded the 2.9 kb human cytosolic PLA₂ (7, kindly provided by Dr. James D. Clark, Genetics Institute, Inc., Cambridge, MA), was labeled with [α -³²P]dCTP (Dupont/NEN, Boston, MA) by nick translation. Filters were hybridized with ³²P-labeled cPLA₂ probe overnight at 42°C in 5× SSPE (1× SSPE: 180 mM NaCl, 10 mM Na₂HPO₄·7H₂O, and 1 mM EDTA), 50% formamide, 5× Denhardt's solution, 1% SDS, 10% dextran sulfate, and 200 µg/ml denatured salmon sperm DNA. The blot was subsequently washed with 2× SSPE at room temperature for 30 min, 2× SSPE, 2% SDS at 50°C for 30 min and a final wash in 0.1× SSPE, 1% SDS at 52°C. After autoradiography for 15 h, bound probes were stripped from the blots by boiling and the filter was sequentially rehybridized with ³²P-labeled cDNA probes for rat nonpancreatic secretory PLA₂ (10, 14), murine IL-6 (14), and rat GAPDH (14). Autoradiographic densities were quantified by scanning densitometry (Scan Maker, Microtek, Torrance, CA) and appropriate software (Adobe Photoshop,

Adobe Systems, Mountainview, CA; and Scan Analysis, Biosoft, Cambridge, UK). cPLA₂ mRNA abundance was corrected for background and normalized for changes in GAPDH transcript abundance.

Protein phosphorylation and ³⁵S labeling of cPLA₂ and immunoprecipitation. Subconfluent mesangial cells in 100-mm plates were incubated for 1 h in serum-free, phosphate-free RPMI 1640 media, supplemented with 12.5 mM Hepes, pH 7.0, and NaHCO₃ (1.1 mg/ml). [³²P]orthophosphate (0.35 mCi/10-cm dish) was added for an additional 2 h. The cells were stimulated with vehicle or IL-1 (50 ng/ml) for the final 30 min of labeling, washed twice with ice cold PBS, and incubated for 10 min at 4°C in 0.4 ml of lysis buffer (150 mM NaCl, 50 mM Tris, pH 8.0, 1 mM EDTA, 1 mM EGTA, 50 mM NaF, 30 mM sodium pyrophosphate, 0.2 mM sodium orthovanadate, and 1.0% NP-40). Extracts from three 10-cm dishes were combined and spun at 200,000 g for 60 min. The resulting supernatants from control and stimulated mesangial cell lysates were matched for protein, and cleared with protein A-Sepharose (Pharmacia). The samples (1 ml) were then incubated overnight at 4°C with excess (10 µl) rabbit antisera against cPLA₂ or preimmune serum (7, kindly provided by Dr. James D. Clark). Immunocomplexes were precipitated using protein A-Sepharose, washed, and analyzed on 8% SDS-polyacrylamide gels. The ³²P-labeled proteins were detected by autoradiography using intensifying screens. In some experiments, the extent of phosphorylation was quantitated using storage phosphor technology and image analysis (15). The dried gel was placed on the photostimulatable storage phosphor imaging plate in a Molecular Dynamics (Sunnyvale, CA) exposure cassette for 2 d at room temperature and subsequently scanned by a Molecular Dynamics model 400A phosphor imager using a 10 mW helium-neon laser. Luminescence at 390 nm was collected, digitalized and stored. A digital image of the original incident radiation detected by the storage phosphor plates was generated, and bands of interest were quantified using Molecular Dynamics image quant software. Storage phosphor imaging plates have a large linear dynamic range and high sensitivity that allows accurate quantitative results (15). In some experiments, PLA₂ activity was quantified in cell lysates (110 µl) before and after cPLA₂ was immunoprecipitated using specific and preimmune antisera (3 µl).

To assess cytokine-induced changes in de novo protein synthesis, cells were washed, depleted of methionine for 30 min and transferred to fresh methionine-, cysteine-free RPMI media containing [³⁵S]-methionine (0.5 mCi/10-cm dish) for 1 h. IL-1 (50 ng/ml) was added for an additional 1 h. Cell-free lysates were prepared and immunoprecipitated, and immunoprecipitates were separated by electrophoresis as described above. ³⁵S-labeled proteins were imaged by fluorography at -70°C after soaking the gels with a fluor amplifying reagent (Amersham Corp.).

Immunoblot analysis of cPLA₂. 2 h before the experiment, cells maintained in complete medium were placed in RPMI media containing 10 mM Hepes, and treated with the indicated agonists. In some experiments, cells were serum-deprived for 24 h and then treated with IL-1 (10 ng/ml) or vehicle for an additional 24 h. Cells were then washed with PBS containing 4 mM EDTA, PMSF, leupeptin, and pepstatin A, and lysed as previously described (10). The samples (80 µg protein) were loaded on 8% SDS-polyacrylamide gel. To observe a phosphorylation-stimulated shift in cPLA₂ mobility (16), electrophoresis was continued until the 49-kD prestained protein marker (ovalbumin) had migrated to the end of the gel (a distance of 17 cm). The proteins were electrophoretically transferred (90 min at 85 V 4°C) to nitrocellulose membrane in a buffer containing 15% MeOH, 25 mM Tris, 192 mM glycine, 0.05% SDS, and 0.5 mM Na₃VO₄. The blot was blocked at 22°C for 2 h in PBS containing 0.2% Tween-20, 5% nonfat milk, and 0.5 mM Na₃VO₄. cPLA₂ protein was detected by incubation with a rabbit polyclonal antibody (1:2,000 dilution) against a recombinant fragment of human cPLA₂ (17) for 1 h followed by a 1 h incubation with a peroxidase-labeled, anti-rabbit IgG antibody (1:2,000 dilution, Kirkegaard & Perry, Gaithersburg, MD). The primary antibody cross-reacts with rat cPLA₂. Immunolocalized proteins were detected using a chemiluminescent detection system (ECL, Amersham Corp.).

All blots were stained subsequently with India ink to verify that proteins were equivalent and transferred.

Results

IL-1 stimulates [³H]arachidonate release from intact cells. We first determined if IL-1 activated a PLA₂ before the previously described induction at 6–10 h of the group II sPLA₂. Enzyme activity was initially assessed by measuring intracellular and extracellular [³H]arachidonic acid release from equilibrium-labeled glomerular mesangial cells in the presence or absence of the calcium ionophore A23187. Calcium ionophore was included because IL-1 does not increase cytosolic Ca²⁺ (18) and most PLA₂s are Ca²⁺ dependent. IL-1 alone in incubations as long as 30 min did not significantly increase [³H]arachidonate release. In contrast, cells exposed to IL-1 followed by ionophore showed enhanced [³H]arachidonate release (Fig. 1). The small increase in arachidonic acid release detected at 2 min was followed by a statistically significant increase at 5 and 30 min. As a positive control, cells were also incubated with the phorbol ester PMA in the absence or presence of ionophore (Fig. 1). PMA is known to activate cPLA₂ in vitro and “primes” intact cells for enhanced arachidonate metabolism (4, 19). Primed stimulation is the synergistic enhancement of cellular responses, when combinations of stimuli are utilized. PMA alone had little effect on mesangial cellular [³H]-arachidonate release but synergized with A23187 to stimulate a twofold increase in cytosolic arachidonate release (Fig. 1). These data suggest that IL-1, similar to PMA, primes cells for enhanced arachidonate release either by modifying a Ca²⁺-dependent PLA₂ or by activating a PLA₂ modulatory protein. Because IL-1 primes mesangial cells within 5 min for enhanced ionophore-stimulated arachidonate release, a PLA₂ that is distinct from the nonpancreatic group II sPLA₂ must be activated.

In vitro, the trifluoromethyl ketone analogue of arachidonic acid (AA-COCF₃) is a slow binding, but specific and potent inhibitor of cPLA₂ (11). We tested the effects of this compound in intact [³H]arachidonate-labeled cells. A 2-h preincubation with AA-COCF₃ completely inhibited IL-1-stimulated but not basal [³H]arachidonate release (Table I). Another arachidonate analogue, in which the trifluoromethyl ketone group is substituted with a -COCH₃ group (AA-COCH₃) and which is devoid of inhibitory activity, had no effect on IL-1-stimulated [³H]arachidonate release (Table I). Two inferences can be drawn from these data. First, IL-1 specifically activates cPLA₂ to stimulate arachidonic acid release in intact cells. Second, basal [³H]arachidonic acid release may result from a cPLA₂-independent mechanism since AA-COCF₃ had no effect in vehicle-stimulated cells (Table I), but almost completely inhibits cPLA₂ activity in the in vitro phosphatidylcholine vesicle assay using cell-free extracts (see below).

IL-1 stimulates cPLA₂ activity. Increased intracellular arachidonate may result from either activation of acylhydrolases and/or inhibition of acyltransferases (2). Since we and others have shown that IL-1 increases acyltransferase activity (20, 21), we focused on IL-1-stimulated deacylation reactions in cell-free extracts. Mesangial cells were stimulated for 2 h with IL-1 (10 ng/ml) or vehicle in the absence of serum. As a positive control, extracts of cells stimulated with PMA (300 nM) for the final 10 min of serum deprivation, were assayed as shown in Table II. Both IL-1 and PMA stimulated PLA₂ activity in subconfluent mesangial cells. IL-1 consistently increased

Table I. [³H]Arachidonic Acid Release in the Presence and Absence of a Fatty Acid Analogue Inhibitory for cPLA₂

Condition	[³ H]Arachidonic acid released cpm
Inactive FA analogue (AA-COCH ₃)	
Vehicle	10,781±486
IL-1	17,653±548
Inhibitory fatty acid analogue (AA-COCF ₃)	
Vehicle	9,889±913
IL-1	10,809±1078

Mesangial cells were labeled with [³H]arachidonic acid as described in Fig. 1. The cells were deprived serum for 2 h in the presence of the inhibitory fatty acid or the inactive fatty acid analogue. (Final concentrations, 25 μM.) Cells were then stimulated with vehicle or IL-1 (20 ng/ml) for 20 min. A23187 (0.25 μM) was subsequently added for 10 min. [³H]arachidonate release was quantified as described in Fig. 1. Results are the mean±SD of triplicate wells representative of duplicate experiments. Total ³H incorporation was the same in control and IL-1-stimulated cells in both treatment protocols.

activity by twofold or greater. As previously reported (4), PMA gave a two- to threefold increase in PLA₂ activity. PLA₂ activity in extracts of cells incubated with IL-1 increased progressively from 10 to 60 min (Fig. 2 A). At 30 min, IL-1 stimulated PLA₂ activity at concentrations between 1.0 and 10 ng/ml, and maximal activation was observed at 50–100 ng/ml (Fig. 2 B). LPS contamination of reagents did not account for these results. cPLA₂ activity in extracts of control cells and cells treated with heated IL-1 was similar (not shown). Addition of polymyxin B, a binder of LPS, with IL-1 did not affect cPLA₂ activity assayed in the phosphatidylcholine vesicle assay.

These results clearly show that IL-1-treated mesangial cells rapidly activate an enzymatic activity which releases arachidonate from the sn-2 position of exogenous [¹⁴C]-phosphatidylcholine in a standard assay for cPLA₂ activity. To characterize the IL-1-stimulated acylhydrolase activity, protein-matched cell extracts from control, IL-1-, and PMA-stimulated mesangial cells were fractionated on a Mono Q HR 5/5

Table II. Stimulation of cPLA₂ Activity in Rat Mesangial Cells by IL-1 and PMA

Condition	PLA ₂ activity pmol/min per mg
Control	238.9±27.9
IL-1	497.8±99.4*
PMA	664.3±92.8*

Subconfluent cultures of mesangial cells were incubated in serum-free media for 2 h, with or without IL-1 (10 ng/ml). In some cultures, PMA (300 nM) was added for the final 10 min of the serum-free incubation. Cell-free extracts were prepared by homogenization and ultracentrifugation as described in Methods, matched for protein, and assayed for PLA₂ activity. Results are expressed as mean values±SEM for triplicate determinations in four individual experiments. * *P* < 0.025, † *P* < 0.005 compared to the control value.




Figure 2. Time course and dose dependence of cPLA₂. (A) IL-1 stimulates cPLA₂ activity in a time-dependent manner. Mesangial cells were incubated in serum-free media for 2 h before the addition of IL-1 (10 ng/ml). Cell-free extracts were prepared from IL-1-treated or control cells at the indicated times, matched for protein, and assayed for PLA₂ activity. Data are presented as the mean value ± SEM of two to five individual experiments assayed in duplicate or triplicate (*n* = number of determinations). ***P* < 0.001, compared to the concurrent control value. (B) IL-1 stimulates cPLA₂ activity in a concentration-dependent manner. IL-1 at the indicated concentrations was added to mesangial cells incubated in serum-free media for 2 h. After 30 min, the cells were harvested and cell-free extracts prepared. The protein-matched samples were then assayed for PLA₂ activity. Results shown are the mean value ± SEM from two to five individual experiments. ***P* < 0.001 compared to the control values (*n* = number of determinations).

anion exchange column. PLA₂ activity was eluted with a NaCl gradient. > 95% of enzyme activity bound to the column. As shown in Fig. 3, the PLA₂ activity from each extract comigrated as a single peak, which was eluted with 0.4–0.45 M NaCl. This elution profile is consistent with that previously described for cPLA₂ in U937 and mesangial cells (4–6, 12). PLA₂ activity in fractionated extracts from both IL-1- and PMA-treated cells maintained higher specific activities than control extracts. IL-1-stimulated sPLA₂ is eluted with 1.2 M NaCl (M. Konieczkowski and J. R. Sedor, unpublished re-

sults). No PLA₂ activity was detected in fractions eluted with this salt concentration. Recoveries in all cases were > 75%.

As a complementary approach to verify that the IL-1-stimulated PLA₂ activity was cPLA₂, we incubated IL-1-stimulated extracts for 30 min at room temperature with either a rabbit polyclonal antiserum against human cPLA₂ (7) or preimmune antiserum. The specific antiserum recognizes rat cPLA₂ epitopes. After immunoprecipitation, residual supernatant PLA₂ activity was undetectable in control or IL-1-stimulated mesangial cell extracts. In contrast, PLA₂ activity was similar before and after immunoprecipitation with preimmune serum in extracts of IL-1-stimulated cells (control extracts, 141.1 ± 12.7 pmol/min per mg; untreated IL-1-stimulated extracts, 238.5 ± 13.6 pmol/min per mg; and residual cPLA₂ activity in IL-1-stimulated extracts pretreated with preimmune sera, 310.2 ± 21.5 pmol/min per mg, *n* = 2).

The effects of several PLA₂ inhibitors on cPLA₂ activity were tested in the phosphatidylcholine vesicle assay. The specific cPLA₂ inhibitor (11), AA-COCF₃, completely inhibited IL-1-stimulated cPLA₂ activity. In contrast, a control arachidonate analogue, AA-COCH₃, which is devoid of inhibitory activity (11), had no effect. cPLA₂ activity was 721.7 ± 57.7, 720.8 ± 33.8, and 95.0 ± 9.2 pmol/mg per min (mean ± SEM, *n* = 3) in vehicle (DMSO)-, AA-COCH₃ (10 μM)-, and AA-COCF₃ (10 μM)-treated extracts of IL-1-stimulated mesangial cells. Compounds that effectively inhibit sPLA₂ activation, less effectively inhibited cPLA₂ activity in extracts of both control and IL-1-stimulated cells. Aristolochic acid (150 μM), quinaquine (100 μM), and 7, 7-dimethyl-5,8-eicosadienoic acid (5 μM) inhibited IL-1-stimulated activity by 36 ± 4.1%, 28 ± 2.8%, and 18 ± 0.3%, respectively. A qualitatively similar pattern of inhibition was observed when these compounds were added to vehicle-stimulated cells (not shown). These data provide further evidence that IL-1 specifically can activate cPLA₂.

Other biochemical characteristics of the cytokine-activated cPLA₂ were determined. Unlike the IL-1-induced sPLA₂ activity (10), the PLA₂ activity rapidly stimulated by IL-1 hydrolyzed arachidonate from both [¹⁴C]phosphatidylcholine and [¹⁴C]phosphatidylethanolamine (data not shown). A pH (pH 4.5–9.0) dependence analysis of the IL-1-stimulated cPLA₂ demonstrated little activity below pH 7.0 and peak activity at an alkaline pH (approximately pH 9.0). Significant PLA₂ activity could not be measured in the extracts in the absence of Ca²⁺ of either vehicle- or IL-1-stimulated mesangial cells (data not shown). These characteristics are similar to those previously reported for both the mesangial cell and kidney enzymes (4, 12, 22).

Regulation of cPLA₂ by IL-1. The preceding data suggest that IL-1 causes a stable modification of cPLA₂ since the cytokine-activated enzyme stimulation was maintained after cell disruption, ultracentrifugation, and column fractionation. We explored two potential mechanisms which could result in a stable enhancement of cPLA₂ activity: (a) de novo synthesis of cPLA₂ or a stimulatory protein such as PLA₂-activating protein, and (b) a posttranslational modification of cPLA₂ which increases enzyme activity.

As shown in Fig. 4, Northern analysis demonstrated cPLA₂ mRNA in both control and cytokine-treated mesangial cells maintained in 17% serum. In four separate experiments, IL-1 had no consistent effect on cPLA₂ mRNA expression or activity (not shown) in incubations as long as 24 h, but induced expression of sPLA₂ at 24 h (Fig. 4) and IL-6 at 1 and 2 h (not




Figure 3. Characterization of PLA₂ activity from PMA and IL-1-treated rat mesangial cells by Mono-Q anion exchange chromatography. Cells were stimulated with PMA and IL-1 as described in Table I. Protein-matched, cell-free extracts were prepared and fractionated using a linear 0.15–1.0 M NaCl gradient. Each 0.5 ml fraction was assayed for PLA₂ activity. Peak PLA₂ activity coeluted at fraction 11 for each treatment. After fractionation, activity in IL-1- and PMA-stimulated extracts was greater than control samples.

shown). Moreover, IL-1 did not induce cPLA₂ protein when newly synthesized proteins in vehicle- and IL-1-stimulated (1 h) mesangial cells were radiolabeled with [³⁵S] methionine and enzyme levels were analyzed by immunoprecipitation (data not shown). Taken together, these data demonstrate that the rapid stimulation of cPLA₂ activity in serum-stimulated cells by IL-1 did not result from increased enzyme mass.

To assess whether the cytokine-stimulated cPLA₂ activity resulted from de novo synthesis of a PLA₂-activating protein,

cycloheximide was added for 30 min or 2 h before the addition of IL-1 (50 ng/ml) for a subsequent 30 min incubation. Cycloheximide (2 μg/ml) pretreatment had no effect on IL-1-stimulated cPLA₂ activity. The percent stimulation of enzyme activity by IL-1 above the control value was 239±78 and 211±37 (mean±SEM, n = 2) in the absence and presence of cycloheximide, respectively.

We next determined whether IL-1-stimulated cPLA₂ activity resulted from a post-translational modification, and hy-




Figure 4. IL-1 does not increase cPLA₂ mRNA levels in mesangial cells maintained in serum. Mesangial cells were incubated with IL-1 (10 ng/ml) or vehicle and the cells harvested for RNA at the indicated times. Total cellular RNA (20 μg) was analyzed for cPLA₂ message by Northern analysis as described in Methods. The filter was then rehybridized using probes for the 14-kD nonpancreatic secretory PLA₂ (sPLA₂), as a positive control for IL-1-stimulated transcript induction, and rat GAPDH (1.2 kb) to assess integrity and sample equivalence. cPLA₂ mRNA expression was corrected for changes in GAPDH transcript abundance. A representative Northern analysis is shown in A and mean (±SEM) change in relative cPLA₂ mRNA abundance in vehicle (Δ)- and IL-1 (○)-stimulated cells is shown from four separate experiments in B.

pothesized that IL-1 altered enzyme activity by phosphorylation for two reasons. First, cPLA₂ contains consensus phosphorylation sequences for several kinases (7, 16); second, IL-1 activates serine-threonine-kinases (23, 24). Mesangial cells were labeled in the absence of serum with [³²P]orthophosphate for 2 h and stimulated with IL-1 (50 ng/ml) for an additional 30 min. cPLA₂ was immunoprecipitated from the high-speed supernatants of control and IL-1-treated cells using polyclonal cPLA₂ antiserum or preimmune sera. As shown in Fig. 5 A, a specific protein was precipitated by the anti-cPLA₂ antiserum but not preimmune serum in both the control and IL-1-treated cells. The specific band was 97 kD, corresponding to the molecular mass of phosphorylated cPLA₂. ³²P incorporation into this band was nearly twofold greater than control in response to IL-1, a finding demonstrated in duplicate experiments. This IL-1-induced increase in cPLA₂ phosphorylation correlated with the magnitude of enzyme activation by IL-1 in whole cells at 30 min and is consistent with the idea that protein phosphorylation may regulate cPLA₂ activity. Others have demonstrated that agonist-induced phosphorylation and activation of cPLA₂ is associated with decreased mobility in SDS-polyacrylamide gels (16). Immunoblot analysis demonstrated that treatment of mesangial cells with IL-1 for 30 min caused shift in cPLA₂ mobility from a more rapidly to a more slowly migrating species (Fig. 5 B). As previously described (16), PMA similarly induced a shift in cPLA₂ mobility.

We next determined if phosphorylation of cPLA₂ by IL-1 resulted in increased enzyme activity (Table III). cPLA₂ activity was twofold higher in high-speed supernatants derived from IL-1-stimulated cells compared to vehicle-treated cells. In two separate experiments, treatment of the extracts with human acid phosphatase (0.4 IU) significantly reduced IL-1-stimulated enzyme activity but had no significant effect on enzyme activity in control cells. Taken together, these data suggest that IL-1 enhances cPLA₂ by phosphorylation. Basal cPLA₂ activity either is not dependent on phosphorylation or results from a phosphorylation event which is insensitive to acid phosphatase.

Mesangial cells used in the preceding experiments were either serum stimulated or acutely serum deprived (2 h). For two reasons, we next assessed IL-1-induced cPLA₂ expression in cells that were held for 24–48 h in 0.5% FBS. First, we have demonstrated that growth factors in serum attenuate induction of sPLA₂ mRNA activity (14). Second, serum contains glucocorticoids and dexamethasone prevents IL-1 induction of cPLA₂ in other cell types (25, 26). Fig. 6 depicts the results of a representative Northern and immunoblot analysis of cPLA₂ expression in control and IL-1-stimulated, serum-deprived cells. IL-1 increased cPLA₂ mRNA in 24-h incubations (Fig. 6 A) but not in incubations shorter than 6 h (not shown). Relative cPLA₂ mRNA levels, corrected for changes in GAPDH mRNA, were 2.4±0.4-fold (mean±SEM, n = 2) higher in IL-1-stimulated cells when compared to mesangial cells only exposed to vehicle. IL-1 concordantly increased cPLA₂ protein 1.7±0.1-fold (mean±SEM, n = 2) (Fig. 6 B). Under these conditions, cPLA₂ mobility is similar in vehicle- and IL-1-treated cells, suggesting that the amounts of hormones/growth factors contained in 0.5% of serum can maintain cPLA₂ in its phosphorylated state. Consistent with this hypothesis, we have noted that mesangial cells must be completely serum-deprived to maximally down-regulate cPLA₂ activity (M. Konieczkowski and J. R. Sedor, unpublished results). cPLA₂ enzyme activity was similarly doubled in serum-deprived mesangial




Figure 5. IL-1 stimulates the phosphorylation of cPLA₂. (A) Mesangial cells were labeled with [³²P]orthophosphate in the absence of serum for 2 h, and then IL-1 (50 ng/ml) was added for an additional 30 min. Cells were rapidly washed with ice-cold PBS and lysed in 1% NP-40. Lysates were incubated with preimmune serum and anti-serum to cPLA₂. Protein-antibody complexes were precipitated with protein A-Sepharose and analyzed on 8% denaturing polyacrylamide gels. [³²P]phosphorylated proteins were imaged using storage phosphor technology and changes in phosphorylation were quantified using image analysis as described in the Methods. (B) Mesangial cells were treated with IL-1 for 30 min and lysed. Proteins from cell lysates were analyzed by immunoblotting as described in Methods. cPLA₂ phosphorylation and activation causes a decreased mobility in SDS-polyacrylamide gels (22). IL-1 and PMA both induced cPLA₂ to migrate more slowly by phosphorylating the protein.

cells treated with IL-1 for 24 h (385.6±13.2 and 620.3±15.2 pmol/min per mg, extracts from control and IL-1-treated cells, mean±SEM, triplicate determinations). This finding is consistent with our previous results using the phosphatidylcholine vesicle assay (10). Although IL-1 markedly induces sPLA₂ by 24 h, phosphatidylcholine vesicles are a poor substrate for

Table III. cPLA₂ Activity in the Presence and Absence of Acid Phosphatase Treatment

Pretreatment	Stimulus	cPLA ₂ activity pmol/min per mg
None	Control (6)	322.1±33.3
	IL-1 (6)	633.1±79.2
Acid phosphatase	Control (9)	245.3±22.6 [‡]
	IL-1 (7)	331.1±68.3*

0.4 U of human acid phosphatase (Calbiochem Corp., 500 IU/ng) was added to 2.5–5 µg of mesangial cell extract prepared as described in Methods. The extracts and acid phosphatase were incubated for 15 min at 30°C, followed by the addition of 30 nCi [¹⁴C]arachidonyl-phosphatidylethanolamine and 5 mM CaCl₂. The samples were incubated subsequently for an additional 30 min at 37°C, and the cPLA₂ activity was quantified as described. cPLA₂ activity is expressed as the mean±SEM for triplicate determinations in two separate experiments. IL-1 significantly stimulated cPLA₂ activity over control, * *P* < 0.01, IL-1 stimulated vs. IL-1 stimulated and acid phosphatase pretreated; [‡] *P* > 0.1, vehicle stimulated vs. vehicle stimulated and acid phosphatase pretreated.

this PLA₂ isoform, suggesting the changes in PLA₂ activity measured by this assay are predominantly if not exclusively a result of cPLA₂ activation.

Discussion

IL-1 is a potent stimulus of eicosanoid metabolism (1, 9, 10). In the present study, we have demonstrated that IL-1, in the presence of ionophore, quickly stimulates arachidonate release from intact mesangial cells. In whole-cell lysates, IL-1 rapidly activates a PLA₂ isoform which we have characterized as cPLA₂. This increase in cytokine-stimulated cPLA₂ activity does not require new protein synthesis but is calcium dependent. IL-1 doubles ³²P incorporation into immunoprecipitable cPLA₂ protein consistent with the -fold stimulation of enzyme activity in similarly timed incubations. Importantly, the IL-1-stimulated enhancement of PLA₂ activity was signifi-

cantly reduced by acid phosphatase treatment. Taken together, these data suggest that IL-1-stimulated eicosanoid metabolism is mediated, at least acutely, by phosphorylation and activation of cPLA₂. These results support the premise, advanced by Lin et al. (16), that agonist-stimulated increases in intrinsic cPLA₂ activity are only apparent when a rise in cytosolic Ca²⁺ allows translocation of the enzyme to its substrate (4, 7, 27). IL-1 does not increase cytosolic Ca²⁺ in mesangial cells (18) and by itself does not increase arachidonate release (10, this paper). In contrast, arachidonic acid release stimulated by the Ca²⁺ mobilizing agonists, vasopressin (10) and ionophore (this paper), is doubled in IL-1-treated cells.

Previous studies have suggested that PLA₂ activity can be regulated by changes in its phosphorylation state. Sequence analysis of cPLA₂ has demonstrated consensus phosphorylation sites for the mitogen-activated protein (MAP) kinase and protein kinase C (7, 16), and cPLA₂ activity can be both positively and negatively regulated by agents that increase activity of certain kinases. Protein kinase C activators prime or potentiate agonist-induced arachidonate release, an effect that can be blocked in part by either protein kinase C down-regulation or PKC inhibitors (19, 28–30). EGF, similar to IL-1 and PMA, causes an increase in cPLA₂ activity that results from stable posttranslational modification (4, 12), and EGF receptor tyrosine kinase activity and autophosphorylation appears necessary for PLA₂ activation (31–33). One study has suggested that PLA₂ is directly activated by the EGF receptor tyrosine kinase (34). Calcium-dependent PLA₂ activity can be potentiated by the addition of either the catalytic subunit of the cAMP-dependent protein kinase or by casein kinase II and is inhibited by incubation with the Ca²⁺/calmodulin-dependent protein kinase II (35). Lin and co-workers have reported that treatment of CHO cells, overexpressing cPLA₂ with ATP, A23187, PMA, and thrombin increases arachidonate release from whole cells, stimulates cPLA₂ activity in vitro, and concomitantly increases ³²P incorporation into cPLA₂ exclusively on serine residues (16). The protein kinase C inhibitor staurosporine prevents ATP-stimulated cPLA₂ phosphorylation and activation, and treatment of ATP-stimulated cell lysates with acid phosphatase eliminates agonist-induced increases in cPLA₂ activity. Recent in vitro data has indicated that both MAP kinase and protein




Figure 6. Northern (A) and immunoblot (B) analysis of cPLA₂ mRNA and protein, respectively, in vehicle and IL-1-stimulated mesangial cells. Cells were held in medium containing 0.5% FBS for 24 h and subsequently stimulated with IL-1 for 24 h. cPLA₂ mRNA and protein expression were determined as described in Methods.

kinase C can activate cPLA₂ by phosphorylation (36, 37). Collectively, these data strongly implicate posttranslational changes in cPLA₂ phosphorylation state as an important regulator of enzyme activity.

Since cell lysates rather than purified protein have been used as the enzyme source, our data, as well as most of the data just summarized, are also consistent with activation of a cPLA₂ regulatory protein by phosphorylation. PLA₂ stimulatory proteins have been isolated from mammalian cells. Best characterized is a 28-kD protein, PLA₂-activating protein (PLAP) (38). Rapid, agonist-stimulated, de novo synthesis of PLAP is the primary mechanism of PLA₂ activation by PLAP (38). However, we found that cycloheximide does not prevent IL-1-stimulated cPLA₂ activation. A PLAP has also been purified from *Aplysia* (39). This PLA₂ stimulatory activity is increased in *Aplysia* extracts treated with phorbol dibutyrate, and in vitro, this protein is a substrate for protein kinase C (39). Phosphorylation also inactivates or enhances the degradation of lipocortins, molecules which bind to phospholipids and prevent PLA₂ activation by "substrate depletion." However phosphorylation of lipocortin I does not correlate temporally with EGF-stimulated activation of PLA₂ in cells overexpressing the wild-type EGF receptor (33). In vitro reconstitution experiments have demonstrated that both MAP kinase and protein kinase C phosphorylate and increase cPLA₂-specific catalytic activity (36, 37), suggesting that direct phosphorylation of cPLA₂ stimulates, at least in part, increased enzymatic activity in intact cells.

IL-1-dependent protein phosphorylations have been reported previously, and IL-1 stimulates protein serine-, threonine-kinase activity in a variety of cell types. The identity of the kinase(s) involved in IL-1-mediated transmembrane signaling is still speculative. IL-1-stimulated arachidonate release in fibroblasts does not require either protein kinase C or cAMP-dependent protein kinase activity (40). Cleveland maps of phosphorylated talin in fibroblasts demonstrate distinct phosphopeptides after PMA and IL-1 treatment, suggesting that the IL-1 activates kinases distinct from C kinase (41). Recently, IL-1 has been shown to rapidly stimulate MAP kinase activity (23, 24) and one study has suggested that IL-1 increases protein phosphorylation by inhibition of protein phosphatase activity (42). cPLA₂ is phosphorylated and activated in vitro by MAP kinase (36, 37), and we suggest that IL-1-stimulated MAP kinase phosphorylates cPLA₂ to increase its activity. Phosphorylation of distinct cPLA₂ domains may independently regulate enzyme activity. IL-1 and vasopressin synergistically increase arachidonate release; vasopressin increases protein kinase C activity in mesangial cells but IL-1 does not. In vitro protein kinase C and MAP kinase phosphorylate distinct residues within recombinantly expressed cPLA₂ domains (37). However other workers have suggested protein kinase C and MAP kinase identically phosphorylate the same amino acid residue (36).

Stimulation of PLA₂ activity by IL-1 may be cell specific. Similar to our findings, IL-1 increased a PLA₂ activity, with Ca²⁺ requirements similar to cPLA₂, in rheumatoid synovial fibroblasts (43). Treatment of a human lung fibroblast line with IL-1 results in cPLA₂ phosphorylation (25). However, in contrast to our findings in mesangial cells (10, this paper), IL-1 only stimulated PGE₂ synthesis in fibroblasts after prolonged (> 5 h) incubation and did not induce sPLA₂ activity (25). Human IL-1β added to guinea pig eosinophils negatively regulated PLA₂ activity through an indirect, Ca²⁺ independent

mechanism (44). In a preliminary report, Clark and co-workers have demonstrated that de novo PLAP synthesis mediates IL-1-stimulated PLA₂ activity in EL-4 cells (45). This cell specific regulation of IL-1-stimulated PLA₂ activity suggests that arachidonic acid hydrolysis may be an important determinant in the cellular response to inflammation.

cPLA₂ activity clearly is regulated at multiple levels. In addition to a rapid posttranslational modification of cPLA₂, IL-1 induces, in 24-h incubations a small (approximately twofold) but consistent increase in cPLA₂ mRNA, protein, and activity. This effect was only detected in mesangial cells held in 0.5% FBS but not those cells maintained in serum. Growth factors (14) and/or glucocorticoids (25, 26) present in serum may negatively regulate IL-1-induced cPLA₂ expression. Long-term incubations (> 8 h) with macrophage colony-stimulating factor (46), TNF (25), epidermal growth factor (47) and PMA (47) also increases cPLA₂ expression. cPLA₂ induction by IL-1 and TNF is not surprising, in that both cytokines activate the transcriptional factors, AP-1 and NFκB, and the 5' flanking region of the cPLA₂ gene contains AP-1 and κB binding elements (48).

In summary, IL-1 regulates the activity of cPLA₂ in a biphasic, time-dependent manner. A rapid increase in catalytic activity results from protein phosphorylation. More prolonged incubation with IL-1 can increase cPLA₂ mass in resting but not serum-activated cells. cPLA₂ activation provides one mechanism for biological events induced by this cytokine. Arachidonic acid release and enhanced eicosanoid metabolism are hallmarks of glomerular inflammation. Increased prostaglandin, HETE, and leukotriene synthesis mediate both the hemodynamic alterations and the cellular activation which characterize inflammatory injury. Recent studies also have demonstrated that these same fatty acid mediators comprise an important intracellular, transmembrane signaling system. Arachidonic acid and/or its metabolites activate GTP-binding proteins (49), regulate the activity of Ras-GTPase activating protein (50), stimulate kinase activity (51), control ion channel function (52), and can activate latent transactivating factors (53). These interactions between arachidonate and other second messenger pathways provide mechanisms by which IL-1-mediated activation of cPLA₂, not only results in the synthesis of proinflammatory lipid mediators, but also initiates a specific program of phenotypic responses in the mesangial cell.

Acknowledgments

We thank Drs. Peter Lomedico, Alvin Stern, and Swapan Roy for generously providing recombinant IL-1α, Dr. James Clark for the cPLA₂ cDNA and antisera, and Kathleen Kashnier for expert secretarial assistance. Drs. Cathleen Carlin, Steven Emancipator, and Michael Ganz critically reviewed the manuscript.

This work was supported by the National Institute of Health grant DK38558, K04 DK01933 and the Community Dialysis Centers of Cleveland. Dr. Sedor is an Established Investigator of the American Heart Association.

References

1. Dinarello, C. A. 1991. Interleukin-1 and interleukin-1 antagonism. *Blood*. 77:1627-1652.
2. Irvine, R. F. 1982. How is the level of free arachidonic acid controlled in mammalian cells? *Biochem. J.* 204:3-16.
3. Davidson, F. F., and E. A. Dennis. 1990. Evolutionary relationships and implications for the regulation of phospholipase A₂ from snake venom to human secreted forms. *J. Mol. Evol.* 31:228-238.
4. Gronich, J. H., J. V. Bonventre, and R. A. Nemenoff. 1988. Identification

- and characterization of a hormonally regulated form of phospholipase A₂ in rat renal mesangial cells. *J. Biol. Chem.* 263:16645-16651.
5. Kramer, R. M., E. F. Roberts, J. Manetta, and J. E. Putnam. 1991. The Ca²⁺-sensitive cytosolic phospholipase A₂ is a 100-kDa protein in human monoblast U937 cells. *J. Biol. Chem.* 266:5268-5272.
 6. Clark, J. D., N. Milona, and J. L. Knopf. 1990. Purification of a 110-kilodalton cytosolic phospholipase A₂ from the human monocytic cell line U937. *Proc. Natl. Acad. Sci. USA.* 87:7708-7712.
 7. Clark, J. D., L. L. Lin, R. W. Kriz, C. S. Ramesha, L. A. Sultzman, A. Y. Lin, N. Milona, and J. L. Knopf. 1991. A novel arachidonic acid-selective cytosolic PLA₂ contains a Ca²⁺-dependent translocation domain with homology to PKC and GAP. *Cell.* 65:1043-1051.
 8. Emancipator, S. N., and J. R. Sedor. 1992. Cytokines and renal disease. In *Cytokines in Health and Disease: Physiology and Pathophysiology*, D. G. Remick and S. L. Kunkel, editors. Marcel Dekker, Inc., New York. 467-488.
 9. Sedor, J. R., Y. Nakazato, and M. Konieczkowski. 1992. IL-1 and the mesangial cell. *Kidney Int.* 41:595-599.
 10. Nakazato, Y., M. S. Simonson, W. H. Herman, M. Konieczkowski, and J. R. Sedor. 1991. Interleukin-1 α stimulates prostaglandin biosynthesis in serum-activated mesangial cells by induction of a non-pancreatic (Type II) phospholipase A₂. *J. Biol. Chem.* 266:14119-14127.
 11. Street, I. P., H. Lin, F. Laliberte, F. Ghomashchi, Z. Wang, H. Perrier, N. M. Tremblay, Z. Huang, P. K. Weech, and M. H. Gelb. 1993. Slow- and tight-binding inhibitors of the 85-kDa human phospholipase A₂. *Biochemistry.* 32:5935-5940.
 12. Bonventre, J. V., J. H. Gronich, and R. A. Nemenoff. 1990. Epidermal growth factor enhances glomerular mesangial cell soluble phospholipase A₂ activity. *J. Biol. Chem.* 265:4934-4938.
 13. Werber, H. I., S. N. Emancipator, M. L. Tykocinski, and J. R. Sedor. 1987. The interleukin 1 gene is expressed by rat glomerular mesangial cells and is augmented in immune complex glomerulonephritis. *J. Immunol.* 138:3207-3212.
 14. Konieczkowski, M., and J. R. Sedor. 1993. Cell-specific regulation of type II phospholipase A₂ expression in rat mesangial cells. *J. Clin. Invest.* 92:2524-2532.
 15. Johnston, R. F., S. C. Pickett, and D. L. Barker. 1990. Autoradiography using storage phosphor technology. *Electrophoresis.* 11:355-360.
 16. Lin, L. L., A. Y. Lin, and J. L. Knopf. 1992. Cytosolic phospholipase A₂ is coupled to hormonally regulated release of arachidonic acid. *Proc. Natl. Acad. Sci. USA.* 89:6147-6151.
 17. Winitz, S., S. K. Gupta, N. X. Qian, L. E. Heasley, R. A. Nemenoff, and G. L. Johnson. 1993. Expression of a mutant G₁₂ α subunit inhibits ATP and thrombin stimulation of cPLA₂-mediated arachidonic acid release independent of Ca²⁺ and MAP kinase regulation. *J. Biol. Chem.* In press.
 18. Kester, M., M. S. Simonson, P. Mene, and J. R. Sedor. 1989. Interleukin-1 generates transmembrane signals from phospholipids through novel pathways in cultured rat mesangial cells. *J. Clin. Invest.* 83:718-723.
 19. Bonventre, J. V., and M. Swidler. 1988. Calcium dependency of prostaglandin E₂ production in rat glomerular mesangial cells. *J. Clin. Invest.* 82:168-176.
 20. Nakazato, Y., and J. R. Sedor. 1992. IL-1 α increases arachidonyl-CoA: lysophospholipid acyltransferase activity and stimulates [³H]arachidonate incorporation into phospholipids in rat mesangial cells. *Life Sci.* 50:2075-2082.
 21. Bursten, S. L., W. E. Harris, K. Bomszyk, and D. Lovett. 1991. Interleukin-1 rapidly stimulates lysophosphatidate acyltransferase and phosphatidate phosphohydrolase activities in human mesangial cells. *J. Biol. Chem.* 266:20732-20743.
 22. Gronich, J. H., J. V. Bonventre, and A. Nemenoff. 1990. Purification of a high-molecular-mass form of phospholipase A₂ from rat kidney activated at physiological calcium concentrations. *Biochem. J.* 271:37-43.
 23. Guy, G. R., S. P. Chua, N. S. Wong, S. B. Ng, and Y. H. Tan. 1991. Interleukin 1 and tumor necrosis factor activate common multiple protein kinases in human fibroblasts. *J. Biol. Chem.* 266:14343-14352.
 24. Bird, T. A., P. R. Sleath, P. C. deRoos, S. K. Dower, and G. D. Virca. 1991. Interleukin-1 represents a new modality for the activation of extracellular signal-regulated kinases/microtubule-associated protein-2 kinases. *J. Biol. Chem.* 266:22661-22670.
 25. Lin, L. L., A. Y. Lin, and D. L. DeWitt. 1992. Interleukin-1 α induces the accumulation of cytosolic phospholipase A₂ and the release of prostaglandin E₂ in human fibroblasts. *J. Biol. Chem.* 267:23451-23454.
 26. Hoeck, W. G., C. S. Ramesha, D. J. Chang, N. Fan, and R. A. Heller. 1993. Cytosolic phospholipase A₂ activity and gene expression are stimulated by tumor necrosis factor: dexamethasone blocks the induced synthesis. *Proc. Natl. Acad. Sci. USA.* 90:4475-4479.
 27. Hack, N., P. Clayman, and K. Skorecki. 1990. A role for G-proteins in the epidermal growth factor stimulation of phospholipase A₂ in rat kidney mesangial cells. *Biosci. Rep.* 10:353-362.
 28. Parker, J., L. W. Daniel, and M. Waite. 1987. Evidence of protein kinase C involvement in phorbol diester-stimulated arachidonic acid release and prostaglandin synthesis. *J. Biol. Chem.* 262:5385-5393.
 29. Burch, R. M., A. L. Ma, and J. Axelrod. 1988. Phorbol esters and diacyl-glycerols amplify bradykinin-stimulated prostaglandin synthesis in swiss 3T3 fibroblasts. *J. Biol. Chem.* 263:4764-4767.
 30. Margolis, B. L., J. V. Bonventre, S. G. Kremer, J. E. Kudlow, and K. L. Skorecki. 1988. Epidermal growth factor is synergistic with phorbol esters and vasopressin in stimulating arachidonate release and prostaglandin production in renal glomerular mesangial cells. *Biochem. J.* 249:587-592.
 31. Goldberg, H. J., M. M. Viegas, B. L. Margolis, J. Schlessinger, and K. L. Skorecki. 1990. The tyrosine kinase activity of the epidermal-growth-factor receptor is necessary for phospholipase A₂ activation. *Biochem. J.* 267:461-465.
 32. Hack, N., B. L. Margolis, A. Ullrich, J. Schlessinger, and K. L. Skorecki. 1991. Distinct structural specificities for functional coupling of the epidermal growth factor receptor to calcium-signalling versus phospholipase A₂ responses. *Biochem. J.* 275:563.
 33. Clark, S., and M. Dunlop. 1991. Modulation of phospholipase A₂ activity by epidermal growth factor (EGF) in CHO cells transfected with human EGF receptor: role of receptor cytoplasmic subdomain. *Biochem. J.* 274:715-721.
 34. Peppelenbosch, M. P., L. G. J. Tertoolen, J. den Hertog, and S. W. deLaat. 1992. Epidermal growth factor activates calcium channels by phospholipase A₂/5-lipoxygenase-mediated leukotriene C₄ production. *Cell.* 69:295-303.
 35. Piomelli, D., and P. Greengard. 1991. Bidirectional control of phospholipase A₂ activity by Ca²⁺/calmodulin-dependent protein kinase II, cAMP-dependent protein kinase, and casein kinase II. *Proc. Natl. Acad. Sci. USA.* 88:6770-6774.
 36. Lin, L. L., M. Wartmann, A. Y. Lin, J. L. Knopf, A. Seth, and R. J. Davis. 1993. cPLA₂ is phosphorylated and activated by MAP kinase. *Cell.* 72:269-278.
 37. Nemenoff, R. A., S. Winitz, N. X. Qian, V. V. Putten, G. L. Johnson, and L. E. Heasley. 1993. Phosphorylation and activation of a high molecular weight form of phospholipase A₂ by p42 microtubule-associated protein 2 kinase and protein kinase C. *J. Biol. Chem.* 268:1960-1964.
 38. Clark, M. A., L. E. Ozgur, T. M. Conway, J. Disposito, S. T. Crooke, and J. S. Bomalaski. 1991. Cloning of a phospholipase A₂-activating protein. *Proc. Natl. Acad. Sci. USA.* 88:5418-5422.
 39. Calignano, A., D. Piomelli, T. C. Sactor, and J. H. Schwartz. 1991. A phospholipase A₂-stimulating protein regulated by protein kinase C in aplysia neurons. *Mol. Brain Res.* 9:347-351.
 40. Cisar, L. A., R. J. Schimmel, and E. Mochan. 1991. Interleukin-1 stimulation of arachidonic acid release from human synovial fibroblasts; blockade by inhibitors of protein kinases and protein synthesis. *Cell. Signalling.* 3:189-199.
 41. Qwarnstrom, E. E., S. A. MacFarlane, R. C. Page, and S. K. Dower. 1991. Interleukin 1 β induces rapid phosphorylation and redistribution of talin: a possible mechanism for modulation of fibroblast focal adhesion. *Proc. Natl. Acad. Sci. USA.* 88:1232-1236.
 42. Guy, G. R., X. Cao, S. P. Chua, and Y. H. Tan. 1992. Okadaic acid mimics multiple changes in early protein phosphorylation and gene expression induced by tumor necrosis factor or interleukin-1. *J. Biol. Chem.* 267:1846-1852.
 43. Hulkower, K. I., W. C. Hope, T. Chen, C. M. Anderson, J. W. Coffey, and D. W. Morgan. 1992. Interleukin-1 β stimulates cytosolic phospholipase A₂ in rheumatoid synovial fibroblasts. *Biochem. Biophys. Res. Commun.* 184:712-718.
 44. Debbaghi, A., R. Hidi, B. B. Vargaftig, and L. Touqui. 1992. Inhibition of phospholipase A₂ activity in guinea pig eosinophils by human recombinant IL-1 β . *J. Immunol.* 149:1374-1380.
 45. Bomalaski, J. S., M. R. Steiner, P. L. Simon, and M. A. Clark. 1992. Interleukin-1 induces increased phospholipase A₂ activity, synthesis of phospholipase A₂ activating protein (PLAP), and release of linoleic acid and interleukin-2 from the murine (EL-4) T helper cell line. *Clin. Res.* 40:185A. (Abstr.)
 46. Nakamura, T., L. Lin, S. Kharbada, J. Knopf, and D. Kufe. 1992. Macrophage colony stimulating factor activates phosphatidylcholine hydrolysis by cytoplasmic phospholipase A₂. *EMBO (Eur. Mol. Biol. Organ.) J.* 11:4917-4922.
 47. Maxwell, A. P., H. J. Goldberg, A. H. N. Tay, Z. Li, G. S. Arbus, and K. L. Skorecki. 1993. Epidermal growth factor and phorbol myristate acetate increase expression of the mRNA for cytosolic phospholipase A₂ in glomerular mesangial cells. *Biochem. J.* 295:763-766.
 48. Tay, A., H. Goldberg, P. Maxwell, Z. Li, and K. Skorecki. 1993. Isolation and characterization of 5' flanking region of the rat cytosolic phospholipase A₂ (cPLA₂) gene. *J. Am. Soc. Nephrol.* 4:502. (Abstr.)
 49. Abramson, S. B., J. Leszczynska-Piziak, and G. Weissmann. 1991. Arachidonic acid as a second messenger. Interactions with a GTP-binding protein of human neutrophils. *J. Immunol.* 147:231-236.
 50. Han, J. W., F. McCormick, and I. G. Marcara. 1991. Regulation of ras-GAP and the neurofibromatosis-1 gene product by eicosanoids. *Science (Wash. DC).* 252:576-579.
 51. Khan, W. A., G. C. Blobe, and Y. A. Hannun. 1992. Activation of protein kinase C by oleic acid. *J. Biol. Chem.* 267:3605.
 52. Kim, D., and D. E. Clapham. 1989. Potassium channels in cardiac cells activated by arachidonic acid and phospholipids. *Science (Wash. DC).* 244:1174-1176.
 53. Hannigan, G. E., and B. R. G. Williams. 1991. Signal transduction by interferon- α through arachidonic acid metabolism. *Science (Wash. DC).* 251:204-207.