

Table S3. NSHL X-linked, Y-linked and modifier loci

LocusName	Position	References
DFN1	Xq22	[1]
DFN2	Xq22	[2]
DFN3	Xq21.1	[3]
DFN4	Xp21.2	[4]
DFN6	Xp22	[5]
DFNY1	Y	[6]
DFNM1	1q24	[7]

References

- Tranebjærg L, Schwartz C, Eriksen H, Andreasson S, Ponjavic V, et al. (1995) A new X linked recessive deafness syndrome with blindness, dystonia, fractures, and mental deficiency is linked to Xq22. *J Med Genet* 32: 257-263.
- Tyson J, Bellman S, Newton V, Simpson P, Malcolm S, et al. (1996) Mapping of DFN2 to Xq22. *Hum Mol Genet* 5: 2055-2060.
- Bach I, Robinson D, Thomas N, Ropers H, Cremers F (1992) Physical fine mapping of genes underlying X-linked deafness and non fra (X)-X-linked mental retardation at Xq21. *Hum Genet* 89: 620-624.
- Lalwani A, Brister J, Fex J, Grundfast K, Pikus A, et al. (1994) A new nonsyndromic X-linked sensorineural hearing impairment linked to Xp21.2. *Am J Hum Genet* 55: 685-694.
- del Castillo I, Villamar M, Sarduy M, Romero L, Herraiz C, et al. (1996) A novel locus for non-syndromic sensorineural deafness (DFN6) maps to chromosome Xp22. *Hum Mol Genet* 5: 1383-1387.
- Wang Q, Lu C, Li N, Rao S, Shi Y, et al. (2004) Y-linked inheritance of non-syndromic hearing impairment in a large Chinese family. *J Med Genet* 41(6): e80.
- Riazuddin S, Castelein C, Ahmed Z, Lalwani A, Mastroianni M, et al. (2000) Dominant modifier DFNM1 suppresses recessive deafness DFNB26. *Nat Genet* 26: 431-434.