

## Supporting Information

### Synthetic Utility of an Isolable Nucleoside Phosphonium Salt

Suyeal Bae and Mahesh K. Lakshman\*


Department of Chemistry, The City College and The City University of New York,  
160 Convent Avenue, New York, New York 10031-9198.

#### Table of Contents



Information	Page
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>O</i> <sup>2</sup> -tris(dimethylamino)phosphonium-2'-deoxyxanthosine hexafluorophosphate ( <b>3</b> ) in CDCl <sub>3</sub>	S3
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>O</i> <sup>2</sup> -tris(dimethylamino)phosphonium-2'-deoxyxanthosine hexafluorophosphate ( <b>3</b> ) in CDCl <sub>3</sub>	S4
<sup>31</sup> P NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>O</i> <sup>2</sup> -tris(dimethylamino)phosphonium-2'-deoxyxanthosine hexafluorophosphate ( <b>3</b> ) in CDCl <sub>3</sub>	S5
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(morpholin-4-yl)-2'-deoxyinosine ( <b>4a</b> ) in CDCl <sub>3</sub>	S6
<sup>13</sup> C NMR spectrum <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(morpholin-4-yl)-2'-deoxyinosine ( <b>4a</b> ) in CDCl <sub>3</sub>	S7
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(piperidin-1-yl)-2'-deoxyinosine ( <b>4b</b> ) in CDCl <sub>3</sub>	S8
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(piperidin-1-yl)-2'-deoxyinosine ( <b>4b</b> ) in CDCl <sub>3</sub>	S9
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(pyrrolidin-1-yl)-2'-deoxyinosine ( <b>4c</b> ) in CDCl <sub>3</sub>	S10
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(pyrrolidin-1-yl)-2'-deoxyinosine ( <b>4c</b> ) in CDCl <sub>3</sub>	S11
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(1,4-dioxa-8-azaspiro[4.5]dec-8-yl)-2'-deoxyinosine ( <b>4d</b> ) in CDCl <sub>3</sub>	S12
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(1,4-dioxa-8-azaspiro[4.5]dec-8-yl)-2'-deoxyinosine ( <b>4d</b> ) in CDCl <sub>3</sub>	S13
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-[4-(4-nitrophenyl)-piperizin-1-yl]-2'-deoxyinosine ( <b>4e</b> ) in CDCl <sub>3</sub>	S14
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-[4-(4-nitrophenyl)-piperizin-1-yl]-2'-deoxyinosine ( <b>4e</b> ) in CDCl <sub>3</sub>	S15
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -methyl- <i>N</i> <sup>2</sup> -[(2- <i>N,N</i> -dimethylamino)ethyl]-2'-deoxyguanosine ( <b>4f</b> ) in CDCl <sub>3</sub>	S16

<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -methyl- <i>N</i> <sup>2</sup> -[(2- <i>N,N</i> -dimethylamino)ethyl]-2'-deoxyguanosine ( <b>4f</b> ) in CDCl <sub>3</sub>	S17
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -propyl-2'-deoxyguanosine ( <b>4g</b> ) in CDCl <sub>3</sub>	S18
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -propyl-2'-deoxyguanosine ( <b>4g</b> ) in CDCl <sub>3</sub>	S19
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(morpholin-4-yl)-2'-deoxyinosine ( <b>5a</b> ) in CDCl <sub>3</sub>	S20
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(piperidin-1-yl)-2'-deoxyinosine ( <b>5b</b> ) in CDCl <sub>3</sub>	S21
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(pyrrolidin-1-yl)-2'-deoxyinosine ( <b>5c</b> ) in CDCl <sub>3</sub>	S22
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-(1,4,dioxo-8-azaspiro[4.5]dec-8-yl)-2'-deoxyinosine ( <b>5d</b> ) in CDCl <sub>3</sub>	S23
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -methyl- <i>N</i> <sup>2</sup> -[(2- <i>N,N</i> -dimethylamino)ethyl]-2'-deoxyguanosine ( <b>5f</b> ) in CDCl <sub>3</sub>	S24
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -propyl-2'-deoxyguanosine ( <b>5g</b> ) in CDCl <sub>3</sub>	S25
<sup>1</sup> H NMR spectrum of <i>O</i> <sup>6</sup> -benzyl- <i>N</i> <sup>2</sup> -(3-benzyloxy-1-propyl)-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2'-deoxyguanosine ( <b>7</b> ) in CDCl <sub>3</sub>	S26
<sup>13</sup> C NMR spectrum of <i>O</i> <sup>6</sup> -benzyl- <i>N</i> <sup>2</sup> -(3-benzyloxy-1-propyl)-3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2'-deoxyguanosine ( <b>7</b> ) in CDCl <sub>3</sub>	S27
<sup>1</sup> H NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -(3-hydroxy-1-propyl)-2'-deoxyguanosine ( <b>8</b> ) in CDCl <sub>3</sub>	S28
<sup>13</sup> C NMR spectrum of 3',5'-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)- <i>N</i> <sup>2</sup> -(3-hydroxy-1-propyl)-2'-deoxyguanosine ( <b>8</b> ) in CDCl <sub>3</sub>	S29
<sup>1</sup> H NMR spectrum of ~1:1 diastereomeric mixture of 3-[3,5-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-deoxy-β- <i>D</i> - <i>erythro</i> -pentofuranosyl]-8-hydroxy-5,6,7,8-pyrimido[1,2- <i>a</i> ]purin-10(3 <i>H</i> )-one ( <b>9</b> ) in CDCl <sub>3</sub>	S30
<sup>13</sup> C NMR spectrum of ~1:1 diastereomeric mixture of 3-[3,5-bis- <i>O</i> -( <i>tert</i> -butyldimethylsilyl)-2-deoxy-β- <i>D</i> - <i>erythro</i> -pentofuranosyl]-8-hydroxy-5,6,7,8-pyrimido[1,2- <i>a</i> ]purin-10(3 <i>H</i> )-one ( <b>9</b> ) in CDCl <sub>3</sub>	S31



SB-1205-07-57-column-1H





SB-1205-04-74-column-C13




SB-1205-07-57-column-P31



SB-1205-05-09-1H




SB-1205-05-09-13C





ppm (t1)

SB-1205-07-12-1H








SB-1205-07-12-13C





SB-1205-07-13-1H





SB-1205-07-13-13C



SB-1205-07-14-1H




SB-1205-07-14-13C





ppm (t1)

SB-1205-07-15-column-1H




SB-1205-07-15-column-13C





ppm (t1)

SB-1205-07-16-column-1H







SB-1205-07-16-column-13C





SB-1205-07-17-1H





ppm (t1)

SB-1205-07-17-13c





SB-1205-07-19-1H





ppm (t1)

SB-1205-07-22-1H





ppm (t1)

SB-1205-07-24-column-1H





ppm (t1)

SB-1205-07-23-1H



ppm (t1)



SB-1205-07-27-1H





ppm (t1)



SB-1205-07-26-1H





SB-1205-07-66-column-1H





ppm (t1)



SB-1205-07-66-column-13C





SB-1205-07-60-column-1H





SB-1205-07-60-column-13C



SB-1205-08-11-column-1H



SB-1205-08-11-column-13C



ppm (t1)