

Fc γ Receptor IIIb Enhances Fc γ Receptor IIa Function in an Oxidant-dependent and Allele-sensitive Manner

Jane E. Salmon,* Sean S. Millard, Nina L. Brogle, and Robert P. Kimberly*

The Department of Medicine, The Hospital for Special Surgery and The New York Hospital; and *The Graduate Program in Immunology, The Cornell University Medical College, New York 10021

Abstract

Two classes of receptors for IgG, Fc γ RIIa and Fc γ RIIIb, both of which exist in two allelic forms, are expressed on human neutrophils. Neutrophils from normal donors, homozygous for the different allelic phenotypes of Fc γ RIIIb, have significantly different levels of Fc γ receptor-mediated phagocytosis of IgG-opsonized erythrocytes (EA). However, the observation that Fc γ RIIIb mediates phagocytosis of specific mAb-targeted erythrocytes poorly suggests that this receptor may influence EA internalization by Fc γ RIIa in an allele-sensitive fashion. Donors homozygous for the NA1 allele of Fc γ RIIIb showed greater activation of Fc γ RIIa after Fc γ RIIIb cross-linking than donors homozygous for the NA2 allele of Fc γ RIIIb. This increase in receptor-specific internalization reflects both an increase in ligand binding by Fc γ RIIa and an increase in internalization efficiency of targets bound. Activation of Fc γ RIIa by Fc γ RIIIb is transferable by supernatants from activated cells and is blocked by inhibitors of reactive oxygen species and the H₂O₂-myeloperoxidase-chloride system and by serine protease inhibitors. Thus, cross-linking of Fc γ RIIIb, which leads to neutrophil degranulation and the generation of reactive oxygen intermediates, in turn alters Fc γ RIIa avidity and efficiency. These oxidant-mediated changes in Fc γ RIIa function provide a novel mechanism for receptors to collaborate in both an autocrine and paracrine fashion. The allele sensitivity of these effects suggests that Fc γ receptor polymorphisms may be inherited disease susceptibility factors in host defense against infection and in the development of autoimmunity. (*J. Clin. Invest.* 1995. 95:2877-2885.) **Key words:** phagocytosis • proteases • Fc γ receptors • oxidants • immunoglobulin G

Address correspondence to Jane E. Salmon, M.D., The Hospital for Special Surgery, 535 East 70th Street, New York, NY 10021. Phone: 212-606-1422; FAX: 212-717-1192.

Received for publication 11 May 1994 and in revised form 23 January 1995.

1. *Abbreviations used in this paper:* AI, attachment index; E, erythrocytes; EA, IgG-sensitized bovine erythrocytes; E_B, biotinylated erythrocytes; E_{AB}, streptavidin coated E_B; E-hIgG2, E coated with human IgG2; E-IV.3, E coated with IV.3 Fab; Fc γ R, receptors for Fc portion of IgG in human cells; Fc γ RIIa, 40-kD receptor on human neutrophils and monocytes for Fc portion of IgG; Fc γ RIIIb, 50-78-kD receptor on human neutrophils for Fc portion of IgG; PABA, *p*-aminobenzamide; PE, phycoerythrin; PI, phagocytic index; TLCK, *N*^ε-tosyl-L-lysyl-chloromethyl ketone.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/95/06/2877/09 \$2.00

Volume 95, June 1995, 2877-2885

Introduction

Receptors for the Fc region of IgG, Fc γ R,¹ provide the crucial link between humoral immunity and the IgG-triggered effector functions by inflammatory cells. Three families of human Fc γ R (Fc γ RI, Fc γ RII, and Fc γ RIII) have been identified. Within these Fc γ R families, multiple distinct genes and alternative splicing variants lead to a variety of receptor isoforms that have differences in structure and distinct functional capacities (for reviews see references 1-4). In addition to this diversity, two receptor isoforms found on phagocytic cells have codominantly expressed, allelic polymorphisms—the NA1 and NA2 alleles of Fc γ RIIIb and the HR-LR alleles of Fc γ RIIa—which influence Fc γ R function.

Neutrophils from normal donors, homozygous for the two different allelic phenotypes of Fc γ RIIIb, have significantly different levels of Fc γ receptor-mediated phagocytosis independent of the allelic phenotype of the second type of Fc γ receptor on neutrophils, Fc γ RIIa (5). Because experiments using antireceptor mAb-targeted erythrocytes indicate that Fc γ RIIIb mediates phagocytosis poorly (6), the importance of Fc γ RIIIb NA phenotype in quantitative EA phagocytosis implies that a model of "one receptor-one function" is inadequate and that Fc γ RIIIb may collaborate interactively with other receptors. Among the different paradigms of receptor cooperation, there may be physical association of receptors that enhances binding and signaling and interaction independent of physical association. For example, the IL-2 receptor can assemble multiple distinct receptor chains into a complex, and different combinations of α , β , and γ chains have distinct binding and signaling capacities (7, 8). Similarly, TGF β signals through a heteromeric complex composed of receptor I and receptor II, but a separate nonsignaling TGF β -binding membrane protein (TGF β receptor type III) captures the ligand and delivers it to the signaling complex (9). In contrast to these systems, signaling through the T cell receptor upregulates the avidity of CD2 for CD58 (LFA-1) through inside-out signaling requiring specific domains of the CD2 cytoplasmic tail (10, 11). Similarly, activation of β_2 and β_3 integrins as a consequence of receptor phosphorylation is induced by cross-linking other classes of receptors (12-14). Even quantitative surface receptor expression of one receptor may be modified by cross-linking other receptors (15, 16).

Given these precedents for receptor interactions and the suggestion that cross-linking of Fc γ RIIIb with antireceptor mAb F(ab')₂ might augment the phagocytic activity of Fc γ RIIa that has been engaged independently (17), we sought evidence that Fc γ RIIIb can augment Fc γ RIIa-specific function and determined the basis for this effect. Taking advantage of the NA1-NA2 polymorphism as a model system to demonstrate that quantitative augmentation varies with differences in the primary structure of Fc γ RIIIb, we have shown that cross-linking of

Fc γ RIIIb directly leads to activation of Fc γ RIIa, a property that can be transferred in supernatants to other neutrophils and that can be blocked by inhibitors of reactive oxygen species and serine proteases. Through coordinate degranulation and generation of reactive oxygen species, Fc γ RIIIb elicits a cell program that can rapidly activate receptors on both the same and adjacent cells. This novel, oxidant-dependent interaction provides an efficient mechanism for amplification of Fc γ receptor function in neutrophils.

Methods

Subjects. Peripheral blood was collected from 32 disease-free volunteers who ranged in age from 20 to 56 yr (34 \pm 8 yr, mean \pm SD). Protocols for these studies were approved by the Institutional Committee on Human Rights in Research.

Determination of Fc γ RIIIb and Fc γ RIIa alleles. Determination of Fc γ RIIIb alleles, NA1 and NA2, was performed by leukoagglutination as described previously (5). The assignment of NA type was confirmed by immunoprecipitation and flow cytometry with mAbs CLB-FcR gran 1, CLB-gran 11, and GRM1 (18, 19). Phenotyping of donors for the LR-HR alleles of Fc γ RIIa was performed by quantitative flow cytometry using mAbs 41.H16 and IV.3 as described previously (20, 21). Phenotypic assignment was corroborated by anti-CD3 mitogenesis assays.

Reagents. HBSS, RPMI-1640, and IgG-free FCS were from GIBCO Laboratories (Grand Island, NY). FCS was heat inactivated at 56°C for 60 min. Phorbol dibutyrate (PDBu), catalase, superoxide dismutase (SOD), aminotriazole, *p*-aminobenzamidine (PABA), sodium azide, and *N*^α-tosyl-L-lysyl-chloromethyl ketone (TLCK) were purchased from Sigma Chemical Co. (St. Louis, MO). Sulfo-NHS-biotin, NHS-LC-biotin, and streptavidin were obtained from Pierce Chemical Co. (Rockford, IL).

Anti-Fc γ R mAbs IV.3 (anti-Fc γ RII, CD32 [22]), and 3G8 (anti-Fc γ RIII, CD16 [23]) IgG, Fab fragments, or F(ab')₂ were purchased from Medarex, Inc. (Annandale, NJ). Anti-CD16 mAb Leu11b (IgM) and anti-CD35 mAb E11 F(ab')₂ were obtained from Becton Dickinson & Co. (Mountain View, CA) and Research Diagnostics Inc. (Flinders, NJ), respectively. Anti-CD16 mAb 286.5 (24) was generously provided by Dr. Howard B. Fleit (Stony Brook, NY), and F(ab')₂ fragments were prepared by digestion with ficin according to the manufacturer's specifications (Pierce Chemical Co.). Silver stain analysis of SDS-PAGE gels of all Fab fragments and F(ab')₂ preparations indicated that there was no intact IgG. Purified human IgG2 myeloma proteins (hIgG2) were obtained from The Binding Site, Inc. (San Diego, CA). Anti-CD11b/CD18 mAb IB4 (IgG2a, anti-CD18) and IB4 F(ab')₂ were generously provided by Irene Graham (Washington University, St. Louis, MO) (25), and mAb MN41 (IgG1, anti-CD11b) was a gift from Jill Buyon (Hospital for Joint Diseases, New York) (26). Murine IgG1, isotype control for the flow cytometry studies, was obtained from Sigma Chemical Co. Phycoerythrin (PE)-conjugated goat anti-mouse IgG F(ab')₂ and FITC-conjugated rabbit anti-human IgG F(ab')₂ was purchased from Tago Immunochemicals (Burlingame, CA).

Preparation of cells. Fresh anticoagulated human peripheral blood was separated by centrifugation through a discontinuous two step Ficoll-Hypaque gradient (5). Neutrophils (PMNs) were isolated from the lower interface and washed with HBSS. Contaminating erythrocytes were lysed with hypotonic saline (0.02% NaCl) for 20 s, followed by 0.16% NaCl and a final wash with HBSS. After final washes, PMN were resuspended to 5 \times 10⁶ cells/ml.

Preparation of erythrocytes. Erythrocytes were coupled to IV.3 Fab (anti-Fc γ RII mAb) or human IgG2 myeloma protein by a biotin-avidin technique, as described previously (21, 27). To prepare biotinylated E (E_B), 0.5 ml of E (1 \times 10⁹ cells/ml) were incubated with sulfo-NHS-biotin (500 μ g/ml) for 20 min at 4°C, followed by three washes. E_B at 1 \times 10⁹/ml were incubated with an equal volume of streptavidin (250 μ g/ml) for 30 min at 4°C. The streptavidin coated E_B (E_{BA}) were then

washed and resuspended to 1 \times 10⁹ E/ml for immediate use. IV.3 Fab and human IgG2 were biotinylated with NHS-LC-biotin (0.01 mg biotin/mg protein) for 60 min at room temperature. To bind the biotinylated mAb to the E_{BA}, E_{BA} (12.5 μ l at 1 \times 10⁹/ml) were combined with 5 μ l of biotinylated protein (0.01–5 μ g) for 45 min (27). After three washes, the anti-Fc γ RII coated E_{BA} (E-IV.3) and the hIgG2 coated E_{BA} (E-hIgG2) were then resuspended in 125 μ l (1 \times 10⁸ E/ml) and used immediately. The density of opsonization of E-IV.3 and E-hIgG2 was standardized with flow cytometry for all experiments as described previously (21, 27). To assure maximal levels of internalization by neutrophils, densely opsonized E-IV.3 and E-hIgG2 (as determined by immunofluorescent flow cytometry) were used in the assays of phagocytosis.

Assay of phagocytosis. Quantitation of phagocytosis by PMN was performed as described previously (5, 21). Briefly, cells were resuspended in RPMI at 5 \times 10⁶ cells/ml. For Fc γ RIIIb-induced activation, the PMN were preincubated for 5 min with 3G8 F(ab')₂ (10 μ g/ml), which remained present throughout the assay of phagocytosis. Saturating concentrations of all anti-CD16, -CD35, or -CD18 F(ab')₂ fragments (as determined by flow cytometry) were used to cross-link PMN surface receptors in all experiments and were present throughout the assay. In the experiments with PDBu-treated PMN, PDBu (15 ng/ml) was added simultaneously with the phagocytic particle. For inhibition experiments the following reagents were added before the stimulus: SOD (150 μ g/ml), catalase (31,000 U/ml), azide (0.1%), PABA (10 mM), or TLCK (0.5 mM). Duplicate samples without inhibitors were the controls for each experimental condition.

To assess internalization of E target particles, PMN (100 μ l) were combined with E-IV.3 or E-hIgG2 (125 μ l). Because PMN from donors homozygous for the HR allele of Fc γ RIIa have minimal binding and internalization of E-hIgG2 (21), they were excluded from studies using this probe. The leukocyte-erythrocyte mixtures (ratios of 1:25) were centrifuged at 44 g for 3 min and then incubated at 37°C for 15 min to allow for maximum internalization. After hypotonic lysis of noninternalized E, phagocytosis was quantitated by light microscopy. At least 400 cells per slide were counted in duplicate. The data are expressed as phagocytic index (PI, number of ingested erythrocytes per 100 PMN).

To study the capacity of supernatants from stimulated PMN to activate resting cells, PMN were treated with 3G8 F(ab')₂ or PDBu for 5 min, washed twice, and cultured for 15 min at 37°C in RPMI. Supernatants were collected and combined with fresh PMN and target particles. After incubation for 15 min with or without catalase (31,000 U/ml) phagocytosis was quantitated by light microscopy.

Assay of attachment. To quantitate adherence of E target particles to PMN, cells were prepared and combined as described in the assay of phagocytosis above. After centrifugation at 44 g for 3 min, the PMN-erythrocyte mixtures were maintained at room temperature for 10 min and then gently resuspended. Adherence of E to PMN was quantitated by light microscopy. Data are expressed as attachment index (AI, number of adherent or internalized erythrocytes per 100 PMN). Under these conditions, < 1% of phagocytes had internalized E.

To distinguish the effect of activation stimuli on Fc γ RIIa-mediated internalization from any effect on Fc γ RIIa-mediated adherence, incubation mixtures with PMN-erythrocyte ratios of 1:1 were used. In preliminary studies, it was determined that under these conditions all erythrocytes were bound to PMN, thus eliminating the potential to increase phagocytosis by increasing adherence. Triplicate tubes were set up, with the first tube used to determine AI and document the absence of free (nonadherent) E by light microscopy. Under these conditions, with fixed and limited adherence, the AI for E-IV.3 and E-hIgG2 were 106 \pm 25 attached E/100 PMN and 102 \pm 30, respectively. The remaining replicates were incubated with or without activating agents [3G8F(ab')₂ (10 μ g/ml)] and/or inhibitors [SOD (150 μ g/ml)] at 37°C for 15 min and after lysis of noninternalized E, internalization was quantitated.

Immunofluorescent flow cytometry. Fresh leukocytes (5 \times 10⁵ in PBS with 0.1% BSA) were incubated with saturating amounts of specific mAb or isotype controls for 30 min at 4°C. After two washes with cold PBS containing 0.1% BSA, cells were incubated with saturating concentrations of PE-conjugated goat anti-mouse IgG F(ab')₂ for 30

min at 4°C, followed by washing twice with cold PBS/0.1% BSA. E coated with human IgG were stained with FITC-conjugated rabbit anti-human IgG F(ab')₂ and E coated with IV.3 Fab were stained with PE-conjugated goat anti-mouse IgG F(ab')₂, followed by washes with cold PBS/0.1% BSA.

After staining, cell-associated immunofluorescence was quantitated on a Cytofluorograf IIS (Becton Dickinson and Co., San Jose, CA) with a 2151 computer as described previously (5). For each experiment, the instrument was calibrated with quantitative FITC and PE microbead standards (Flow Cytometry Standards Corp., Research Triangle Park, NC) to allow for assessment of both absolute and relative levels of immunofluorescence.

Data analysis. For assessment of the relative phagocytic capacity of individuals with different FcγRIIIb alleles, all experiments were performed in a matched-pairs experimental design. Accordingly, each subject homozygous for a given FcγRIIIb allele was studied in comparison to a second subject, homozygous for the other allele and matched for the same phenotype of FcγRIIa (e.g., NA1-LR vs. NA2-LR).

The effects of activating stimuli on FcγR-mediated phagocytosis and attachment are presented as % control [$100 \times \text{experimental PI (or AI)}/\text{control PI (or AI)}$]. Data are displayed as mean ± SEM. The effects of stimuli or inhibitors were compared using a paired *t* test (two-tailed). A probability of 0.05 was used to reject the null hypothesis that there is no difference between the groups or conditions.

Results

Allele-sensitive differences in FcγRIIIb-induced activation of FcγRIIa. PMN from donors homozygous for the NA1 and NA2 alleles of FcγRIIIb in PMN differ in their capacities to mediate internalization of IgG-sensitized erythrocytes. To determine the relative roles of FcγRIIIb and FcγRIIa in contributing to this difference, we compared FcγRIIa-mediated internalization in NA homozygous donors and the capacity of the two alleles of FcγRIIIb to enhance FcγRIIa function. PMN from normal donors homozygous for NA1 and NA2 alleles and matched for identical FcγRIIa alleles were studied simultaneously in a matched-pairs design. In NA1 and NA2 homozygotes both baseline expression of FcγRIIa and FcγRIIIb and specific FcγRIIa-mediated phagocytosis are equivalent (21). Cross-linking of FcγRIIIb with 3G8 F(ab')₂ in donors homozygous for the NA1 allele markedly increased internalization of erythrocytes coupled to anti-FcγRII mAb IV.3 Fab (E-IV.3), an FcγRII-specific probe (21, 27) (Fig. 1 A; *P* < 0.006). In contrast, in donors homozygous for the NA2 allele, cross-linking of FcγRIIIb did not activate FcγRIIa-mediated internalization of E-IV.3 (NA1 vs. NA2, *P* < 0.005; *n* = 11). Cell surface expression of FcγRIIa did not change after FcγRIIIb cross-linking in either group (NA1 103 ± 2% control; NA2 101 ± 1% control; *n* = 4). Furthermore, the capacity to activate FcγRIIa function was not intrinsically different in individuals homozygous for either NA1 or NA2; PDBu-induced enhancement of E-IV.3 internalization (which is independent of FcγRIIIb) was identical for both groups (NA1 vs. NA2: 400 ± 306% control vs. 434 ± 212%, *n* = 6 pairs). These findings indicate that the capacity of FcγRIIIb to augment FcγRIIa function varies with FcγRIIIb structure and is a specific property of FcγRIIIb.

The suggestion of an FcγRIIIb-induced increase in the attachment index of E-IV.3 in NA1 individuals (Fig. 1 A) raised the possibility that the FcγRIIIb cross-linking might alter ligand binding capacity of FcγRIIa in addition to amplification of phagocytosis. Therefore, to examine FcγRIIIb-driven activation of FcγRIIa in the context of an FcγRIIa-specific natural ligand that might be more sensitive to changes in receptor avidity,




Figure 1. Cross-linking of FcγRIIIb activates FcγRIIa-mediated phagocytosis: allele-sensitivity. PMN from normal donors homozygous for NA1 and NA2 alleles and matched for identical FcγRIIa alleles were studied simultaneously in a matched-pairs design. PMN from each donor were preincubated with 3G8 F(ab')₂ (10 μg/ml) or control medium for 5 min and then combined with (A) E-IV.3 or (B) E-hIgG2. The effects of 3G8 F(ab')₂-induced activation on E-IV.3 phagocytosis and attachment is presented as % control [$100 \times \text{PI}_{3\text{G8 F(ab')}_2}$ (or AI)/ $\text{PI}_{\text{control}}$ (or AI)]. Control PI and AI for E-IV.3 were 114 ± 16 E-IV.3 internalized/100 PMN and 203 ± 32 E-IV.3 attached/100 PMN, respectively, and for E-hIgG2 105 ± 12 E-hIgG2 internalized/100 PMN and 152 ± 26 E-hIgG2 attached/100 PMN, respectively. Baseline AI and PI for NA1 and NA2 homozygotes were indistinguishable. (A) Cross-linking FcγRIIIb with 3G8 F(ab')₂ in donors homozygous for the NA1 allele increased internalization of E-IV.3 (NA1 vs. control *P* < 0.006), whereas in donors homozygous for the NA2 allele there was no increase in FcγRIIa-mediated internalization (NA1 vs. NA2 *P* < 0.005, *n* = 11 pairs). In NA1 homozygotes, there was a trend toward increased adherence of E-IV.3 in activated cells (NA1 vs. control 0.1 > *P* > 0.05; NA1 vs. NA2 *P* = NS; *n* = 6 pairs), but no increase in surface expression of FcγRII (*n* = 4). (B) In donors homozygous for the NA1 allele, cross-linking FcγRIIIb with 3G8F(ab')₂ amplified internalization of E-hIgG2 more than in NA2 donors (NA1 vs. NA2 *P* < 0.02; NA1 vs. control *P* < 0.001; NA2 vs. control *P* < 0.05, *n* = 12 pairs). Cross-linking of both the NA1 and NA2 alleles of FcγRIIIb resulted in comparable increments in attachment of E-hIgG2 (*n* = 5).

we explored the attachment and phagocytosis of erythrocytes coupled to human IgG2 (E-hIgG2). We and others have previously shown that E-hIgG2 do not bind FcγRIIIb but do bind efficiently to the LR (131-H) allelic form of FcγRIIa (21, 28). In PMN from homozygous NA1 individuals 3G8 F(ab')₂ preincubation significantly increased phagocytosis of E-hIgG2 (*P* < 0.001; Fig. 1 B). In contrast to the experiments with E-IV.3, however, NA2 donors also were capable of increasing FcγRIIa-mediated phagocytosis of E-hIgG2 (*P* < 0.05). When matched for FcγRIIa alleles, stimulated PMN from NA1 donors showed significantly greater increases in FcγRIIa function than PMN from NA2 (*P* < 0.02). The increase in E-hIgG2 internalization was greater than that in E-IV.3 for each group, which may reflect the clear increase in the attachment index for E-hIgG2 in both NA1 and NA2 donors (Fig. 1 B). Assessment of the percentage of cells supporting attachment demonstrated that the increase in attachment index reflects both a change in the percent attachment and the number of E-hIgG2 attached per cell.

To distinguish the effect of FcγRIIIb-mediated activation on attachment to FcγRIIa per se from its effect on internalization and to establish that both processes were indeed affected, internalization was assessed at limiting PMN-erythrocyte ratios. With a ratio of 1:1, all erythrocytes are bound to PMN,




Figure 2. Cross-linking CD16, but not CD35 or CD18, activates Fc γ RIIa. PMN were treated with saturating concentrations of the anti-CD16 mAbs Leu11b (IgM) or 286.5 F(ab')₂, anti-CD18 (IB4 F(ab')₂), anti-CD35 (E11 F(ab')₂), or control medium and incubated with either E-IV.3 or E-hIgG2. Phagocytosis was determined by light microscopy. Data are expressed as % control ($100 \times \text{PI}_{\text{stimulated}}/\text{PI}_{\text{control}}$). Internalization of both Fc γ RIIa-specific probes is enhanced by two distinct methods of cross-linking Fc γ RIIIb. Leu11b: E-hIgG2 $P < 0.02$, $n = 16$; E-IV.3 $P < 0.004$, $n = 12$; 286.5 F(ab')₂: E-hIgG2 $P < 0.02$, $n = 13$; E-IV.3 $P < 0.001$, $n = 16$; and E11 F(ab')₂ ($n = 8-9$) and IB4 F(ab')₂ ($n = 6-11$) $P = \text{NS}$.

which eliminates the potential to increase phagocytosis by increasing adherence. Using this system of controlled adherence (E-IV.3 AI: 106 ± 25 attached E/100 PMN and E-hIgG2 AI: 102 ± 30), treatment with 3G8 F(ab')₂ still resulted in a consistent increase in Fc γ RIIa-mediated phagocytosis (E-IV.3, $191 \pm 22\%$ control PI, $n = 5$, $P < 0.003$; E-hIgG2, $167 \pm 15\%$ control PI, $n = 6$, $P < 0.007$). Thus, cross-linking of Fc γ RIIIb resulted in both increased attachment and enhanced efficiency of internalization of E's attached to Fc γ RIIa (Fig. 1).

To exclude the possibility that activation of Fc γ RIIa might reflect a unique property of mAb 3G8, several different anti-CD16 mAbs of differing isotypes were studied. PMN from NA1 homozygotes or NA1-NA2 heterozygotes were treated with saturating concentrations of mAb 286.5 F(ab')₂ (a mIgG1 recognizing a ligand-binding site epitope) (24), Leu11b (a mIgM), or control medium. Internalization of both E-IV.3 and E-hIgG2 was significantly augmented by both of these anti-CD16 mAbs (Fig. 2). The lack of a requirement for an intact IgG heavy chain in the cross-linking antibody is underscored by experiments with the IgM anti-CD16 mAb Leu11b. Furthermore, the activation of Fc γ RIIa by intact Leu11b argues against the possibility that contaminating residual proteases from the digestion of IgG are the activators of Fc γ RIIa. In contrast to results with 3G8 F(ab')₂, 286.5 F(ab')₂, and Leu11b, incubation of PMN with F(ab')₂ fragments of anti-CD35 (E11) or anti-CD18 (IB4) mAbs did not alter Fc γ RIIa function (Fig. 2). These results emphasize the importance of CD16 cross-linking as a specific trigger for the activation of Fc γ RII.

Activation of Fc γ RIIa-mediated phagocytosis is oxidant dependent. Gresham et al. (29) have demonstrated that phorbol esters can stimulate neutrophil internalization of IgG-sensitized E by both Fc γ RIIa and Fc γ RIIIb through an oxidant-dependent mechanism. Given the ability of PDBu to augment Fc γ RIIa-

specific function, we evaluated the possibility that activation of Fc γ RIIa by PDBu and Fc γ RIIIb might use a similar mechanism. Internalization by Fc γ RIIa was unaffected by incubation with 0.5 mM H₂O₂, but in the presence of aminotriazole (20 mM), an inhibitor of catalase that can be released from endogenous neutrophil stores and metabolize H₂O₂, internalization was significantly increased by 0.5 mM H₂O₂ (E-IV.3 = $147 \pm 19\%$ control PI, $n = 11$, $P < 0.03$; E-hIgG2 = 133 ± 10 , $n = 10$; $P < 0.01$). Given these observations providing direct evidence that oxidants can affect receptor-specific function, we examined the effects of SOD and catalase, inhibitors of reactive oxygen metabolites. Neither SOD nor catalase altered baseline receptor-specific phagocytosis (Fig. 3). PDBu significantly augmented Fc γ RIIa-specific phagocytosis, an effect blocked by SOD and catalase (Fig. 3) in a manner similar to their effects on EA phagocytosis (29). In the presence of aminotriazole, PDBu resulted in an even greater enhancement of Fc γ RIIa-mediated internalization compared with PDBu alone (50% increase). Fc γ RIIIb-mediated phagocytosis was negligible at baseline and unaffected by phorbols (30). Most importantly, the increase in Fc γ RIIa phagocytosis induced by cross-linking Fc γ RIIIb was also inhibited in the presence of either SOD or catalase (Fig. 3). In contrast, heat-denatured and inactivated SOD did not block either PDBu- or Fc γ RIIIb-induced stimulation (PDBu with denatured SOD = $372 \pm 69\%$ control PI E-hIgG2; 3G8 F(ab')₂ with denatured SOD = $214 \pm 38\%$ control PI). These findings suggest that activation of Fc γ RIIa is mediated by generation of reactive oxygen intermediates and provide strong evidence that Fc γ RIIIb triggers their generation.

Reactive oxygen intermediates are found in cell supernatants, and therefore we considered the possibility that the erythrocyte probe might be modified by exposure to active supernatants and lead to the spurious conclusion that intrinsic Fc γ RIIa properties are being altered. Supernatants from PDBu-treated PMN were active in enhancing Fc γ RIIa-mediated binding and phagocytosis of unstimulated PMN (Fig. 4). As predicted, the activation induced by supernatants was oxidant dependent and blocked by catalase (Fig. 4). Similar results from experiments with supernatants from 3G8 F(ab')₂-stimulated PMN ($144 \pm 11\%$ control, $P < 0.04$) indicated that this effect was not simply a carry over of the stimulus per se from stimulated to unstimulated cells, but due to soluble factors in supernatants from activated cells. Having established the activity of stimulated supernatants, we tested their capacity to directly modify phagocytic probes. Treatment of both E-IV.3 and E-hIgG2 with supernatants from stimulated PMN did not alter their internalization ($107 \pm 8\%$ control and $97 \pm 2\%$ control, respectively), indicating not only that the oxidants are acting on the phagocytes rather than on the E's but also that oxidants may act in a paracrine fashion.

Because 3G8 F(ab')₂ pretreatment can induce an increase in both ligand binding and phagocytic efficiency, we examined the question of whether oxidants contributed to both effects. Inclusion of SOD in incubation mixtures of stimulated PMN abrogated the increase in attachment of E-hIgG2 (Fig. 5 A). Similarly, with limiting PMN/erythrocyte ratios, SOD blocked the augmentation of Fc γ RIIa-mediated phagocytosis of E-IV.3 or E-hIgG2 by Fc γ RIIIb cross-linking (Fig. 5 B). Thus both components of Fc γ RIIIb-stimulated phagocytosis by Fc γ RIIa are oxidant dependent.

"Auto-opsinization" does not mediate Fc γ RIIIb-induced augmentation of phagocytosis. We considered the possibility




Figure 3. Activation of Fc γ RIIa-mediated phagocytosis is oxidant dependent. PMN treated with PDBu (15 ng/ml), 3G8 F(ab')₂ (10 μ g/ml), or control medium and incubated with either E-IV.3 (A) or E-hIgG2 (B) in the presence of SOD (150 μ g/ml), catalase (31,000 U/ml), or buffer. Data are expressed as % control ($100 \times \text{PI}_{\text{stimulated}}/\text{PI}_{\text{control}}$). Activation of Fc γ RIIa by either PDBu and cross-linking Fc γ RIII is markedly decreased in the presence of inhibitors of reactive oxygen metabolites. PDBu-induced activation: E-IV.3: SOD vs. no SOD $P < 0.04$, $n = 5$; E-hIgG2: SOD vs. no SOD $P < 0.05$; cata-

lase vs. no catalase $P < 0.02$, $n = 5$. 3G8 F(ab')₂-induced activation: E-IV.3: SOD vs. no SOD $P < 0.02$; $n = 5$; catalase vs. no catalase $P < 0.04$, $n = 5$; E-hIgG2 internalization: SOD vs. no SOD $P < 0.02$; $n = 6$; catalase vs. no catalase $P < 0.01$, $n = 6$. PMN from NA1 homozygotes or NA1-NA2 heterozygotes were used in experiments with 3G8F(ab')₂.

that stimulated PMN might release the complement component C3, which becomes activated and bound to the erythrocyte probe ("auto-opsonization") (31, 32). Pretreatment of the probe with active supernatants did not confer enhanced phagocytosis. Nonetheless, we blocked the ligand-binding site for C3bi on complement receptor 3 with mAb MN41 IgG (26), but this did not prevent enhancement of Fc γ RIIa function (E-hIgG2 internalization by PDBu-stimulated PMN: 443% unstimulated PI, $n = 4$, $P < 0.03$). Recognizing that CD11b/CD18 may be activated by H₂O₂ (33) and that PDBu amplification of EA phagocytosis can be blocked by pretreatment of PMN with IB4 F(ab')₂ (34), an mAb against the β -chain of CR3 (25), we tested the effects of IB4 on Fc γ RIIIb-induced effects. As

shown in Fig. 6, internalization of E-IV.3 and E-hIgG2 was enhanced to a similar extent with or without IB4 F(ab')₂ (25 μ g/ml). Taken together these experiments do not support a role for CD11b/CD18 in the activation of Fc γ RIIa. However, a role for leukocyte response integrin or integrin-associated protein, which have the potential to amplify Fc γ R function in other systems (35-37), cannot be excluded.

Chlorinated oxidants and serine proteases participate in Fc γ RIIa activation. Proteolysis by serine proteases has been proposed as a mechanism for enhanced ligand binding by Fc γ RIIa in human monocytes (38, 39). Because PMN can use the H₂O₂-myeloperoxidase-chloride system to generate chlorinated oxidants capable of activating such protease zymogens and inactivating protease inhibitors (40-43), we considered the possibility that the generation of hypochlorous acid might be the mechanism underlying oxidant-dependent enhancement of Fc γ RIIa-mediated ligand binding. H₂O₂ is a substrate for the generation of HOCl, and O₂⁻ plays a role in regulating myeloperoxidase activity and may increase H₂O₂ availability by oxidizing other substrates (44, 45). While inhibition of activation by both catalase and superoxide dismutase is consistent with a role for HOCl, the more potent inhibition of Fc γ RIIa activation by catalase supports this model.

To examine the contribution of HOCl to oxidant-dependent Fc γ RIIIb-driven activation of Fc γ RIIa, we assessed the ability of methionine, a scavenger that rapidly reacts with HOCl to yield methionine sulfide (46), and sodium azide, an inhibitor of myeloperoxidase, to abrogate activation. Constitutive internalization of E-hIgG2 was not significantly altered by coincubation with 20 mM methionine ($131 \pm 13\%$ control), whereas 3G8 F(ab')₂-induced amplification of E-hIgG2 phagocytosis was significantly decreased ($P < 0.008$; Fig. 7). Methionine also inhibited the 3G8 F(ab')₂-induced increase of E-IV.3 phagocytosis ($P < 0.04$; Fig. 7) and decreased PDBu-induced enhancement (no methionine vs. methionine: 271 ± 55 vs. $141 \pm 13\%$ of unstimulated PI, $n = 6$, $P < 0.04$). Coincubation with sodium azide (0.1%) significantly decreased the Fc γ RIIIb-induced enhancement of Fc γ RIIa phagocytic function (E-hIgG2 $P < 0.01$; E-IV.3 $P < 0.001$, Fig. 7). The PDBu-induced activation also




Figure 4. Supernatants from stimulated PMN activate Fc γ RIIa. PMN were treated with PDBu (15 ng/ml) for 5 min, washed twice, and cultured for 15 min at 37°C in RPMI. Supernatants were collected and combined with fresh PMN and E-IV.3. After incubation for 15 min with or without catalase (31,000 U/ml) attachment and phagocytosis were determined ($n = 5-8$). Data are expressed as % control [$100 \times \text{PI}_{\text{stimulated}}$ (or AI)/ $\text{PI}_{\text{control}}$ (or AI)]. For AI: control vs. PDBu $P < 0.02$, PDBu vs. catalase $P < 0.04$. For PI: control vs. PDBu $P < 0.0001$, PDBu vs. catalase $P < 0.01$.




Figure 5. The increased efficiency of both Fc γ RIIa binding and internalization is oxidant dependent. (A) PMN in the presence or absence of SOD (150 μ g/ml) were treated with 3G8 F(ab')₂ (10 μ g/ml) or control medium and incubated with E-hIgG2 and attachment was assessed. Inclusion of SOD abrogated the Fc γ RIIb-driven amplification of E-IgG2 binding (SOD vs. no SOD: $P < 0.04$, $n = 5$). (B) To specifically assess phagocytosis, we fixed and limited attachment (1:1 ratios of E to PMN) as described in Methods. Rosetted PMN were incubated with 3G8 F(ab')₂ or

control medium, in the presence or absence of SOD. The 3G8 F(ab')₂-induced increase in the efficiency of Fc γ RIIa-mediated internalization was inhibited by SOD (no SOD vs. SOD: E-IV.3 $P < 0.03$, $n = 6$; E-hIgG2 0.1 $P > 0.05$, $n = 3$).

decreased 50% ($P < 0.01$). These data support a role for chlorinated oxidants that may interact directly with cellular targets to induce Fc γ RIIa activation or may facilitate the capacity of proteases to alter the receptor–ligand interactions.

A role for proteases in Fc γ RIIa function has been suggested by the observations that the serine protease inhibitor, TLCK, decreases Fc γ RIIa-mediated binding of EA in human monocytes (38, 39). As expected for the E-IV.3 probe, neither basal levels ($86 \pm 14\%$ control PI, $n = 6$) nor 3G8 F(ab')₂-stimulated levels of internalization (no TLCK vs. TLCK: 156 ± 10 vs. $138 \pm 12\%$ of unstimulated PI) were significantly affected by TLCK. In contrast, basal internalization of E-hIgG2, a probe sensitive to changes in ligand-binding site, was moderately decreased in the presence of 0.5 mM TLCK ($76 \pm 7\%$ control, $n = 5$, $P < 0.03$), while 3G8 F(ab')₂-stimulated enhancement was dramatically reduced from $217 \pm 22\%$ of unstimulated PI to $126 \pm 11\%$ ($n = 8$, $P < 0.02$). As further evidence for the role of inhibition of serine proteases, we performed a series of experiments in the presence of PABA, a specific reversible serine protease inhibitor. Similar to TLCK, PABA significantly reduced 3G8 F(ab')₂ activation of internalization of E-hIgG2 (no PABA vs. PABA: 239 ± 104 vs. $130 \pm 31\%$ of unstimulated PI, $n = 7$, $P < 0.01$), whereas the effect on E-IV.3 was minimal (no PABA vs. PABA: 211 ± 41 vs. $170 \pm 61\%$ of unstimulated

PI, $n = 7$, $P = \text{NS}$). Taken together, these observations suggest that activation-induced proteolytic effects contribute to augmentation of Fc γ RIIa function primarily through changes in binding detected by native ligand. Facilitation of this process by chlorinated oxidants provides a rapid mechanism by which neutrophils can use the Fc γ RIIb-activated NADPH oxidase system and granular constituents (47, 48) in a cooperative manner to modulate the binding capacity of Fc γ RIIa without altering quantitative receptor expression. The ability of methionine and azide to inhibit the activation-induced increase in E-IV.3 phago-




Figure 6. Anti-CD18 mAb IB4 F(ab')₂ does not inhibit activation of Fc γ RIIa. PMN were pretreated with IB4 F(ab')₂ (10 μ g/ml) before incubation with E-IV.3 or E-hIgG2. Internalization of E-IV.3 was enhanced to a similar extent with or without IB4 F(ab')₂ (control vs. PDBu $P < 0.02$; IB4 F(ab')₂ vs. IB4 F(ab')₂+PDBu, $n = 4$, $P < 0.004$). Similarly,

phagocytosis of E-hIgG2 could be amplified in the presence of IB4 F(ab')₂ [IB4 F(ab')₂ vs IB4 F(ab')₂+PDBu, $n = 6$, $P < 0.01$].




Figure 7. Chlorinated oxidants participate in Fc γ RIIa activation. PMN were incubated with the HOCl scavenger methionine (20 mM); sodium azide (0.1%), an inhibitor of myeloperoxidase; or buffer for 5 min at 37°C before activation with 3G8 F(ab')₂. Internalization of E-hIgG2 and E-IV.3 in stimulated and unstimulated cells was quantitated. Duplicate samples of activated cells without methionine (or azide) and unactivated cells with methionine (or azide) were the controls for each experimental condition. Activated phagocytosis of both Fc γ RIIa-specific probes was inhibited in the presence of methionine (E-hIgG2: no methionine vs. methionine, $n = 8$, $P < 0.008$; E-IV.3: no methionine vs. methionine, $n = 6$, $P < 0.04$). Similarly, azide inhibited activation of Fc γ RIIa (E-hIgG2: no azide vs. azide, $n = 7$, $P < 0.01$; E-IV.3: no azide vs. azide, $n = 9$, $P < 0.001$).

cytosis suggests that effects of oxidants, such as HOCl, extend beyond changes in ligand binding.

Discussion

Neutrophils from normal donors, homozygous for the two different allelic phenotypes of Fc γ RIIIb, have significantly different levels of Fc γ receptor-mediated phagocytosis (1). However, experiments using antireceptor mAb-targeted erythrocytes indicate that Fc γ RIIIb mediates phagocytosis poorly (2, 30) and raise the possibility that Fc γ RIIIb may collaborate with Fc γ RIIa on neutrophils for internalization of EA. Such an interactive mechanism would require that Fc γ RIIIb have the capacity to influence Fc γ RIIa-mediated phagocytosis and that this capacity demonstrates quantitative differences between donors homozygous for the two different Fc γ RIIIb alleles. Indeed, initial observations have suggested that cross-linking of Fc γ RIIIb with antireceptor mAb can augment the phagocytic activity of Fc γ RIIa on PMN (17). Therefore, we systematically explored both the capacity of Fc γ RIIIb to influence Fc γ RIIa in an allele-sensitive fashion and the mechanisms underlying this effect.

Our data indicate that cross-linking of Fc γ RIIIb does activate Fc γ RIIa for phagocytosis. Anti-CD16 mAbs of differing isotypes enhanced Fc γ RIIa function, demonstrating that priming is not a unique property of single mAb and is not dependent on IgG heavy chain. Furthermore, F(ab')₂ fragments of anti-CD35 or -CD18 mAbs did not alter Fc γ RIIa function, underscoring the importance of CD16 cross-linking as a specific trigger for the activation of Fc γ RII. The increase in Fc γ RIIa-specific internalization reflects both an increase in ligand-mediated binding of erythrocytes and an increase in internalization efficiency of targets bound. This activation does not involve Fc γ RIIIb serving as a ligand-binding, "capture" receptor delivering ligand directly to Fc γ RIIa (49–51), because Fc γ RIIIb does not bind hIgG2 although we recognize the possibility that this might occur with some human IgG1 and IgG3. The mechanism of activation is dependent on the Fc γ RIIIb-driven generation of oxidants, and at least part of the increase in avidity of Fc γ RIIa appears to rely on a proteolytic modification. In addition, an oxidant-dependent increase in Fc γ RIIa-mediated internalization also occurs. The capacity of supernatants from stimulated PMN to enhance Fc γ RIIa function in unstimulated PMN precludes the need for a physical association between Fc γ RIIa and Fc γ RIIIb. These oxidant-dependent mechanisms provide the opportunity not only for autocrine stimulation of Fc γ receptor function but also for paracrine stimulation of different cells and perhaps even different receptor species.

The mechanism of altered avidity involves, at least in part, proteolysis. Previous work by Van de Winkel et al. (38, 39) suggested that a proteolytic process was involved in Fc γ RIIa-mediated binding in monocytes. Although those studies did not formally exclude the possibility that a change in surface charge or other properties rather than direct receptor modification was involved, our studies indicating a change in avidity for ligand-driven binding of E-hIgG2 but not for mAb-driven binding of E-IV.3 support the concept of direct modification of Fc γ RIIa. Of course, modification of an as yet unrecognized accessory molecule participating in binding cannot be excluded. The ability of catalase and SOD to block this effect strongly implicates oxidants in this process, and the inhibition of Fc γ RIIa activation in the presence of methionine, a scavenger for HOCl, or azide, an inhibitor of myeloperoxidase, suggests that neutrophils can

use the H₂O₂-myeloperoxidase system to generate chlorinated oxidants, such as HOCl, which activate proteinase zymogens and inactivate proteinase inhibitors to begin a protease cascade. Because quantitative receptor expression did not change, the modulation of Fc γ RIIa is unlike that for other leukocyte receptors and counterreceptors, such as CD16, CD43, L-selectin, and TNF receptor, which regulate surface expression through enzymatic cleavage (15, 16, 52). Rather, conformational changes affecting the ligand binding site appear most likely.

In addition to a change in avidity, our data clearly demonstrate an increase in receptor-mediated internalization efficiency. Although the incomplete blockade of activation of TLCK or PABA could reflect either incomplete enzyme inhibition or the presence of several different mechanisms of activation, both the increase in E-IV.3 internalization in the absence of an increase in binding and the increase in phagocytosis in the limiting attachment paradigm unambiguously demonstrate an enhancement of the efficiency of receptor-mediated internalization. As with the change in avidity, this process involves the Fc γ RIIIb-mediated generation of oxidants and is inhibitable by both catalase and SOD. Given the rapid time frame for these effects, it is unlikely that modulation of transcription factors plays a significant role although oxidants are known to affect NF- κ B and others (53–55). In contrast, oxidants can rapidly influence levels of quantitative tyrosine phosphorylation, and tyrosine phosphorylation is essential for Fc γ receptor-initiated functions (56–58). Indeed, synergy between Fc γ R and Fc γ RIIIb for activation of the respiratory burst and phagocytosis requires tyrosine phosphorylation of Fc γ R (30, 59). Of course, changes in phosphorylation states could provide mechanisms for both avidity modulation ("inside-out" signaling [12–14]) and for internalization.

Our observations provide the basis for better understanding of the mechanisms whereby Fc γ RIIIb homozygous donors can differ in quantitative EA phagocytosis. More importantly, however, our data indicate a mechanism for receptor collaboration in an autocrine fashion without the requirement of either ligand-mediated or -independent direct physical interaction between receptor species. Of course, our results do not preclude a physical interaction of some fraction of Fc γ RIIIb and Fc γ RIIa, as suggested by resonance energy transfer studies (60, 61), nor do they preclude other molecules such as CR3 from playing a role in Fc γ RIIa priming (59); however, they do suggest a more general mechanism for Fc γ RIIIb that has the potential to influence or "collaborate" with a broader range of receptors (30) in both an autocrine and a paracrine fashion. Indeed, the target cells for a paracrine effect need not be restricted to other phagocytes, and thus these paracrine effects may provide a foundation for Fc γ receptor alleles on phagocytes to influence other components of the immune system.

In this context, these observations suggest that Fc γ R polymorphisms may be considered inherited disease susceptibility factors in host defense against infection and in the development of autoimmunity. For example, the current work showing that NA2 homozygotes are relatively ineffective in amplifying Fc γ RIIa is consistent with the observation that individuals homozygous for Fc γ RIIIb-NA2, especially when combined with homozygosity for Fc γ RIIa-HR and with a terminal complement component deficiency, are more likely to develop serious meningococcal infection (62). Interactions with other components of the immune system through modulation of protein tyrosine kinase activities, transcription factors, and other mechanisms remain to be explored.

Acknowledgments

We greatly appreciate the participation of the normal volunteers without whose assistance these studies could not have been completed. We thank Dr. Stephen J. Weiss and Dr. Jeffrey C. Edberg for useful suggestions, Dr. Howard Fleit and Dr. Jill Buyon for providing mAbs, Carl Triscari for assistance with flow cytometry, and Dr. Charles L. Christian for his continued support.

This work was supported in part by grants ROI-AR38889 (J. E. Salmon) and ROI-AR33062 (R. P. Kimberly) awarded by the National Institutes of Health and an award from the Gustavus and Louise Pfeiffer Research Foundation. The Flow Cytometry Core Facility at the Hospital for Special Surgery is supported in part by the Cornell Multipurpose Arthritis and Musculoskeletal Diseases Center (P60-AR38520).

References

1. Ravetch, J. V., and J. P. Kinet. 1991. Fc receptors. *Annu. Rev. Immunol.* 9:457-492.
2. Hogarth, P. M., M. D. Hulett, and N. Osman. 1992. Fc γ receptors: gene structure and receptor function. *Immunol. Res.* 11:217-225.
3. Schreiber, A. D., M. D. Rossman, and A. I. Levinson. 1992. The immunobiology of human Fc γ receptors on hematopoietic cells and tissue macrophages. *Clin. Immunol. Immunopathol.* 62:S66-S72.
4. van de Winkel, J. G. J., and P. J. A. Capel. 1993. Human IgG Fc receptor heterogeneity: molecular aspects clinical implications. *Immunol. Today* 14:215-221.
5. Salmon, J. E., J. C. Edberg, and R. P. Kimberly. 1990. Fc γ receptor III on human neutrophils. Allelic variants have functionally distinct capacities. *J. Clin. Invest.* 85:1287-1295.
6. Edberg, J. C., J. E. Salmon, and R. P. Kimberly. 1992. Functional capacity of Fc γ receptor III (CD16) on human neutrophils. *Immunol. Rev.* 11:239-251.
7. Takeshita, T., H. Asao, K. Ohtani, N. Ishii, S. Kumaki, N. Tanaka, H. Munakata, M. Nakamura, and K. Sugamura. 1992. Cloning of the γ chain of the human IL-2 receptor. *Science (Wash. DC)*. 257:379-382.
8. Taniguchi, T., and Y. Minami. 1993. IL2/IL2 receptor systems: a current overview. *Cell* 73:5-8.
9. Lopez-Casillas, F., J. L. Wrana, and J. Massague. 1993. Betaglycan presents ligand to the TGF β signaling receptor. *Cell* 73:1435-1444.
10. Hahn, W. C., Y. Rosenstein, V. Calvo, S. J. Burakoff, and B. E. Bierer. 1992. A distinct cytoplasmic domain of CD2 regulates ligand avidity and T-cell responsiveness to antigen. *Proc. Natl. Acad. Sci. USA*. 89:7179-7183.
11. Hahn, W. C., and B. E. Bierer. 1993. Separable portions of the CD2 cytoplasmic domain involved in signaling and ligand avidity regulation. *J. Exp. Med.* 178:1831-1835.
12. Gismondi, A., F. Mainiero, S. Morrone, G. Palmieri, M. Piccoli, L. Frati, and A. Santoni. 1992. Triggering through CD16 or phorbol esters enhances adhesion of NK cells to laminin via very late antigens. *J. Exp. Med.* 176:1251-1257.
13. Chatila, T., R. S. Geha, and M. A. Arnaout. 1989. Constitutive and stimulus-induced phosphorylation of CD11/CD18 leukocyte adhesion molecules. *J. Cell Biol.* 109:3435-3444.
14. Buyon, J. P., S. G. Slade, J. Reibman, S. B. Abramson, M. R. Phillips, G. Weissman, and R. Winchester. 1990. Constitutive and induced phosphorylation of the α - and β -chains of the CD11/CD18 leukocyte integrin family: relationship to adhesion-dependent functions. *J. Immunol.* 144:191-197.
15. Kishimoto, T. K., M. A. Jutila, E. L. Berg, and E. C. Butcher. 1989. Neutrophil Mac-1 and MEL-14 adhesion proteins are inversely regulated by chemotactic factors. *Science (Wash. DC)*. 245:1238-1241.
16. Bazil, V., and J. L. Strominger. 1994. Metalloprotease and serine protease are involved on cleavage of CD43, CD44, and CD16 from stimulated human granulocytes. *J. Immunol.* 152:1314-1322.
17. Salmon, J. E., N. L. Brogle, J. C. Edberg, and R. P. Kimberly. 1991. Fc γ receptor III induces actin polymerization in human neutrophils and primes phagocytosis mediated by Fc γ receptor II. *J. Immunol.* 146:997-1004.
18. Tetteroo, P. A. T., C. E. Van der Schoot, F. J. Visser, M. J. E. Bos, and A. G. E. Kr. Von dem Borne. 1987. Three different types of Fc γ receptors on human leukocytes defined by workshop antibodies: Fc γ R_{low} of neutrophils, Fc γ R_{low} of NK/K lymphocytes and Fc γ RII. *In Leucocyte Typing III*. A. J. McMichael, editor. Oxford Press, Oxford. 702-706.
19. Edberg, J. C., P. B. Redecha, J. E. Salmon, and R. P. Kimberly. 1989. Human Fc γ RIII (CD16). Isoforms with distinct allelic expression, epitopes displays and membrane anchors on PMN and NK cells. *J. Immunol.* 143:1642-1649.
20. Gosselin, E. J., M. F. Brown, C. L. Anderson, T. F. Zipf, and P. M. Guyre. 1990. The monoclonal antibody 41H16 detects the Leu 4 responder form of human Fc γ RII. *J. Immunol.* 144:1817-1822.
21. Salmon, J. E., J. C. Edberg, N. L. Brogle, and R. P. Kimberly. 1992. Allelic polymorphisms of human Fc γ receptor IIA and Fc γ receptor IIIB. Independent mechanisms for differences in human phagocyte function. *J. Clin. Invest.* 89:1274-1281.
22. Looney, R. J., D. H. Ryan, K. Takahashi, H. B. Fleit, H. J. Cohen, G. N. Abraham, and C. L. Anderson. 1986. Identification of a second class of IgG Fc receptors on human neutrophils. A 40 kilodalton molecule also found on eosinophils. *J. Exp. Med.* 163:826-836.
23. Fleit, H. B., S. D. Wright, and J. C. Unkeless. 1982. Human neutrophil Fc γ receptor distribution and structure. *Proc. Natl. Acad. Sci. USA*. 79:3275-3279.
24. Fleit, H. B., C. D. Kobasiuk, N. S. Peress, and S. A. Fleit. 1992. A common epitope is recognized by monoclonal antibodies prepared against purified neutrophil Fc γ RIII (CD16). *Clin. Immunol. Immunopathol.* 62:16-24.
25. Wright, S. D., P. E. Rao, W. C. Van Vorhees, L. S. Craigmyle, K. Iida, M. A. Talle, E. F. Westberg, G. Goldstein, and S. C. Silverstein. 1983. Identification of the C3bi receptor on human monocytes and macrophages by using monoclonal antibodies. *Proc. Natl. Acad. Sci. USA*. 80:5699-5703.
26. Diamond, M. S., S. C. Johnson, M. L. Dustin, P. McCaffery, and T. A. Springer. 1987. Differential effects on leukocyte functions of CD11a, CD11b, and CD18 mAb. *In Leucocyte Typing IV, White Cell Differential Antigens*. W. Knapp, editor. Oxford Press, Oxford. 570-574.
27. Edberg, J. C., and R. P. Kimberly. 1992. Receptor specific probes for the study of Fc γ receptor specific function. *J. Immunol. Methods*. 148:179-187.
28. Parren, P. W. H. I., P. A. M. Warmerdam, L. C. M. Boeijs, J. Arts, N. A. C. Westerdaal, A. Vlug, P. J. A. Capel, L. A. Aarden, and J. G. J. van de Winkel. 1992. On the interaction of IgG subclasses with the low affinity Fc γ RIIA (CD32) on human monocytes, neutrophils, and platelets. Analysis of a functional polymorphism to human IgG2. *J. Clin. Invest.* 90:1537-1546.
29. Gresham, H. D., J. A. McGarr, P. G. Shackelford, and E. J. Brown. 1988. Studies on the molecular mechanisms of human Fc receptor-mediated phagocytosis. Amplification of ingestion is dependent on the generation of reactive oxygen metabolites and is deficient in polymorphonuclear leukocytes from patients with chronic granulomatous disease. *J. Clin. Invest.* 82:1192-1201.
30. Edberg, J. C., and R. P. Kimberly. 1994. Modulation of Fc γ and complement receptor function by the glycosyl-phosphatidylinositol-anchored form of Fc γ RIII. *J. Immunol.* 152:5826-5835.
31. Ezekowitz, R. A. B., R. B. Sim, M. Hill, and S. Gordon. 1983. Local opsonization by secreted macrophage complement components. Role of receptors for complement in uptake of zymosan. *J. Exp. Med.* 159:244-260.
32. Borregard, N., L. Kjeldsen, K. Rygaard, L. Bastholm, M. H. Nielsen, O. W. Bejerrum, and A. H. Johnsen. 1992. Stimulus-dependent secretion of plasma proteins from human neutrophils. *J. Clin. Invest.* 90:86-96.
33. Skoglund, G., I. Cotgreave, J. Ricon, M. Patarroya, and M. Ingelman-Sundberg. 1988. H₂O₂ activates CD11b/CD18-dependent cell adhesion. *Biochem. Biophys. Res. Commun.* 157:443-449.
34. Gresham, H., I. L. Graham, D. C. Anderson, and E. J. Brown. 1991. Leukocyte adhesion-deficient neutrophils fail to amplify phagocyte function in response to stimulation. Evidence for CD11b/CD18-dependent and -independent mechanisms of phagocytosis. *J. Clin. Invest.* 88:588-597.
35. Gresham, H. D., J. L. Goodwin, P. M. Allen, D. C. Anderson, and E. J. Brown. 1989. A novel member of the integrin receptor family mediates Arg-Gly-Asp-stimulated neutrophil phagocytosis. *J. Cell Biol.* 108:1935-1943.
36. Brown, E. J., L. Cooper, T. Ho, and H. Gresham. 1990. Integrin-associated protein: a 50-kD plasma antigen physically and functionally associated with integrins. *J. Cell Biol.* 111:2785-2794.
37. Van Strijp, J. A. G., D. G. Russell, E. Tuomanen, E. J. Brown, and S. D. Wright. 1993. Ligand specificity of purified complement receptor type three (CD11b/CD18, $\alpha_m\beta_2$, (Mac-1)). *J. Immunol.* 151:3324-3336.
38. Van de Winkel, J. G. J., R. van Ommen, T. W. J. Huizinga, M. de Raad, W. B. Tuijnman, P. J. T. A. Groenen, P. J. A. Capel, R. A. P. Koene, and W. J. M. Tax. 1989. Proteolysis induces increased binding affinity of the monocyte type II receptor for human IgG. *J. Immunol.* 143:571-578.
39. Van de Winkel, J. G. J., M. Jansze, and P. J. A. Capel. 1990. Effect of protease inhibitors on human monocyte IgG Fc receptor II. Evidence that serine protease activity is essential for Fc γ RII-mediated binding. *J. Immunol.* 145:1890-1896.
40. Ossanna, P. J., S. T. Test, N. R. Matheson, S. Regiani, and S. J. Weiss. 1986. Oxidative regulation of neutrophil elastase- α -1-proteinase inhibitor interactions. *J. Clin. Invest.* 77:1939-1951.
41. Weiss, S. J. 1989. Tissue destruction by neutrophils. *N. Engl. J. Med.* 320:365-376.
42. Desrochers, P. E., and S. J. Weiss. 1988. Proteolytic inactivation of α -1-proteinase inhibitor by a neutrophil metalloproteinase. *J. Clin. Invest.* 81:1646-1650.
43. Desrochers, P. E., K. Mookhtiar, H. E. Van Wart, K. A. Hasty, and S. J. Weiss. 1992. Proteolytic inactivation of α_1 -proteinase inhibitor and α_2 -antichymotrypsin by oxidatively activated human neutrophil metalloproteinases. *J. Biol. Chem.* 267:5005-5012.
44. Thomas, E. L., D. B. Learn, M. M. Jefferson, and W. Weathered. 1988.

Superoxide-dependent oxidation of extracellular reducing agents by isolated neutrophils. *J. Biol. Chem.* 263:2178–2186.

45. Kettle, A. J., and C. C. Winterbourne. 1988. Superoxide modulates the activity of myeloperoxidase and optimizes the production of hypochlorous acid. *Biochem. J.* 252:259–263.

46. Test, S. T., and S. J. Weiss. 1986. The generation and utilization of chlorinated oxidants by human neutrophils. *Adv. Free Radical Biol. Med.* 2:91–116.

47. Crockett-Torabi, E., and J. C. Fantone. 1990. Soluble and insoluble immune complexes activate human neutrophil NADPH oxidase by distinct Fc γ receptor-specific mechanisms. *J. Immunol.* 145:3026–3032.

48. Huizinga T. W. J., K. M. Dolman, N. J. M. van der Lindne, M. Kleijer, J. H. Nuijens, A. E. G. Kr. von dem Borne, and D. Roos. 1990. Phosphatidylinositol-linked FcRIII mediates exocytosis of neutrophil granule proteins, but does not mediate initiation of the respiratory burst. *J. Immunol.* 144:1432–1437.

49. Anderson, C. L., L. Shen, D. M. Eicher, W. D. Wewers, and J. K. Gill. 1990. Phagocytosis mediated by three distinct classes of Fc receptors on human leukocytes. *J. Exp. Med.* 171:1333–1345.

50. Boros, P., J. A. Odin, T. Muryoi, S. K. Masur, and J. C. Unkeless. 1991. IgM anti-Fc γ R autoantibodies trigger neutrophil degranulation. *J. Exp. Med.* 173:1473–1482.

51. Huizinga, T. W. J., F. van Kemenade, L. Koenderman, K. M. Dolman, A. E. G. Kr. von dem Borne, P. A. T. Tetteroo, and D. Roos. 1989. The 40 kDa Fc γ receptor (FcRII) on human neutrophils is essential for the IgG-induced respiratory burst and IgG-induced phagocytosis. *J. Immunol.* 142:2365–2369.

52. Porteu, F., and C. Nathan. 1990. Shedding of tumor necrosis factor receptors by activated human neutrophils. *J. Exp. Med.* 172:599–607.

53. Schreck, R. P., P. Rieber, and P. A. Baeuerle. 1991. Reactive oxygen intermediates as apparently widely used messengers in the activation of the nF- κ B transcription factor and HIV-1. *EMBO (Eur. Mol. Biol. Organ.) J.* 10:2247–2258.

54. Satrianom, J. A., M. Schuldiner, K. Hora, Y. Xing, Z. Shan, and D. Schlondorff. 1993. Oxygen radicals as second messengers for expression of the monocyte chemoattractant protein, JE/MCP-1, and the monocyte colony-stimulating factor, CSF-1, in response to tumor necrosis factor- α and immunoglobulin G. Evidence for involvement of reduced nicotinamide adenine dinucleotide phosphate (NADPH)-dependent oxidase. *J. Clin. Invest.* 92:1564–1571.

55. Schenk, H., M. Klein, W. Erdbrugger, W. Droge, and K. Schulze-Osthoff. 1994. Distinct effects of thioredoxin and antioxidants on the activation of transcription factors NF- κ B and AP-1. *Proc. Natl. Acad. Sci. USA.* 91:1672–1676.

56. Huang, M.-M., Z. Indik, L. F. Brass, J. A. Hoxie, and A. D. Schreiber. 1992. Activation of Fc γ RII induces tyrosine phosphorylation of multiple proteins including Fc γ RII. *J. Biol. Chem.* 267:5467–5473.

57. Scholl, P. R., D. Ahern, and R. S. Geha. 1992. Protein tyrosine phosphorylation induced via the IgG receptors Fc γ RI and Fc γ RII in the human monocytic cell line THP-1. *J. Immunol.* 149:1751–1757.

58. Rankin, B. M., S. A. Yocum, R. S. Mittler, and P. A. Kiener. 1993. Stimulation of tyrosine phosphorylation and calcium mobilization by Fc γ receptor cross-linking. Regulation by the phosphotyrosine phosphatase CD45. *J. Immunol.* 150:605–616.

59. Zhou, M.-J., and E. J. Brown. 1994. CR3 (Mac-1, $\alpha_M\beta_2$, CD11b/CD18) and Fc γ RIII cooperate in generation of a neutrophil respiratory burst: requirement for Fc γ RIII and tyrosine phosphorylation. *J. Cell Biol.* 125:1407–1416.

60. Zhou, M.-J., R. F. Todd, J. G. J. van de Winkel, and H. R. Petty. 1993. Cocapping of the leukoadhesin molecules complement receptor type 3 and lymphocyte function-associated antigen-1 with Fc γ receptor III on human neutrophils: possible role of lectin-like interactions. *J. Immunol.* 150:3030–3041.

61. Kindzelskii, A. L., R. F. Todd, L. A. Boxer, and H. R. Petty. 1994. CR3 promotes inter-receptor proximity on neutrophils. *FASEB (Fed. Am. Soc. Exp. Biol.) J.* 8:501a.(Abstr.)

62. Fijen, C. A. P., R. G. M. Bredius, and E. J. Kuijper. 1993. Polymorphism of IgG Fc receptors in meningococcal disease. *Ann. Intern. Med.* 119:636.