

Table S1. Number of participants, deaths, follow-up duration, and death rates per 1,000 person-years for each eGFR and urinary ACR category, stratified by CHD status (January 2010 Data Freeze)

eGFR Categories	No baseline history of Coronary Heart Disease*				Baseline history of Coronary Heart Disease			
	ACR categories				ACR categories			
	<10	10 to <30	30 to 300	>300	<10	10 to <30	30 to 300	>300
90 to <120								
No.	3,765	2,330	683	81	643	572	181	34
Deaths	48	52	30	7	22	28	9	3
FU (years)	3.54	3.54	3.56	3.16	3.63	3.68	3.513	3.31
Death rate	3.60	6.30	12.35	27.34	9.43	13.29	14.15	26.67
60 to <90								
No.	3,212	1,893	674	101	823	693	307	71
Deaths	70	84	42	11	35	40	35	13
FU (years)	3.63	3.61	3.50	3.36	3.63	3.65	3.50	3.28
Death rate	6.00	12.29	17.80	32.43	11.72	15.80	32.55	55.84
45 to <60								
No.	336	252	180	50	128	143	106	34
Deaths	10	15	19	9	10	18	23	8
FU (years)	3.69	3.49	3.52	3.14	3.68	3.27	3.05	3.63
Death rate	8.07	17.07	30.01	57.32	11.72	38.49	71.11	64.79
30 to <45								
No.	79	87	75	27	33	41	53	23
Deaths	4	10	9	4	3	5	14	6
FU (years)	3.38	3.80	3.21	3.52	3.60	3.39	3.25	3.30
Death rate	14.96	30.26	37.40	42.04	25.25	36.00	71.11	79.15
15 to <30								
No.	8	8	15	39	9	13	16	27
Deaths	1	2	4	6	0	4	8	8
FU (years)	3.25	2.73	2.66	3.52	3.80	3.32	3.15	3.30
Death rate	38.48	91.57	100.42	45.56	0.00	92.79	158.8	89.66

Supplementary Table S1; Warnock et al, AJKD, Kidney Function, Albuminuria, and All-Cause Mortality in the REGARDS (Reasons for Geographic and Racial Differences in Stroke) Study”

Coronary heart disease defined as self-reported myocardial infarction or history of coronary artery procedure, or electrocardiographic evidence of a previous myocardial infarction; FU, follow time, expressed as median years. Death rates are expressed per 1,000 person-years.