

APPENDIX 2 (as supplied by the authors): Duration of the impact of herpes zoster pain on specific EQ-5D health domains and on the activities of daily living of 261 patients with herpes zoster

	Impaired at recruitment *	Duration (days) †		
	n	Median	95% CI	Q1 – Q3
Herpes zoster pain	261	32.5	30.5 – 44.0	23.5 – 80.0
Pain interference with activities of daily living (ZBPI)				
General activities	225	29.7	28.0 – 32.0	19.5 – 56.0
Mood	233	28.5	27.0 – 30.5	18.5 – 50.5
Walking ability	160	28.5	26.5 – 29.5	16.5 – 47.6
Normal work	210	29.0	28.0 – 31.5	19.5 – 50.0
Relations with others	196	27.0	22.6 – 29.0	14.5 – 48.0
Sleep	230	29.8	28.5 – 33.5	20.5 – 57.5
Enjoyment of life	231	28.8	27.0 – 30.5	20.5 – 50.5

Note: ZBPI = Zoster Brief Pain Inventory.

* Kaplan-Meier curves were restricted to subjects who reported impairment in specific health domains or pain-interference in specific activities of daily living at recruitment.

† Calculated from rash onset until the end of the pain interference in the activities of daily living, defined as the first of two consecutive questionnaires with herpes zoster pain interference = 0 before 30 days or herpes zoster pain interference = 0, 1, 2 after 30 days.