

Supplemental Information Table of Contents

Page	Content
S2	(A) Analytical RP-HPLC chromatogram of purified MAX8, (B) ESI (+) mass spectrum of purified MAX8
S3	Dynamic oscillatory rheology of 1.0 wt% MAX8 hydrogel at pH 7.4, 50 mM BTP, 150 mM NaCl, and 37°C alone and with encapsulated proteins: (A) Dynamic frequency sweep at 0.2% strain of (i) MAX8 alone, (ii) MAX8 encapsulating positively charged proteins: lysozyme and lactoferrin, (iii) MAX8 encapsulating neutral proteins: myoglobin and human IgG, and (iv) MAX8 encapsulating negatively charged proteins: α -lactalbumin and bovine serum albumin. (B) Dynamic strain sweep at 6 rad/sec of (i) MAX8 alone, (ii) MAX8 encapsulating positively charged proteins: lysozyme and lactoferrin, (iii) MAX8 encapsulating neutral proteins: myoglobin and human IgG, and (iv) MAX8 encapsulating negatively charged proteins: α -lactalbumin and bovine serum albumin.
S4	Dynamic oscillatory rheology of shear-thinned 1.0 wt% MAX8 hydrogel at pH 7.4, 50 mM BTP, 150 mM NaCl, and 37°C alone and with encapsulated proteins: (A) Dynamic frequency sweep at 0.2% strain of (i) MAX8 alone, (ii) MAX8 encapsulating small proteins: lysozyme, myoglobin, and α -lactalbumin (iii) MAX8 encapsulating human IgG. (B) Dynamic strain sweep at 6 rad/sec of (i) MAX8 alone, (ii) MAX8 encapsulating small proteins: lysozyme, myoglobin, and α -lactalbumin (iii) MAX8 encapsulating human IgG.





