

Type of file: table

Label: 1

Filename: supplemental table 4_9_10.doc

Supplemental Table Falls and fractures compared between active HT versus control.

Event	Estrogen Trial (ET)				p ¹	Estrogen plus Progestogen Trial (EPT)				p ²	HT Trials Combined				
	Active		Placebo			Active		Placebo			Active		Placebo		p
	N	%	N	%		N	%	N	%		N	%	N	%	
<u>Events occurring between 3 to 6 years</u>															
≥2 Falls per year															
No	424	97.7%	430	95.3%	0.06	498	97.3%	456	97.9%	0.55	922	97.5%	886	96.6%	0.28
Yes	10	2.3%	21	4.7%		14	2.7%	10	2.1%		24	2.5%	31	3.4%	
	median		median		0.005 ³	median		median		0.75 ³	median		median		0.03 ³
Number of falls per year	0		0.33			0		0			0		0		
Fractures (total)															
No	355	96.5%	368	94.9%	0.28	417	95.4%	357	94.7%	0.63	772	95.9%	725	94.8%	0.29
Yes	13	3.5%	20	5.2%		20	4.6%	20	5.3%		33	4.1%	40	5.2%	
<u>Events occurring between baseline and 6 years</u>															
≥2 Falls per year															
No	443	98.2%	457	96.6%	0.12	530	98.2%	458	97.5%	0.45	973	98.2%	915	97.0%	0.096
Yes	8	1.8%	16	3.4%		10	1.9%	12	2.6%		18	1.8%	28	3.0%	
	median		median		0.043 ³	median		median		0.16 ³	median		median		0.012 ³
# of falls per year	0.17		0.17			0.17		0.17			0.17		0.17		
Fractures (total)															
No	413	91.6%	422	89.4%	0.26	490	91.1%	403	85.6%	0.006	903	91.3%	825	87.5%	0.006
Yes	38	8.4%	50	10.6%		48	8.9%	68	14.4%		86	8.7%	118	12.5%	

¹ comparison between active and placebo for ET trial

² comparison between active and placebo for EPT trial

³ p-value obtained by nonparametric, Wilcoxon rank-sum test