

**Table S1. Oligonucleotide sequences**

TYPE/DESIGNATION	SEQUENCE
<b>EMSA oligonucleotides</b>	
m4 S3	CCGAAACTGTGGGAACCTGGTAGA
numb S1a	CTTTTTTTTCTCACACTTACCTT
numb S1b	CTTACCTTTTCCCACACCCCGC
numb S2	TGGGTCAGCGTGTGAAATACCTTT
numb S2mut	TGGGTCAGCGTGTCAAATACCTTT
numb S3	ACGAAGAGTTCCCACGACCACCGA
numb S4	CACTTCGGCGTGTGAAATTTCAA
<b>PCR primers for reporter gene constructs</b>	
numb 2.5-F	<b>ggggtacc</b> ccGGCTGCATTGTTTTCAAACGACAG
numb 2.5-R	ga <b>agatctt</b> cTGGAGTATGACACCTCGGTAATCC
numb CD1-F	ga <b>agatctt</b> cTTCTAATATTTGCATCCTGCCAGC
numb CD1-R	<b>ggggtacc</b> ccATTCTTCGATCTCGATCGTAGCT
numb CD2-F	ga <b>agatctt</b> cGATAGTATGATAGTGTATGTTT
numb CD2-R	<b>ggggtacc</b> ccTATCATCACTTCAATGGTTTTCC
<b>PCR primers for in situ hybridization probe</b>	
numb IS-F	gc <b>gaattc</b> GGATTACTTTCATGTCAGCGGAGAC
numb IS-R	ggc <b>gcgccgc</b> GTGGGTAACCTGGTAGCTTGTTCA
<b>PCR primers for CD2 enhancer mutagenesis</b>	
numb Su(H) S3m-F	ACGAAGAGTTGCCACGACCACCGA
numb Su(H) S3m-R	TCGGTGGTCGTGGCAACTCTTCGT
numb Su(H) S4m-F	CACTTCGGCGTGTCAAATTTCAA
numb Su(H) S4m-R	TTGAAAATTTGACACGCCGAAGTG
<b>Su(H) short hairpin oligonucleotides</b>	
Su(H)-NE-F	<b>ctagc</b> agtAACACTTTATCCTGTGGTCAtagttatattcaagcataTCACCGACAGCATAAAGTGTT <b>gcg</b>
Su(H)-NE-R	<b>aattc</b> gcAACACTTTATGCTGTGGTGAtagtcttgaatataactaTGACCGACAGGATAAAGTGTT <b>actg</b>

Terminal lowercase letters denote synthetic restriction sites (bold) used for cloning into the vector.