

Material Supplemental Table. Primers and restriction enzymes used in genotyping and RT-PCR.

	Forward	Reverse
Genomic DNA PCR		
ErbB2	5'-CGGAACCCACATCAGGCC-3'	5'-TTTCCTGCAGCCTACGC-3'
VgEcR	5'-TTTCGACGCAGCGTTACGAA-3'	5'-GTTGACTCATTATACGCC-3'
RXR α	5'-TGGCAGTACATCAAGTGTATCATA-3'	5'-AGCTGATGACCGAGAAAGG-3'
β -Gal	5'-CCCCCGCAGATCCCGG-3'	5'-GTTTTCCCAGTCACGACG-3'
I κ B α SR	5'-CCCCCGCAGATCCCGG-3'	5'-TGGACTACAAAGACGATGACGATAAA-3'
RT-PCR		
I κ B α SR	5'-ATGGACTACAAAGACGATGACGATAAA-3'	5'-CCAGGCCGCTGTCTGGC-3'
RPL-19	5'-AATGCTCGGATGCCTGAGAA-3'	5'-CTCCATGAGGATGCGCTTGT-3'
Real-time PCR		
MMP-3	5'-TGTCCCCTTCCATCTCTCTC-3'	5'-TGGTGATGTCTCAGGTTCCAG-3'
MMP-9	TGCCCATTTTCGACGACGAC-3'	5'-GTGCAGGCCGAATAGGAGC-3'
Acrp30	5'-AGCCGCTTATATGTATCGCTCA-3'	5'-TGCCGTCATAATGATTCTGTTGG-3'
PDGF-AA	5'-TGGCTCGAAGTCAGATCCACA-3'	5'-AGCCCCTACGGAGTCTATCTC-3'
ICAM-1	5'-GGCATTGTTCTCTAATGTCTCCG-3'	5'-GCTCCAGGTATATCCGAGCTTC-3'
Southern Blot		
DNA Fragment	Restrictive Enzyme	DNA Fragment Size
ErbB2	<i>PvuII</i>	1.4kb
VgEcR	<i>BamHI</i> & <i>XbaI</i>	2.0kb
β -Gal	<i>BamHI</i> & <i>XbaI</i>	4.0kb
RXR α	<i>EcoRI</i> & <i>HindIII</i>	1.9kb
I κ B α SR	<i>HindIII</i> & <i>XmaI</i>	1.2kb