

SUPPLEMENTARY DATA

Supplementary Table 1 Plasma damage marker concentrations of patients with diabetes mellitus according to albuminuria stratum.

	<i>Normo- albuminuria</i>	<i>Micro- albuminuria</i>	<i>Macro- albuminuria</i>	<i>p-value for trend in DM</i>
<u><i>Glomerular</i></u>				
- IgG (mg/ml)	6.3 (4.4-40.7)	21.8 (5.2-39.9)	7.0 (2.2-32.2)	0.80
<u><i>Proximal tubular</i></u>				
- KIM-1 (ng/ml))	0.05 (0.0-0.32)	0.16 (0.07-0.30)	0.19 (0.07-0.28)	0.88
- NAG (U/L)	31.8 (23.3-39.7)	32.6 (24.5-48.2)	23.2 (18.5-35.3)	0.19
- NGAL (ng/ml)	197 (117-323)	188 (131-288)	245 (193-430)	0.52
- CysC (mg/L)	0.79 (0.68-0.89)	1.00 (0.80-1.17)	1.30 (1.17-1.44)	<0.001
<u><i>Distal tubular</i></u>				
- H-FABP (ng/ml)	0.84 (0.36-2.63)	1.11 (0.00-2.53)	1.94 (0.47-17.08)	0.001

Values are given as median (25th-75th percentile)