

Supplemental Information

Table S1. Descriptive Information (Mean, Standard Deviation) of Sample, by Study Group

| | Controls N = 24 | Siblings N = 26 | VCFS N = 72 |
|----------------------------|----------------------------|----------------------------|------------------------|
| Gender, female (number, %) | 9 (37.5) | 12 (46.2) | 35 (48.6) |
| Age, Time 1 | 11.8 (1.9) | 12.2 (2.1) | 11.8 (2.1) |
| Age, Time 2 | 15.0 (1.9) | 15.2 (2.0) | 15.1 (2.1) |
| FSIQ, Time 1 | 94.5 (14.0) | 104.9 (14.6) | 70.9 (13.9) |
| FSIQ, Time 2 | 96.8 (11.5) | 104.1 (13.1) | 68.5 (14.9) |
| SOPS, Total | 3.0 (5.2) | 0.40 (0.8) | 6.4 (9.7) |
| SOPS, Positive Symptoms | 0.5 (1.4) | 0.04 (0.20) | 1.2 (2.7) |
| Years Between Timepoints | 3.1 (0.35) | 3.1 (0.43) | 3.3 (0.57) |

FSIQ, Full-scale IQ; SOPS, Scale of Prodromal Symptoms; VCFS, velocardiofacial syndrome




Figure S1. This graph depicts the Receiver Operating Characteristic (ROC) curve for a linear combination of neuroanatomic change scores (the logit score predictor) (excluding overall cranial gray and white matter) as a predictor of SOPS Prodromal Status at Time 2. SOPS, Scale of Prodromal Symptoms.