

The pivotal role of the β 7 strand in the intersubunit contacts of different human small heat shock proteins

Cell Stress and Chaperones

Evgeny V. Mymrikov, Olesya V. Bukach, Alim S. Seit-Nebi and Nikolai B. Gusev

Corresponding author: Nikolai B. Gusev, Department of Biochemistry,
School of Biology, Moscow State University, Moscow 119991, Russian Federation
Tel/Fax 7-495-939-2747; E-mail NBGusev@mail.ru

Online Resource 1

Table 1. Estimation of different elements of the secondary structure of human small heat shock proteins and their Cys-mutants from the data of far UV CD by CONTINLL subprogram of Sreerama and Woody (Sreerama and Woody 2000).

Structure Protein	α -Helix		β -strand		Turns	Unordered
	Regular	Distorted	Regular	Distorted		
Wild type α B-crystallin	0.02	0.05	0.23	0.13	0.23	0.35
Cys mutant of α B-crystallin	0.02	0.05	0.23	0.13	0.23	0.35
Wild type HspB6	0.03	0.06	0.23	0.13	0.22	0.34
Cys-mutant of HspB6	0.03	0.06	0.24	0.13	0.22	0.33
Wild type HspB8	0.02	0.05	0.23	0.13	0.22	0.35
Cys-mutant of HspB8	0.02	0.05	0.23	0.13	0.22	0.35