

Supporting Information for

Chemical Noise Reduction via Mass Spectrometry and Ion/Ion Charge Inversion: Amino Acids

Kerry M. Hassell[†], Yves C. LeBlanc[‡] and Scott A. McLuckey^{†*}

[†]Department of Chemistry
Purdue University
West Lafayette, Indiana, USA 47907-1393

[‡]AB SCIEX
71 Four Valley Drive
Concord, Ontario, Canada L4K4V8

Table of Contents

Figure S-1 – a) Isolated deprotonated arginine prior to ion/ion reaction. b) Post-ion/ion reaction positive ion mass spectrum after reaction of deprotonated arginine with the 10+ charge state of PAMAM generation 3.0.

Figure S-2 - Spectra of a mixture of amino acids (a) ionized as cations and (b) reacted with PAMAM 3.5 anions to convert the cations to anions. The peak indicated with * corresponds to a fragment ion known to be formed from the collision-induced dissociation of several of the amino acid cations present in (a).

Figure S-3 - Operational charge inversion efficiencies (see text for definition) for several amino acid cations in reaction with 1.5 [PAMAM-6H]⁶⁻ anions.

Figure S-4 - Response curves for histidine spiked in precipitated plasma. (a) Protonated histidine abundance versus concentration observed directly via positive nESI and (b) deprotonated histidine abundance versus concentration following charge inversion.




Figure S-1 – a) Isolated deprotonated arginine prior to ion/ion reaction. b) Post-ion/ion reaction positive ion mass spectrum after reaction of deprotonated arginine with the 10+ charge state of PAMAM generation 3.0.




Figure S-2 - Spectra of a mixture of amino acids (a) ionized as cations and (b) reacted with PAMAM 3.5 anions to convert the cations to anions. The peak indicated with * corresponds to a fragment ion known to be formed from the collision-induced dissociation of several of the amino acid cations present in (a).




Figure S-3. – Operational charge inversion efficiencies (see text for definition) for several amino acid cations in reaction with 1.5 [PAMAM-6H]⁶⁻ anions.




Figure S-4 – Response curves for histidine spiked in precipitated plasma. (a) Protonated histidine abundance versus concentration observed directly via positive nESI and (b) deprotonated histidine abundance versus concentration following charge inversion.