

Supporting Information for:

Enhanced Potency of Nucleotide-Dendrimer Conjugates as Agonists of the P2Y₁₄ Receptor: Multivalent Effect in G Protein-Coupled Receptor Recognition

Arijit Das,^a Yixing Zhou,^b Rhonda Carter,^b T. Kendall Harden,^b and Kenneth A. Jacobson^{a*}

Table of Contents

Figure S1. ¹ H NMR spectrum of compound 9 .	p. S2
Figure S2. ¹ H NMR spectrum of compound 12 .	p. S3
Figure S3. MALDI MS of PAMAM dendrimer complex 8 .	p. S4
Figure S4. MALDI MS of PAMAM dendrimer complex 9 .	p. S5
Figure S5. MALDI MS of PAMAM dendrimer complex 10 .	p. S6
Figure S6. MALDI MS of PAMAM dendrimer complex 11 .	p. S7
Figure S7. MALDI MS of PAMAM dendrimer complex 15 .	p. S8

Figure S1. ^1H NMR spectrum of PAMAM-UDPGA conjugate **9**.




Figure S2. ^1H NMR spectrum of PAMAM-UDPGA conjugate **12**.




Figure S3. MALDI MS of PAMAM dendrimer complex **8**.




Figure S4. MALDI MS of PAMAM dendrimer complex **9**.




Figure S4. MALDI MS of PAMAM dendrimer complex **10**.




Figure S6. MALDI MS of PAMAM dendrimer complex **11**.




Figure S7. MALDI MS of PAMAM dendrimer complex **15**.

