

FIGURE LEGENDS


Figure: Hazard Ratio for Incident PAD According to Time Since Smoking Cessation

Data are hazard ratios adjusted for age, hypertension, diabetes, hypercholesterolemia, body mass index, alcohol consumption and physical activity. Current smokers represent the reference group. Error bars represent 95 percent confidence intervals. PAD = Peripheral artery disease

Supplementary Figure: Cumulative Incidence for PAD According to Pack Years of Smoking

Cumulative incidence curves for symptomatic PAD according to never smokers, <10 pack years, 10-<30 pack years and ≥ 30 pack years of smoking exposure. PAD = Peripheral artery disease

Symptomatic Peripheral Arterial Disease



Patients at risk	Years							
	0	2	4	6	8	10	12	14
Never Smokers	20305	20224	20110	19934	19570	17992	9366	1037
<10 Pack-years	8597	8557	8502	8409	8252	7636	4036	480
10 to <30 Pack-years	7464	7407	7329	7223	7043	6438	6410	433
30+ Pack-years	2777	2731	2672	2597	2480	2216	1141	131

Supplementary Table Risk of incident peripheral arterial disease according to smoking status, classified into intermittent claudication only and undergoing invasive procedures, respectively

	Smoking Status				<i>P</i> trend*
	Never (n=20336)	Past (n=14263)	Current, <15 cigs/day (n=1967)	Current, ≥15 cigs/day (n=3259)	
Claudication Only (N=85)					
Age adjusted model (n=39825)	1.0 (Referent)	2.12 (1.25-3.93)	3.83 (1.53-9.61)	12.72 (7.12-22.71)	<0.001
Multivariable model 1 (n=39803) [‡]	1.0 (Referent)	2.22 (1.25-3.94)	4.03 (1.61-10.13)	12.59 (7.05-22.51)	<0.001
Multivariable model 2 (n=38960) [§]	1.0 (Referent)	2.32 (1.30-4.16)	3.45 (1.28-9.33)	12.37 (6.81-22.47)	<0.001
Peripheral Artery Revascularization (N=93)					
Age adjusted model (n=39825)	1.0 (Referent)	4.21 (2.13-8.34)	17.14 (7.95-36.99)	23.07 (11.62-45.79)	<0.001
Multivariable model 1 (n=39803) [‡]	1.0 (Referent)	4.25 (2.15-8.42)	18.24 (8.45-39.41)	23.34 (11.75-46.36)	<0.001
Multivariable model 2 (n=38960) [§]	1.0 (Referent)	4.70 (2.30-9.62)	19.25 (8.57-43.25)	25.64 (12.44-52.87)	<0.001

* *P* for trend across categories of cigarette exposure

† Incidence rates are per 1000 person-years of observation

‡ Additionally adjusted for history of hypertension, history of diabetes, and history of hypercholesterolemia

§ Additionally adjusted for body mass index, alcohol consumption, and physical activity