

Supplementary table S2. Integration with respect to transcriptional orientation and exons

<b>Set</b>	<b>Sites in genes</b>	<b>Sites in exons</b>	<b>Percent in exons</b>	<b>Sites in same orientation as transcription</b>	<b>Percent same orientation as transcription</b>
Pre-transplantation (liquid)	427	33	7.73	228	53.40
Pre-transplantation (methylcellulose)	219	13	5.94	112	51.14
Pre-transplantation (all)	641	46	7.18	336	52.42
Post-transplantation (uncultured)	103	8	7.77	50	48.54
Post-transplantation (methylcellulose)	23	0	0.00	13	56.52
Post-transplantation (all)	109	8	7.34	52	47.71
HIV MEFs	1745	106	6.07	860	49.28
Mouse lenti tumors	138	10	7.25	77	55.80
MRC pre-transplantation (liquid)	632	34	5.38	301	47.63
MRC pre-transplantation (methylcellulose)	370	23	6.22	193	52.16
MRC pre-transplantation (all)	997	56	5.62	490	49.15
MRC post-transplantation (uncultured)	239	8	3.35	114	47.70
MRC post-transplantation (methylcellulose)	75	4	5.33	35	46.67
MRC post-transplantation (all)	277	12	4.33	129	46.57
MRC HIV MEF	2492	127	5.10	1273	51.08
MRC mouse lenti tumors	247	13	5.26	133	53.85