

Table S3. Life history data used for generation time (G) estimates. Values of l_i (age-specific survival rates), b_i (birth rates), and p_i (probability of a gene being inherited from a parent of age i), for all age classes, i , used to calculate the mean age of breeding adults, G [see 55].

Age Class				Age Class			
i	l_i	b_i	p_i	i	l_i	b_i	p_i
1	1.000	0.0	0.0	18	0.071	14.0	0.091
2	0.682	0.0	0.0	19	0.064	12.0	0.071
3	0.514	0.0	0.0	20	0.059	15.0	0.081
4	0.409	0.0	0.0	21	0.054	12.0	0.059
5	0.337	0.0	0.0	22	0.049	13.0	0.059
6	0.284	0.0	0.0	23	0.045	14.5	0.060
7	0.244	0.0	0.0	24	0.041	15.0	0.057
8	0.212	0.0	0.0	25	0.038	13.0	0.045
9	0.185	0.0	0.0	26	0.035	15.0	0.048
10	0.164	0.0	0.0	27	0.032	13.0	0.038
11	0.145	0.0	0.0	28	0.029	12.0	0.032
12	0.130	0.0	0.0	29	0.027	12.0	0.030
13	0.116	0.0	0.0	30	0.025	11.0	0.025
14	0.105	4.0	0.038	31	0.023	12.0	0.025
15	0.095	5.0	0.043	32	0.021	12.0	0.023
16	0.086	8.5	0.067	33	0.020	12.0	0.021
17	0.078	12.0	0.086				