

Electronic Supplementary Materials for *Oecologia*

Integration of an invasive consumer into an estuarine food web: Direct and indirect effects of the New Zealand mud snail

Valance E. F. Brenneis^{*1,2}, Andrew Sih¹, and Catherine E. de Rivera²

¹*University of California, Davis. Department of Environmental Science & Policy
One Shields Avenue, Davis, CA 95616*

²*Portland State University. Department of Environmental Science & Management
P.O. Box 751, Portland, OR 97207*

*Corresponding author - Tel: (503) 725-4982 Fax: (503) 725-9040 E-mail:

valanceb@pdx.edu

Appendix A. Field survey site information, fish catch, and benthic invertebrate densities. Values for July, August, and September 2008 sampling events within a bay are averaged.

	Youngs Bay, OR	Baker Bay, WA
Latitude	46.10'16"	46.16'40"
Longitude	123.53'00"	124.01'00"
Temperature range	17.2 - 18.9°C	15.1 - 18.0°C
Salinity range	3.1 - 6.4 psu	8.1 - 14.0 psu
Conductivity	5.6 - 11.2 ms	14.0 - 23.1 ms
Total fish catch		
Pacific staghorn sculpin	10	10
Threespine stickleback	5,172	62
Shiner perch	1,997	84
Starry flounder	1	31
English sole	0	196
Benthic invertebrate density		
<i>P. antipodarum</i> (Mean # m ⁻² ± SE)	172.4 ± 33.7	1.6 ± 1.1
<i>Americorophium spp.</i>	3,880.1 ± 916.5	36.1 ± 14.4
<i>Eogammarus spp.</i>	8.02 ± 4.2	0
<i>Macoma spp.</i>	3.2 ± 1.9	8.0 ± 3.4
<i>Polycheate spp.</i>	111.4 ± 18.4	20.0 ± 10.6
<i>Mysid spp.</i>	7.2 ± 4.1	7.2 ± 5.6
Other gammarid amphipods	0	24.9 ± 12.4