


IgG standard	
IgG (ng/mL)	Abs 490/650 nm
64	1.197
	1.168
32	0.756
	0.764
16	0.439
	0.445
8	0.237
	0.232
4	0.12
	0.129
2	0.064
	0.069
1	0.051
	0.052



anti-monkey IgG coated plate		
day after trasnplantation	Fold dilution of plasma	Abs 490/650 nm
0	100000	0.994
		0.999
	500000	0.290
		0.290
33	100000	0.984
		1.014
	500000	0.271
		0.274
75	100000	0.996
		1.041
	500000	0.288
		0.294
89	100000	1.036
		1.004
	500000	0.280
		0.294
103	100000	1.129
		1.156
	500000	0.341
		0.346

MazF coated plate		
day after trasnplantation	Fold dilution of plasma	Abs 490/650 nm
0	10000	0.022
		0.006
	50000	0.006
		0.010
	100000	0.006
		0.010
33	10000	-0.001
		0.001
	50000	0.013
		0.007
	100000	0.003
		0.007
75	10000	-0.005
		-0.002
	50000	-0.001
		-0.004
	100000	0.012
		0.021
89	10000	0.009
		0.005
	50000	0.003
		0.003
	100000	0.000
		0.001
103	10000	0.018
		0.019
	50000	0.013
		0.017
	100000	0.007
		0.009
500000	10000	0.006
		0.008
	50000	-0.005
		-0.001
	100000	0.007
		0.004
500000	0.005	
	0.002	

Figure S1 Raw data of absorbance 490/650nm for Figure 3