

Nucleotide sequences of two genes encoding the light harvesting chlorophyll a/b binding protein of rice

Sheng Luan and Lawrence Bogorad

The Biological Laboratories, Harvard University, 16 Divinity Avenue, Cambridge, MA 02138, USA
Submitted February 3, 1989 EMBL accession nos X13908, X13909

Southern blots suggest that the rice genome contains about four genes coding for light harvesting chlorophyll a/b binding protein (LHCP). From a genomic library kindly provided by Dr. Susan Wessler at the University of Georgia, we have cloned and sequenced two of the four genes. Here we report the nucleotide sequences of the coding regions and some of 3' and 5' flanking regions of these genes, cab1R and 2R. The coding regions for mature proteins are very similar to one another and also to those of LHCP genes from wheat(1) and pea(2). The presumed transit sequences and flanking sequences of the two genes differ substantially.

1R GCG GGC TCG CAG ATC TTC AGC GAG GGC GGG CTC GAC TAC CTC GGC AAC CCG ACC CTG
 A G S Q I F S E G G L D Y L G N P S L
 2R GCG GGg TCG CAG ATC TTC AGC GAG GGC GGG CTC GAC TAC CTC GGC AAC CCG AGC CTG

1R ATC CAC GCG CAG AGC ATC CTC GCC ATC TGG GCG GTG CAG GTG GTG CTC ATG GGC GCC
 I H A Q S I L A I W A V Q V V L M G A
 2R ATC CAC GCG CAG AGC ATC CTC GCC ATC TGG GCG GTG CAG GTG GTG CTC ATG GGC GCC

1R GTC GAG GGG TAC CGG ATT GCT GGC GGG CCG CTC GGC GAG GTC GTC GAC CCG CTG TAC
 V E G Y R I A G G P L G E V V D P L Y
 2R GTC GAG GGG TAC CGG ATT GCc GGC GGG CCG CTC GGC GAG GTC GTC GAC CCG CTC TAC

1R CCC GGC GGC GCC TTC GAC CCG CTC GGC CTC GCC GAT GAC CCC GAG GCG TGC GCG GAG
 P G G A F D P L G L A D D P E A C/f A E
 2R CGg GGC GGC GCC TTC GAC CCG CTC GGC CTC GCC GAT GAC CCg GAG GCc TtC GGG GAG

1R CTC AAG GTG AAG AAG ATC AAG AAC GGC CGC CTC GCC ATG TTC TCC ATG TTC GGC TTC
 L K V K K/e I K N G R L A M F S M F G F
 2R CTC AAG GTG AAG gAG ATC AAG AAC GGC CGC CTC GCC ATG TTC TCC ATG TTC GGC TTC

1R TTC GTC CAG GCC ATC GTC ACC GGC AAG GGC CCC CTC GAG AAC CTC GCC GAC CAC CTC
 F V Q A I V T G K G P L E N L A D H L
 2R TTC GTC CAG GCC ATC GTC ACC GGC AAG GGC CCC CTC GAG AAC CTC GCC GAC CAC CTC

1R GCC GAC CCC GTC AAC AAC AAC GGC TGG GCC TAC GCC ACC AAC TTC GTC CCC GGC AAG
 A D P V N N N A W A Y A T N F V P G K
 2R GCC GAC CCC GTC AAC AAC AAC GGC TGG GCC TAC GCC ACC AAC TTC GTC CCC GGC AAG

1R TGA
 -
 2R TGA
 940
 1R AGTGGGGACCGTAGCTTAGCAGTGGTTAATTGTGGTGGATGGATTGTGGCCAGCGAGTTGGTGTCTTGGGT
 2R gcgccGccgCcgcgtgctgccatGgcgAcgcacgcgcGtc
 964
 1089
 1R TGGGGAAAGATGGTTAGTGGCAGGAGATGATGATCGAGTTGGTGGTGTACACTAAGAAGATGAAGAAGAAG
 2R TctGcAgGagaGtgTgcGTgtGtgtAcgcGgtGcagtAGaTGtacGggGgaTctgGccacgtcatctacAtgcc
 1113

Comparing cablR with 2R: The CAAT and TATA sequences are underlined. Small letters indicate amino acid/nucleotide difference. *** represents presumed deletions.

ACKNOWLEDGEMENTS

This research was supported in part by a research grant from the Rockefeller Foundation. S.L. is supported by K.C. Wong Education Foundation.

Reference:

1. Lamppa, GK, Morelli, G, Chua, NH (1985) Mol. Cell. Biol. 5,1370-1378
2. Cashmore, AR (1984) Proc. Natl. Acad. Sci. USA 81, 2960-2964