

## Nucleotide sequences of influenza A virus RNA segment 7: a comparison of five isolates

Suzanne L.Zebedee and Robert A.Lamb

Department of Biochemistry, Molecular Biology and Cell Biology, Northwestern University, Evanston, IL 60208-3500, USA

Submitted August 3, 1988

EMBL accession nos X08088-X08093 (incl.)

The nucleotide sequences of RNA segment 7 from influenza viruses A/WSN/33 (N), A/WS/33 (W), A/FW/1/50 (F), A/Singapore/1/57 (S), A/Port Chalmers/1/73 (P), and mouse-adapted A/Port Chalmers/1/73 (M) were determined directly from virion RNA by dideoxy primer-extension sequencing<sup>1</sup> and they are compared to that of A/Udorn/72 (U) determined previously<sup>2</sup>. Nucleotide numbering starts at residue 1 at 3' end of virion RNA segment 7. Untranslated regions, residues 1-25 and 1007-1027 were not determined. The table shows nucleotide differences; those causing coding changes are stippled. Amino acid changes in the M<sub>1</sub> and M<sub>2</sub> proteins between A/Udorn/72 and other strain(s) are also shown.

Ntd #	Strains						aa # & change	Ntd #	Strains						aa # & change	Ntd #	Strains						aa # & change		
	U	N	W	F	S	P			U	N	W	F	S	P			M	U	N	W	F	S	P		
31	c	t	t	t	t	c		331	a	g	g	g	g	a			785	c	t	t	t	c	c		
37	g	a	a	a	a	a		373	a	a	a	a	g	a			791	t	c	c	t	t	t		
55	t	c	c	c	t			394	a	a	a	a	a	g			794	t	c	t	t	t	t		
70	t	c	c	c	t			406	c	t	t	c	t	c			795	g	a	a	g	g	g	28	V-I"
94	g	g	g	g	g	a		427	g	g	g	g	g	a/g			800	t	a	a	c	c	t		
100	g	a	a	a	a	g		454	c	a	a	a	a	c			803	g	a	a	a	g	g		
124	t	a	a	t	t			463	g	g	g	a	g	g			805	g	a	a	g	g	g	31	S-N"
127	a	g	g	g	g	a		469	t	c	c	c	t	t			815	g	a	g	g	g	g		
136	a	c	c	c	c	a		502	c	t	t	t	t	c			819	t	t	t	c	t	t		
147	c	t	c	c	c	c/t41 A-V*		503	a	c	c	a	a	a			866	c	t	t	t	t	c		
169	a	a	a	a	a	t		514	g	g	g	g	g	a			872	a	t	t	t	c	a		
175	a	a	a	a	a	t		524	g	a	a	a	a	g	167	A-T*	873	t	c	c	c	t	t	54	P-L"
205	g	a	a	a	g	g		532	c	t	c	c	c	t			874	t	g	g	t	t	t	54	P-R"
211	t	t	t	c	t	t		544	a	c	c	a	a	a			879	g	a	a	a	g	g	56	E-K"
214	a	g	g	g	a	a		568	g	a	g	g	g	g			882	c	t	t	c	c	c	57	H-Y"
241	a	g	g	a	a	a		610	a	g	g	g	g	a			888	c	t	t	c	c	c		
274	c	t	t	c	c	c		619	g	a	a	a	a	g			917	a	g	g	a	a	a		
277	c	t	t	t	c	c		637	g	t	t	g	g	g	205	E-D*	920	t	a	a	a	t	t		
286	t	c	c	t	t	t		638	g	a	a	g	g	g	206	V-I*	958	g	a	a	g	g	g	82	S-N"
289	g	a	a	g	g	g		670	a	g	g	a	a	a			970	c	t	t	c	c	c	86	A-V"
309	g	a	a	g	g	g	95 R-K*	677	g	a	a	g	g	g	218	A-T*	977	c	t	t	t	c	c		
328	a	g	g	a	a	a		697	c	c	c	a	g	c	224	S-R*	978	a	g	g	a	a	a	89	S-G"
								712	a	a	a	g	a	a			991	g	a	a	g	g	g	93	S-N"

\* = M<sub>1</sub> aa; \*\* = M<sub>2</sub> aa; + = M<sub>2</sub> aa strain F; # = M<sub>2</sub> aa strains N & W

## REFERENCES

- Lamb, R.A. and Lai, C.-J. (1981) Virology 112, 746-751.
- Air, G.M. (1979) Virology 97, 468-472.
- Lamb, R.A., Lai, C.-J. and Choppin, P.W. (1981) Proc. Natl. Acad. Sci. USA 78, 4170-4174.