**Supplementary Table 1** - Socio-demographic and other characteristics according to participants' pediatric migraine status, Seattle and Tacoma, Washington, USA 1996-2008

	Pediatric Migraine	<sup>1</sup> No Migraine	
	N=220	N=3063	•
Characteristics	%	%	P-value
Age (years)*	31.2 ± 4.9	32.7 ± 4.5	< 0.001
<30	35.5	22.0	< 0.001
30-34	40.0	44.3	
35-39	20.9	27.2	
≥40	3.6	6.5	
Race/ethnicity			
Non-Hispanic White	92.3	85.9	0.008
Other	7.7	14.1	
Annual Household Income (\$)			
<30,000	5.0	3.5	0.38
30,000-69,999	22.3	19.6	
≥ 70,000	68.6	73.6	
Missing	4.1	3.3	
Nulliparous	64.1	63.4	0.83
Education ≤ high school	5.0	3.9	0.43
Unmarried	13.6	8.7	0.01
History of Diabetes Mellitus	2.3	1.3	0.21
History of Chronic Hypertension	8.6	4.0	< 0.001
Family History of Diabetes Mellitus	16.8	14.4	0.33
Family History of Hypertension	50.9	49.4	0.66
Currently Employed	81.8	81.3	0.84
Current Smoker	31.8	27.5	0.17
Multivitamin User	97.3	97.4	0.94
No Current Exercise	10.0	8.0	0.29
Exercise as teenager (13-17 years)			
Yes	94.1	93.8	0.71
No	4.1	4.8	
missing	1.8	1.4	

<sup>\*</sup>Mean ± standard deviation (SD)

<sup>&</sup>lt;sup>1</sup>NOTE: For this analysis, we excluded women with migraine a diagnosed as an adult (diagnosed ≥18 years of age; N=445) and those women with missing information for age at migraine diagnosis (N=5)

**Supplementary Table 2** - Comparison of participants' anthropometric characteristics according to pediatric migraine diagnosis, Seattle and Tacoma, Washington, USA 1996-2008

	Pediatric Migraine N=220	<sup>1</sup> No Migraine N=3063	Student's T-Test
Characteristics	Mean ± SD	Mean ± SD	P-value
Weight at age 18 (kg)	58.8 ± 12.2	57.4 ± 8.8	0.099
Adult Height (m)	1.67 ± 0.07	$1.67 \pm 0.07$	0.371
Body Mass Index at age 18 (kg/m²)	21.1 ± 3.9	20.7 ± 2.9	0.137
Pregravid Weight (kg)	69.9 ± 21.2	65.1 ± 13.6	0.001
Pregravid Body Mass Index (kg/m²)	25.0 ± 7.1	23.4 ± 4.7	0.001
Adult Weight Gain (kg)	11.1 ± 14.3	7.6 ± 10.4	0.001

Mean ± standard deviation (SD).

<sup>&</sup>lt;sup>1</sup>NOTE: For this analysis, we excluded women with migraine a diagnosed as an adult (diagnosed ≥18 years of age; N=445) and those women with missing information for age at migraine diagnosis (N=5)