

# Supporting Information

## Screening of 5-HT<sub>1A</sub> Receptor Antagonists using Molecularly Imprinted Polymers

Naphtali A. O'Connor<sup>†</sup>, David A. Paisner<sup>†</sup>, Donna Huryn<sup>‡</sup>, Kenneth J. Shea<sup>\*,†</sup>

<sup>†</sup> Department of Chemistry, University of California, Irvine, 92717

<sup>‡</sup> Department of Chemistry, University of Pennsylvania, 19104-6323

Ref. 28:

Fletcher, A.; Forster, E. A.; Bill, D. J.; Cliffe, I. A.; Hartley, J. E.; Jones, D. E.; McLenacha, A.; Stanhope, K. J.; Critchley, D. J.; Childs, K. J.; Middlefell, V. C.; Lanfumey, L.; Corradetti, R.; Laporte, A. M.; Gozlan, H.; Hamon, M.; Dourish, C. T., *Behav. Brain. Res.* **1996**, 73, 337-353.

**Table S1.** Data for binding isotherm of **P(WAY)**, **P(BA)** and **P(BL)**.

<b>P(WAY)</b>		<b>P(BA)</b>		<b>P(BL)</b>	
Bound (mM)	Free (mM)	Bound (mM)	Free (mM)	Bound (mM)	Free (mM)
0.39	1.47	0.29	1.57	0.31	1.55
0.36	1.27	0.28	1.35	0.29	1.34
0.30	0.92	0.24	0.98	0.24	0.98
0.24	0.76	0.18	0.82	0.19	0.81
0.22	0.56	0.18	0.6	0.18	0.6
0.19	0.38	0.15	0.42	0.15	0.42
0.14	0.26	0.11	0.29	0.12	0.28
0.09	0.14	0.07	0.16	0.08	0.15
0.07	0.07	0.054	0.086	0.057	0.083