## Supplementary Table 1. Correlations of measures of obstructive sleep apnea with glucose homeostasis

	Fasting Plasma Glucose (mg/dL)	1 hour glucose (mg/dL)	2-hour glucose‡ (mg/dL)	Hemoglobin A1C (%)
Apnea-Hypopnea Index‡	-0.018	-0.136	0.064	0.018
Arousal Index‡	0.102	-0.127	0.172	0.234 (p=0.069)
Lowest SaO <sub>2</sub> ‡	0.061	-0.020	-0.036	0.190

(All numbers represent correlation coefficients. Numbers in bold = significant association: \*=p<0.05,  $\dagger=p<0.01$ )

‡=Spearman Correlation Analysis

Glu 1 hour = Glucose level 1 hour after oral glucose ingestion on OGTT

Glu 2 hour = Glucose level 2 hours after oral glucose ingestion on OGTT

Lowest  $SaO_2$  = lowest oxyhemoglobin saturation

## Supplementary Table 2. Correlation of measures of obstructive sleep apnea with insulin secretory measures

	1-hour insulin‡ (microIU/mL)	2-hour insulin‡ (microIU/mL)	IGI‡	AIRg‡
Apnea-Hypopnea Index‡	0.059	0.146	0.059	0.079
Arousal Index‡	0.019	0.079	0.061	-0.067
Lowest SaO <sub>2</sub> ‡	-0.173	-0.15	-0.067	0.022

(All numbers represent correlation coefficients. Numbers in bold = significant association: \*=p<0.05,  $\dagger=p<0.01$ )

‡=Spearman Correlation Analysis

HOMA-B= Homeostasis model assessment of beta cell function

IGI= Insulinogenic index derived from OGTT

AIRg= Acute insulin response to glucose derived from IVGTT

Lowest  $SaO_2$  = lowest oxyhemoglobin saturation

## Supplementary Table 3. Correlation of measures of obstructive sleep apnea with insulin sensitivity parameters

	Fasting insulin	HOMA-IR‡	WBISI	SI
	(OGTT)‡			
	(microIU/mL)			
Apnea-Hypopnea Index‡	0.204	0.190	-0.16	-0.338*
Arousal Index‡	0.075	0.102	-0.149	-0.224
Lowest SaO <sub>2</sub> ‡	-0.205	-0.187	0.151	0.230

(All numbers represent correlation coefficients. Numbers in bold = significant association: \*=p<0.05,  $\dagger=p<0.01$ )

**‡**=Spearman Correlation Analysis

HOMA-IR=Homeostasis model assessment of insulin resistance

WBISI= Whole body insulin sensitivity index derived from OGTT

 $S_I$  = Insulin sensitivity index from the IVGTT

Lowest  $SaO_2$  = lowest oxyhemoglobin saturation