

Table S3. Distribution of organs testing negative and positive in limiting dilution assays used to generate the plot in Figure 3E of the main text.

| Mouse strain | Parasite strain | Liver | | | | Spleen | | | |
|---|---|------------|----------|------------------|-------------------------|------------|----------|------------------|-------------------------|
| | | LDA result | | Statistical test | | LDA result | | Statistical test | |
| | | Negative | Positive | Chi-square | Fisher's ^(a) | Negative | Positive | Chi-square | Fisher's ^(a) |
| C57/BL6 | wild type | 20% | 80% | | | 10% | 90% | | |
| | <i>mtxnpx</i> ⁻ | 86% | 14% | p<0.05 * | p<0.05 | 100% | 0% | p<0.05 * | p<0.05 |
| | <i>mtxnpx</i> ⁻ / <i>+mTXNPx</i> | 10% | 90% | | | 10% | 90% | | |
| B6.p47 ^{phox-/-} | <i>mtxnpx</i> ⁻ | 100% | 0% | p<0.05 * | p<0.05 | 75% | 25% | p<0.05 * | p<0.05 |
| | <i>mtxnpx</i> ⁻ / <i>+mTXNPx</i> | 0% | 100% | | | 0% | 100% | | |
| B6.RAG2 ^{-/-} IFN- γ ^{-/-} | wild type | 11% | 89% | | | n.a. | n.a. | - | - |
| | <i>mtxnpx</i> ⁻ | 89% | 11% | p<0.05 * | p<0.05 | | | | |
| | <i>mtxnpx</i> ⁻ / <i>+mTXNPx</i> | 11% | 89% | | | | | | |

^(a) Comparison between *mtxnpx*⁻ and wild type plus *mtxnpx*⁻/*+mTXNPx*; * more than 25 % of cells with expected values <5; n.a., not analyzed.