

Supplemental Table 1. Real time PCR primers.

Gene¹	Primer	Sequence²
EMR1	F	GCATAATCGCTGCTGGTTGA
	R	CCAGGCAAGGAGGACAGAGTT
MCP1	F	GCTGGAGAGCTACAAGAGGATCA
	R	CCTCTCTCTTGAGCTTGGTGACA
TNF α	F	AGACCCTCACACTCAGATCATCTTC
	R	CCTCCACTTGGTGGTTTGCT
IL6	F	GTCGGAGGCTTAATTACACATGTTC
	R	GAATTGCCATTGCACAACCTCTTT
IGF-I	F	CAGTTCGTGTGTGGACCGAG
	R	GCTCCGGAAGCAACACTCAT
IGFBP5	F	ATACAACCCAGAACGCCAGCT
	R	ACCTGGGCTATGCACTTGATG
IGFIR	F	GGCACAACTACTGCTCCAAAGAC
	R	CTTTATCACCACCACACTTCTG
PR	F	CAGAAGCCAGCCAGAGCC
	R	CACAGGTAAGCACGCCATAGT
ER α	F	TGCACCATTGACAAGAACCGGA
	R	AGCACCCATTTCAATTCGGCCT
Amphiregulin	R	AACTCCGCTGCTACCGCTGG
	R	TCCACCGGCACTGTGGTCCC
18S	F	GTGGGCCTGCGGCTTAAT
	R	GCGAGAGTCTCGTTCGTTATC

¹ Primers were designed to measure the abundance of egf-like module containing mucin-like hormone receptor-like sequence 1 (EMR1), monocyte chemoattractant protein 1 (MCP-1), TNF α , IL6, IGF-I, IGF binding protein 5 (IGFBP5), IGF-I receptor (IGFIR), progesterone receptor (PR), estrogen receptor (ER α), amphiregulin, and 18S ribosomal RNA (18S) transcripts.

² Primer sequences are shown in the 5' to 3' orientation.