

**Supplementary Table.** Adjusted mean (SE) mean arterial pressure (MAP), by trimester and history of migraine, after excluding women with incident gestational hypertension or preeclampsia

Migraine	<u>First Trimester</u>		<u>Second Trimester</u>		<u>Third Trimester</u>	
	Unadjusted Mean [SE]	Δ (95% CI)	Unadjusted Mean [SE]	Δ (95% CI)	Unadjusted Mean [SE]	Δ (95% CI)
NO	82.56 [0.13]		81.95 [0.11]		84.32 [0.13]	
YES	83.33 [0.33]	0.77 (0.1 to 1.4)	81.88 [0.29]	-0.07 (-0.6 to 0.5)	86.78 [0.31]	2.26 (1.17 to 2.8)
<b>MODEL 1</b>						
Migraine	* Adjusted Mean [SE]	Δ (95% CI)	* Adjusted Mean [SE]	Δ (95% CI)	* Adjusted Mean [SE]	Δ (95% CI)
NO	82.92 [0.16]		82.98 [0.10]		84.87 [0.13]	
YES	83.66 [0.33]	0.74 (0.4 to 1.4)	83.36 [0.29]	-0.11 (-0.7 to 0.4)	87.10 [0.31]	2.23 (1.7 to 2.8)
<b>MODEL 2</b>						
Migraine	** Adjusted Mean [SE]	Δ (95% CI)	** Adjusted Mean [SE]	Δ (95% CI)	** Adjusted Mean [SE]	Δ (95% CI)
NO	82.11 [0.16]		81.39 [0.11]		84.07 [0.13]	
YES	82.88 [0.33]	0.77 (0.2 to 1.4)	81.31 [0.27]	-0.08 (-0.6 to 0.4)	86.34 [0.28]	2.27 (1.7 to 2.8)

<sup>1</sup>Adjusted for age, race/ethnicity, parity, smoking and prenatal vitamin use. Adjusted mean values reported for MODEL 1 are from a regression model based on the entire study cohort (N = 3,373) with indicators variables specified to represent Non-Hispanic white, nulliparous, non-smoking women from 25-34 years of age.

<sup>2</sup>Adjusted for age, race/ethnicity, parity, smoking, prenatal vitamin use, and pre-pregnancy body mass index. Adjusted mean values reported for MODEL 2 are from a regression model based on the entire cohort (N = 3,373) with indicators variables specified to represent Non-Hispanic white, nulliparous, non-smoking women from 25-34 years of age, with a pre-pregnancy BMI < 25 kg/m<sup>2</sup>.