

Repetitive segmental structure of the transducin β subunit: Homology with the *CDC4* gene and identification of related mRNAs

(guanine nucleotide regulatory protein/photoreceptor biochemistry/molecular evolution/signal transduction)

HENRY K. W. FONG*, JAMES B. HURLEY*[†], ROSEMARY S. HOPKINS*, RYN MIAKE-LYE*, MARK S. JOHNSON[‡], RUSSELL F. DOOLITTLE[‡], AND MELVIN I. SIMON*

*Division of Biology, California Institute of Technology, Pasadena, CA 91125; and [†]Department of Chemistry, University of California, San Diego, La Jolla, CA 92093

Contributed by Melvin I. Simon, November 19, 1985

ABSTRACT Retinal transducin, a guanine nucleotide regulatory protein (referred to as a G protein) that activates a cGMP phosphodiesterase in photoreceptor cells, is comprised of three subunits. We have identified and analyzed cDNA clones of the bovine transducin β subunit that may be highly conserved or identical to that in other G proteins. From the cDNA nucleotide sequence of the entire coding region, the primary structure of a 340-amino acid protein was deduced. The encoded β subunit has a M_r of 37,375 and is comprised of repetitive homologous segments arranged in tandem. Furthermore, significant homology in primary structure and segmental sequence exists between the β subunit and the yeast *CDC4* gene product. The M_r 37,375 β subunit polypeptide is encoded by a 2.9-kilobase (kb) mRNA. However, there exists in retina other β -related mRNAs that are divergent from the 2.9-kb mRNA on the basis of oligonucleotide and primer-extended probe hybridizations. All mammalian tissues and clonal cell lines that have been examined contain at least two β -related mRNAs, usually 1.8 and 2.9 kb in length. These results suggest that the mRNAs are the processed products of a small number of closely related genes or of a single highly complex β gene.

The transduction of sensory signals involves a family of guanine nucleotide-binding proteins, referred to as N or G proteins (1, 2). Different G proteins are known to mediate the stimulation or inhibition of target biochemical processes in response to activation of specific receptors by light, hormones, neurotransmitters, and other chemical signals. Retinal transducin is a well-characterized G protein that activates a cGMP phosphodiesterase in photoreceptor rod cells and, like other G proteins, it is comprised of three subunits (3). The transducin α subunit (T_α) contains the guanine nucleotide-binding site and reversibly associates with a complex of the β and γ subunits (T_β and T_γ), two tightly bound polypeptides with reported M_r s of about 36,000 and 8400, respectively.

The structure and function of transducin has been studied by the analysis of cDNA clones for the γ and α subunits (4-10). The molecular cloning of cDNA for the α subunit of bovine transducin has revealed the presence of two closely related retina-specific proteins that possess significant sequence homology to each other, *ras* oncogene proteins and elongation factors of protein synthesis. T_γ and Ras proteins were also noted to have similar COOH-terminal sequences (4).

The expression of T_γ and T_α gene products appears to be retina specific (9, 10). On the other hand, the β subunits of transducin and of other G proteins from various tissues are structurally highly conserved on the basis of immunological

studies, amino acid composition, and proteolytic peptide mapping, suggesting that the β gene product may be ubiquitous (11-14). In addition, functional similarities between transducin β and the β subunit of other G proteins have been reported (15-17). However, two forms of the β polypeptide with M_r s of about 36,000 and 35,000 have been observed (18). To further define the structure and function of the β subunit we have isolated corresponding cDNA clones. In this paper we report and analyze the complete amino acid sequence of the β subunit and characterize the expression of β subunit mRNA in various tissues.

MATERIALS AND METHODS

Purification of T_β and Protein Sequence Analysis. Bovine transducin was isolated from rod outer segment membranes as described (19), and $T_{\beta\gamma}$ was purified by hydrophobic interaction chromatography on heptylagarose. It was partially cleaved with trypsin into fragments with M_r s of 27,000 and 14,000 and the purified M_r 27,000 tryptic peptide fragment of T_β was subjected to automated Edman degradation (20).

Preparation of Oligodeoxynucleotides, Isolation of cDNA, and DNA Sequence Analysis. Oligodeoxynucleotides were synthesized on an automated DNA synthesizer using phosphoramidite chemistry (21). The bovine retinal cDNA library in the λ gt10 phage vector was generously provided to us by Jeremy Nathans (22) and screened with mixed oligonucleotide probes as described (4). DNA sequencing was performed by the dideoxynucleotide chain-termination method (23). Synthetic oligonucleotides, 21 bases in length, were used as sequencing primers.

Sequence Searches and Comparisons. The bovine T_β sequence was searched against an updated version of the protein sequence collection known as NEWAT (24). The sequences and various segments of given proteins were subjected to a rigorous set of alignment and comparison procedures designed to assess the significance of relationships between them (25). In particular, alignments were made with an algorithm similar to that of Needleman and Wunsch (26) in concert with the weighted scoring system derived from the Dayhoff log-odds mutation matrix (27). Dot-matrix plots depicting similar segments were obtained with a program written by Mark S. Johnson.

Primer-Extended Hybridization Probe Synthesis. A cDNA probe was constructed by primer-extension (28) using universal primers and M13 DNA templates that contained the 1386-base-pair (bp) *EcoRI* fragment from the λ TB112 cDNA clone.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Abbreviations: T_α , T_β , and T_γ , α , β , and γ subunits of transducin; bp, base pair(s); kb, kilobase(s); G protein, a family of guanine nucleotide-binding proteins.

[†]Present address: Howard Hughes Medical Institute, University of Washington, Seattle, WA 98195.




Fig. 2. Homologous segments of the β subunit. The amino acid sequence of the β subunit was aligned as ≈ 43 -residue or ≈ 86 -residue (segments a-d) repeat units by using computer programs described previously (25, 26). Identical or highly conserved residues that align more than three times in the homologous segments are boxed. A consensus sequence including some of the most highly conserved residues is presented below the aligned sequence; hydrophobic residues are symbolized with a square. Numbering on the left indicates the amino acid positions.

of the 1.5-kb *EcoRI* fragment from isolate λ TB103 was determined by using synthetic oligonucleotide primers. The sequence of the first 185 nucleotides at the 5' end of the cDNA was obtained from two independent T_β cDNA clones, λ TB112 and λ TB116. The 5' sequences of the cDNA inserts in λ TB112 and λ TB116 overlapped the open reading frame found in λ TB103, which possessed a different 5' sequence. The 5' region of λ TB103 is probably an artifact of cloning, since, unlike the 5' cDNA region of λ TB112 and λ TB116, it is not present in any of the other T_β cDNA clones and does not hybridize to T_β mRNA in RNA transfer blot analysis (results not shown).

The NH_2 -terminal protein sequence of the M_r 27,000 T_β tryptic peptide was found to be precisely encoded within the cDNA sequence and is preceded by a basic residue, arginine, as expected for cleavage by trypsin. Thus, the DNA sequence in Fig. 1 encodes the β subunit of transducin. The open reading frame extends from the first ATG codon at nucleotide 95 until an in-phase termination signal, the TAA codon at nucleotide 1115. The protein is 340 amino acids long, and its M_r is 37,375, which is close to M_r 36,000, the estimated size for the transducin β subunit (3). Further, the amino acid composition agrees closely with the reported amino acid analysis of β subunits (11).

The cDNA sequence in Fig. 1 includes 94 bases of 5'-untranslated DNA but shows only 272 bases of 3'-untranslated DNA.

The *EcoRI* site at nucleotide 1381 was found in each independent T_β clone that was examined, and the cDNA sequence in Fig. 1 does not extend to the 3' terminus of the message.

Primary Structure of the β Subunit. At first glance, the amino acid sequence of the β subunit displayed several identical or highly conserved stretches of amino acids, which appeared to comprise larger segments of repeated homologous sequences. The best alignment by computer analysis resulted in a repeat unit of about 86 residues composed of two ≈ 43 amino acid segments (Fig. 2). A reiterated pattern of amino acid sequences was clear. A consensus pattern is shown in Fig. 2; residues Gly-13, His-14, Asp-36, Trp-42, and Asp-43 are highly conserved, as are the hydrophobic residues at consensus positions 11, 18, 29, 30, 31, and 41. Other identities and conserved substitutions are seen throughout the aligned sequences.

To look for other proteins that share sequence homology with T_β the NEWAT protein sequence database was searched. The only sequence found to show significant resemblance to T_β was the COOH-terminal portion of the yeast CDC4 gene product. The CDC4 protein is composed of 779 amino acids (32, 33). The full-length segment from amino acid 361 to 718 is shown aligned with the 340 amino acid sequence of T_β (Fig. 3). The authentic alignment score was +4.8 standard deviations above the mean of 64 jumbled comparisons of sequences of the same lengths and compo-




Fig. 3. Homology with yeast CDC4. The alignment of the CDC4 sequence with that of T_β was optimized and the appropriate gaps were introduced by computer analysis. The rows of sequences are arranged according to the segments of the β protein. Amino acid residues that are identical are boxed. Numbering on the right indicates the amino acid positions in CDC4.

sitions. Selected segments within these sequences scored even higher. Although the overall resemblance amounts to only 18–19% identity, there is little doubt that these proteins share common ancestry. The repetitive homologous segments within the CDC4 protein and the β subunit and the homology between these proteins emerge clearly from the dot-matrix analyses shown in Fig. 4.

Expression and Diversity of the β Subunit and Related mRNAs. Hybridization of a primer-extended cDNA probe to poly(A)⁺ RNA revealed two forms of mRNA, about 2.9 and 1.8 kb in length, in all bovine and mouse tissues that were examined (Fig. 5) as well as in cloned mouse, rat, and human cell lines (data not shown). To determine which of the two mRNA bands represents the T β cDNA clone that was isolated, the 305-bp *Bam*HI–*Eco*RI restriction fragment, which consisted primarily of 3'-untranslated sequences, was subcloned into the pUC-9 vector, labeled by nick-translation, and hybridized to retinal poly(A)⁺ RNA. In contrast to previous results using the full-length probe, only one major mRNA with the size of 2.9 kb hybridized to the 3'-region plasmid probe (Fig. 6, lane R).

To further characterize the different mRNAs, a series of sequencing primer probes was hybridized to replicate filter blots of size-fractionated retinal poly(A)⁺ RNA (Fig. 6, lanes B–Q). The probes hybridize throughout the length of the reported sequence to the small regions that are indicated in Fig. 1. The filters were washed successively at increasing temperature: first, 0–4°C, 55°C, and, then, 70°C. Only one mRNA hybridized to all T β -specific sequencing primers after stringent washing, and, therefore, this mRNA, about 2.9 kb in length, corresponds to the cDNA clone that was isolated. However, at least two other mRNA species, about 4.3 and 1.8 kb in length, also hybridized to a number of the probes. These same mRNA species were seen when a full-length hybridization probe was used (Fig. 6, lane A). The hybridizations occurred in a specific manner since only the primer probes 6, 12, 14, and 16 hybridized to the additional 4.3- and 1.8-kb mRNAs, whereas the 3'-specific probes 2 and 4 hybridized only to the 2.9-kb mRNA (results for primer 4 not shown). A control probe, 18, showed no hybridization to any of the mRNAs or to the relatively abundant 28S and 18S rRNAs. As the washing temperature was increased to 55°C, the oligonucleotide probes were first melted from the 1.8-kb transcript and washing at 70°C removed most of the remaining probe from the 4.3-kb mRNA. However, the entire series of probes remained hybridized to the 2.9-kb mRNA.

DISCUSSION

It is apparent that the β polypeptide consists of contiguous homologous segments, each about 43 amino acid residues in length. The consensus motif in the repeated segments of the




FIG. 5. β Subunit mRNA in bovine and mouse tissues. (A) Each lane contained 2 μ g of total poly(A)⁺ RNA from a mouse tissue. (B) Each lane contained 8 μ g of total poly(A)⁺ RNA from bovine tissue as indicated. A primer-extended probe was hybridized to the RNA blots at 42°C and washed at 50°C. At the right are indicated the positions and sizes of 28S, 23S, 18S, and 16S rRNA standards (in kilobases).

β protein include the less frequently found amino acid residues tryptophan and histidine at periodic intervals. Furthermore, the relative positions of specific cysteines and aspartate residues are maintained in each segment. The conserved pattern suggests that the protein structure has evolved by duplication and divergence from a basic \approx 43 amino acid sequence. Furthermore, the conserved features might be involved in a specific function of the protein, such as ligand or metal ion binding or an enzymatic activity. Repetitive homologous sequences have been found in a number of proteins, including the epidermal growth factor precursor (34–36) and the ribosomal S1 protein (37). Recently, the *Xenopus* 5S rRNA transcription factor TFIIIA, a zinc-binding protein (38), was noted to have a repeating segmented structure (39). The repeated segments are thought to form zinc-binding and DNA–interaction sites. The specific function of the conserved amino acids in the β subunit is not yet apparent.

The primary structure and segmental sequence of the β subunit bear striking resemblance with a portion of the yeast *CDC4* gene. It is quite evident that *CDC4* also has multiple copies of the conserved amino acid sequence motif that is found in the segments of the β subunit. In *CDC4* the consensus motif also appears to repeat periodically, indicating contiguous segments equal in size or slightly larger than the \approx 43-residue-long segments of the β subunit. The higher-order homologies such as the conserved consensus motif, the periodic repetition of the motif, as well as the overall sequence homology argue that the two proteins descended from a common ancestor. The β subunit and the *CDC4*




FIG. 4. Dot-matrix plots of bovine transducin β subunit sequence vs. itself (Left), transducin β subunit sequence vs. residue 361–718 of yeast *CDC4* (Center), and the same region of yeast *CDC4* vs. itself (Right). The analyses were performed with a moving 20-residue window (segment size = 20) and employed the weighted scoring system derived from the Dayhoff log-odds mutation matrix (27). The cut-off score for plotting a segment was 180 in each case. Note that the diagonal lines denoting similar segments occur at approximately 40- to 45-residue intervals in all three cases.




FIG. 6. Expression of diverse β subunit mRNA. Each lane contained 1 μ g of total poly(A)⁺ RNA from bovine retina. Lane A, hybridization to a primer-extended probe at lowered stringency. The blot was hybridized at 35°C and washed at 40°C in 0.3 M NaCl/0.03 M sodium citrate/0.1% NaDodSO₄ two times for 15 min. Lane R, hybridization to a nick-translated plasmid probe containing 305 bp of 3'-region cDNA. The filter was hybridized at 42°C and washed up to 52°C. Lanes B–Q, hybridization as indicated to sequencing primer probes 2, 6, 12, 14, 16, and 18. Oligonucleotide 18 hybridizes within the hatched region of the λ TB103 insert and was used as a control (see Fig. 1 Upper). Hybridization was performed at 50°C and washing was carried out successively at 0°C (lanes B–G), 55°C (lanes H–L), and 70°C (lanes M–Q). RNA size markers are indicated on the left and right.

protein may also share similar functions. *CDC4* is a yeast cell-division-cycle gene that is required in an early event in the mitotic cycle (40, 41). A primary activity of the β subunit is its interaction with the GTP-binding α subunits. In yeast the Ras proteins, which have regions of amino acid homology with α subunits (42), are thought to play a role analogous to that of the α subunit of the G_s protein—i.e., they activate adenylate cyclase in a GTP-dependent mechanism (43). Perhaps, just as α interacts with the β subunit in G proteins, Ras may interact with the *CDC4* gene product as part of an information-processing system required to integrate metabolic events with the yeast cell cycle. On the other hand, the β subunit may have a role in the control of the cell cycle in addition to its function as part of the G protein complex.

Our results also indicate that there is a diversity of β subunit-related mRNAs that could encode different proteins. It seems likely that there are a number of highly homologous β -related genes on the basis of hybridization with genomic DNA (results not shown); however, we cannot exclude the possibility that multiple cross-hybridizing messages arise from a single highly complex gene. Identification of β -related gene products and investigation of their role in the mechanisms involved in signal transduction and the regulation of cell growth await further study.

We express sincere thanks to Drs. Breck Byers and John Yochem for communicating to us the sequence of the yeast *CDC4* gene and Drs. Michael A. Lochrie for many helpful discussions and ideas, Jeremy Nathans for providing us with the bovine retinal cDNA library, Suzanna Horvath for synthesizing the oligodeoxynucleotides, David B. Teplow for assistance with the peptide sequence analysis, and Eric Phizicky for providing heptylagarose. This work was supported by Grant GM34236 from the National Institutes of Health.

- Rodbell, M. (1980) *Nature (London)* **284**, 17.
- Gilman, A. G. (1984) *Cell* **36**, 577–579.
- Stryer, L. (1985) *Biopolymers* **24**, 29–47.
- Hurley, J. B., Fong, H. K. W., Teplow, D. B., Dreyer, W. J. & Simon, M. I. (1984) *Proc. Natl. Acad. Sci. USA* **81**, 6948–6952.
- Yatsunami, K., Pandya, B. V., Oprlan, D. D. & Khorana, H. G. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 1936–1940.
- Lochrie, M. A., Hurley, J. B. & Simon, M. I. (1985) *Science* **228**, 96–99.
- Tanabe, T., Nukado, T., Nishikawa, Y., Sugimoto, K., Suzuki, H., Takahashi, H., Noda, M., Haga, T., Ichiyama, A., Kangawa, K., Minamino, N., Matsuo, H. & Numa, S. (1985) *Nature (London)* **315**, 242–245.
- Yatsunami, K. & Khorana, H. G. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 4316–4320.
- Medynski, D. C., Sullivan, K., Smith, D., Van Dop, C., Chang, F., Fung, B. K.-K., Seeburg, P. H. & Bourne, H. R. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 4311–4315.

- Van Dop, C., Medynski, D., Sullivan, K., Wu, A. M., Fung, B. K.-K. & Bourne, H. R. (1984) *Biochem. Biophys. Res. Commun.* **124**, 250–255.
- Manning, D. R. & Gilman, A. G. (1983) *J. Biol. Chem.* **258**, 7059–7063.
- Audiger, Y., Pantaloni, C., Bigay, J., Deterre, P., Bockaert, J. & Homburger, V. (1985) *FEBS Lett.* **189**, 1–7.
- Gierschik, P., Codina, J., Simons, C., Birnbaumer, L. & Spiegel, A. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 727–731.
- Huff, R. M., Axton, J. M. & Neer, E. J. (1985) *J. Biol. Chem.* **260**, 10864–10871.
- Kanaho, Y., Tsai, S.-C., Adamik, R., Hewlett, E. L., Moss, J. & Vaughan, M. (1984) *J. Biol. Chem.* **259**, 7378–7381.
- Northrup, J. K., Sternweis, P. C. & Gilman, A. G. (1983) *J. Biol. Chem.* **258**, 11361–11368.
- Bockaert, J., Deterre, P., Pfister, C., Guillon, G. & Chabre, M. (1985) *EMBO J.* **4**, 1413–1417.
- Sternweis, P. C. & Robishaw, J. D. (1984) *J. Biol. Chem.* **259**, 13806–13813.
- Fung, B. K.-K., Hurley, J. B. & Stryer, L. (1981) *Proc. Natl. Acad. Sci. USA* **78**, 152–156.
- Hunkapiller, M. W. & Hood, L. E. (1983) *Methods Enzymol.* **91**, 486–493.
- Hunkapiller, M., Kent, S., Caruthers, M., Dreyer, W., Firca, J., Giffin, C., Horvath, S., Hunkapiller, T., Tempst, P. & Hood, L. (1984) *Nature (London)* **310**, 105–111.
- Nathans, J. & Hogness, D. S. (1983) *Cell* **34**, 807–814.
- Sanger, F., Nicklen, S. & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 5463–5467.
- Doolittle, R. F. (1981) *Science* **214**, 149–159.
- Feng, D. F., Johnson, M. S. & Doolittle, R. F. (1985) *J. Mol. Evol.* **21**, 112–125.
- Needleman, S. B. & Wunsch, C. D. (1970) *J. Mol. Biol.* **48**, 443–453.
- Dayhoff, M. O., ed. (1978) *Atlas of Protein Sequence and Structure* (National Biomedical Research Foundation, Washington, DC), Vol. 5, Suppl. 3.
- Hu, N.-T. & Messing, J. (1982) *Gene* **17**, 271–277.
- Drubin, D. G., Caput, D. & Kirschner, M. W. (1984) *J. Cell Biol.* **98**, 1090–1097.
- Lehrach, H., Diamond, D., Wozney, J. M. & Broedtker, H. (1977) *Biochemistry* **16**, 4743.
- Thomas, P. S. (1980) *Proc. Natl. Acad. Sci. USA* **77**, 5201–5205.
- Peterson, T. A., Yochem, J., Byers, B., Nunn, M. F., Duesberg, P. H., Doolittle, R. F. & Reed, S. I. (1984) *Nature (London)* **309**, 556–558.
- Yochem, J. (1985) Dissertation (University of Washington, Seattle, WA).
- Gray, A., Dull, T. J. & Ullrich, A. (1983) *Nature (London)* **303**, 722–725.
- Scott, J., Urdea, M., Quiroga, M., Sanchez-Pescador, R., Fong, N., Selby, M., Rutter, W. J. & Bell, G. I. (1983) *Science* **221**, 236–240.
- Doolittle, R. F., Feng, D. F. & Johnson, M. S. (1984) *Nature (London)* **307**, 558–560.
- Doolittle, R. F., Woodbury, N. W. & Jue, R. A. (1982) *Biosci. Rep.* **2**, 405–412.
- Hanas, J. S., Hazuda, D. J., Bogenhagen, D. F., Wu, F. Y. H. & Wu, C. W. (1983) *J. Biol. Chem.* **258**, 14120–14125.
- Miller, J., McLachlan, A. D. & Klug, A. (1985) *EMBO J.* **4**, 1609–1614.
- Byers, B. & Goetsch, L. (1974) *Cold Spring Harbor Symp. Quant. Biol.* **38**, 123–131.
- Hartwell, L. H. (1971) *J. Mol. Biol.* **59**, 183–194.
- Hurley, J. B., Simon, M. I., Teplow, D. B., Robishaw, J. D. & Gilman, A. G. (1984) *Science* **226**, 860–862.
- Toda, T., Uno, I., Broach, J., Matsumoto, K. & Wigler, M. (1985) *Cell* **40**, 27–36.