


Appendix 2 (as submitted by authors): Multivariable analysis of venous (A) and arterial (B) thrombotic risk in third-generation combined oral contraceptive (COC) users compared with second-generation COC users. Rate ratio (RR) above 1.0 indicates increased risk. Second-generation COC users: *n* (number of thrombotic events) venous = 23, arterial = 41; *N* (woman-years of follow-up) venous = 33 187, arterial = 33 219; third-generation COC users: *n* (venous) = 384, *n* (arterial) = 382; *N* (venous) = 651 455, *N* (arterial) = 651 376. CI = confidence interval.




Appendix to Gronich N, Lavi I, Rennert G. Higher risk of venous thrombosis associated with drospirenone-containing oral contraceptives: a population-based cohort study. *CMAJ* 2011. DOI 10.1503/cmaj.110463.

Copyright © 2011 Canadian Medical Association or its licensors.

**B** Variable RR (95% CI)

3rd Generation	0.97 (0.70, 1.35)
Age	1.08 (1.07, 1.10)
Diabetes mellitus	1.61 (0.94, 2.74)
Hyperlipidemia	1.29 (0.94, 1.76)
Hypertension	2.46 (1.74, 3.48)
Cancer	1.48 (0.70, 3.15)
Smoking	1.26 (1.02, 1.57)
Obesity	1.45 (1.16, 1.81)
Length of use (months)	0.97 (0.96, 0.98)



Appendix to Gronich N, Lavi I, Rennert G. Higher risk of venous thrombosis associated with drospirenone-containing oral contraceptives: a population-based cohort study. *CMAJ* 2011. DOI 10.1503/cmaj.110463.

Copyright © 2011 Canadian Medical Association or its licensors.