

Table S2: Mean blood pressure at 8 and 32 weeks of age

8 weeks (N=7-12)

Genotype	Intervention	Mean	SD	<i>P vs WT-NT</i>	<i>P vs DD-NT</i>
WT	NT	91.08	8.99		
WT	LP	78.25	5.91	0.088	
WT	AP	89.94	8.45	1.000	
DD	NT	126.13	9.51	0.000	
DD	LP	100.90	9.10	0.225	0.000
DD	AP	106.78	6.01	0.036	0.001
DD/ <i>Ncf1</i> -	NT	106.07	9.77	0.001	0.000
DD/ <i>Ncf1</i> -	LP	98.00	9.54	0.371	0.000
DD/ <i>Ncf1</i> -	AP	98.87	6.81	0.260	0.000

32 weeks (N=3-7)

Genotype	Intervention	Mean	SD	<i>P vs WT-NT</i>	<i>P vs DD-NT</i>
WT	NT	93.71	6.58		
WT	LP	81.06	1.88	0.013	
WT	LN	91.36	6.81	1.000	
WT	AP	95.84	7.95	1.000	
WT	AN	87.30	1.65	0.595	
DD	NT	138.15	15.94	0.000	
DD	LP	93.21	15.44	1.000	0.001
DD	LN	102.30	3.90	0.064	0.004
DD	AP	101.68	13.62	0.709	0.010
DD	AN	101.95	6.72	0.227	0.005
DD/ <i>Ncf1</i> -	NT	104.67	9.88	0.109	0.000
DD/ <i>Ncf1</i> -	LP	96.60	6.41	1.000	0.000
DD/ <i>Ncf1</i> -	LN	91.27	7.05	1.000	0.000
DD/ <i>Ncf1</i> -	AP	96.03	15.45	1.000	0.000
DD/ <i>Ncf1</i> -	AN	86.69	9.56	0.727	0.000