SUPPLEMENTAL MATERIAL

Supplemental Table 2A. Reclassification Tables for Clinical Model + UACR vs Clinical Model

		Clinical Model + UACR				
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model	<5%	21 (80.8%)	3 (11.5%)	2 (7.7%)	26	19.2
N (%)	5-9.99%	5 (9.8%)	34 (66.7%)	12 (23.5%)	51	33.3
	≥10%	0 (0%)	8 (14.5)	47 (85.5%)	55	14.5
Total		<u> </u>			132	

		Cl	Clinical Model + UACR			
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model	<5%	889 (89.0%)	109 (10.9%)	1 (0.1%)	999	11.0
N (%)	5-9.99%	97 (13.8%)	532 (75.9%)	72 (10.3%)	701	24.1
	≥10%	0 (0%)	85 (17.9%)	389 (82.1%)	474	17.9
Total					2,174	

Supplemental Table 2B. Reclassification Tables for Clinical Model + HbA1c vs Clinical Model

		С	Clinical Model + HbA1c			
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	18 (69.2%)	8 (30.8%)	0 (0%)	26	30.8
N (%)	5-9.99%	5 (9.8%)	35 (68.6%)	11 (21.6%)	51	31.4
	≥10%	0 (0%)	6 (10.9%)	49 (89.1%)	55	10.9
Total					132	

		Cl	Clinical Model + HbA1c			
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model	<5%	873 (87.4%)	126 (12.6%)	0 (0%)	999	12.6
N (%)	5-9.99%	112 (16.0%)	474 (67.6%)	114 (16.4%)	701	32.4
	≥10%	3 (0.6%)	102 (21.5%)	369 (77.8%)	474	22.1
Total					2,174	

Supplemental Table 2C. Reclassification Tables for Clinical Model + HbA1c + UACR vs Clinical Model

Stroke		Clinical Model + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	18 (69.2%)	7 (26.9%)	1 (3.8%)	26	30.7
N (%)	5-9.99%	10 (19.6%)	28 (54.9%)	13 (25.5%)	51	45.1
	≥10%	0 (0%)	10 (18.2%)	45 (81.8%)	54	18.2
Total		·			132	

No Stroke		Clinical Model + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	869 (87.0%)	129 (12.9%)	1 (0.1%)	999	13.0
N (%)	5-9.99%	142 (20.3%)	447 (63.8%)	112 (16.0%)	701	36.3
	≥10%	5 (1.1%)	112 (23.6%)	357 (75.3%)	474	24.7
Total					2,174	

Supplemental Table 2D. Reclassification Tables for Clinical Model + HbA1c + UACR vs Clinical Model + HbA1c

Stroke		Clinical Model + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model +	<5%	19 (82.6%)	4 (17.4%)	0 (0%)	23	17.4
HbA1c	5-9.99%	9 (18.4%)	34 (69.4%)	6 (12.2%)	49	30.6
N (%)	≥10%	0 (0%)	7 (11.7%)	53 (88.3%)	60	11.7
Total		·			132	

No Stroke		Clinical Model + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	937 (94.8%)	51 (5.2%)	0 (0%)	988	5.2
HbA1c	5-9.99%	79 (11.3%)	565 (80.5%)	58 (8.3%)	702	19.6
N (%)	≥10%	0 (0%)	72 (14.9%)	412 (85.1%)	484	14.9
Total					2,174	

Supplemental Table 2E. Reclassification Tables for Clinical Model + LA vs Clinical Model

Stroke			Clinical Model + LA			
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	23 (92.0%)	2 (8.0%)	0 (0%)	25	8.0
N (%)	5-9.99%	7 (13.2%)	34 (64.2%)	12 (22.6%)	53	35.8
	≥10%	0 (0%)	4 (6.8%)	55 (93.2%)	59	6.8
Total					137	

No Stroke			Clinical Model + LA			
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	883 (86.5%)	138 (13.5%)	0 (0%)	1021	13.5
N (%)	5-9.99%	113 (13.2%)	527 (72.3%)	89 (12.2%)	729	27.7
	≥10%	0 (0%)	84 (17.4%)	400 (82.6%)	484	17.4
Total					2,234	

Supplemental Table 2F. Reclassification Tables for Clinical Model + MAC vs Clinical Model

Stroke			Clinical Model + MAC			
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model	<5%	25 (100%)	0 (0%)	0 (0%)	25	0
N (%)	5-9.99%	4 (7.5%)	42 (79.2%)	7 (13.2%)	53	20.7
	≥10%	0 (0%)	2 (3.4%)	57 (96.6%)	59	3.4
Total		<u> </u>			137	

No Stroke		(
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model	<5%	985 (96.5%)	36 (3.5%)	0 (0%)	1021	3.5
N (%)	5-9.99%	37 (5.1%)	655 (89.8%)	37 (5.1%)	729	10.2
	≥10%	1 (0.2%)	45 (9.3%)	439 (90.7%)	484	9.3
Total		·			2,234	

Supplemental Table 2G. Reclassification Tables for Clinical Model + LA + MAC vs Clinical Model

Stroke		Clinical Model + LA + MAC				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	22 (88.0%)	3 (12.0%)	0 (0%)	25	12.0
N (%)	5-9.99%	8 (15.1%)	32 (60.4%)	13 (24.5%)	53	39.6
	≥10%	0 (0%)	4 (6.8%)	55 (93.2%)	59	6.8
Total					137	

No Stroke		Clinical Model + LA + MAC				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	882 (86.4%)	137 (13.4%)	2 (0.2%)	1021	13.6
N (%)	5-9.99%	128 (17.6%)	500 (68.6%)	101 (13.9%)	729	31.4
	≥10%	0 (0%)	94 (19.4%)	390 (80.6%)	484	19.4
Total		·			2,234	

Supplemental Table 2H. Reclassification Tables for Clinical Model + LA + HbA1c vs Clinical Model

Stroke		Clin				
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model	<5%	18 (69.2%)	8 (30.8%)	0 (0%)	26	30.8
N (%)	5-9.99%	9 (17.3%)	23 (44.2%)	20 (38.5%)	52	55.8
	≥10%	0 (0%)	6 (11.3%)	47 (88.7%)	53	11.3
Total					131	

No Stroke		Clinic				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	789 (80.8%)	180 (18.4%)	7 (0.7%)	976	19.1
N (%)	5-9.99%	152 (20.7%)	413 (56.2%)	170 (23.1%)	735	43.8
	≥10%	7 (1.6%)	88 (19.8%)	349 (78.6%)	444	21.4
Total					2,155	

Supplemental Table 2I. Reclassification Tables for Clinical Model + LA + HbA1c vs Clinical Model + HbA1c

Stroke		Clinical Model + LA + HbA1c				
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model +	<5%	18 (78.3%)	5 (21.7%)	0 (0%)	23	21.7
HbA1c	5-9.99%	9 (18.8%)	25 (52.1%)	14 (29.2%)	48	48.0
N (%)	≥10%	0 (0%)	7 (11.7%)	53 (88.3%)	60	11.7
Total		·			131	

No Stroke		Clinical Model + LA + HbA1c				
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model +	<5%	846 (86.1%)	136 (13.8%)	1 (0.1%)	983	13.9
HbA1c	5-9.99%	101 (14.6%)	471 (67.9%)	122 (17.6%)	694	32.2
N (%)	≥10%	1 (0.1%)	74 (15.5%)	403 (84.3%)	478	15.6
Total					2,155	

Supplemental Table 2J. Reclassification Tables for Clinical Model + LA + HbA1c + UACR vs Clinical Model

Stroke		Clinical N				
		<5%	Total	% Reclassified		
Clinical Model	<5%	17 (65.4%)	9 (34.6%)	0 (0%)	26	34.6
N (%)	5-9.99%	11 (21.2%)	23 (44.2%)	18 (34.6%)	52	55.8
	≥10%	0 (0%)	9 (17.0%)	44 (83.0%)	53	17.0
Total					131	

No Stroke		Clinical M				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	819 (83.9%)	146 (15.0%)	11 (1.1%)	976	16.1
N (%)	5-9.99%	206 (28.0%)	391 (53.2%)	138 (18.8%)	735	46.8
	≥10%	15 (3.4%)	102 (23.0%)	327 (73.6%)	444	26.4
Total					2,155	

Supplemental Table 2K. Reclassification Tables for Clinical Model + LA + HbA1c + UACR vs Clinical Model + HbA1c

Stroke		Clinical Model + LA + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model +	<5%	21 (63.6%)	12 (36.4%)	0 (0%)	33	36.4
HbA1c	5-9.99%	7 (15.2%)	24 (52.2%)	15 (32.6%)	46	47.8
N (%)	≥10%	0 (0%)	5 (9.6%)	47 (90.4%)	52	9.6
Total					131	

No Stroke		Clinical Model + LA + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			% Reclassified
Clinical Model +	<5%	933 (85.2%)	158 (14.4%)	4 (0.4%)	1095	14.8
HbA1c	5-9.99%	106 (16.0%)	421 (63.4%)	137 (20.6%)	664	36.6
N (%)	≥10%	1 (0.3%)	60 (15.2%)	335 (84.6%)	396	15.5
Total					2,155	

Supplemental Table 2L Reclassification Tables for Clinical Model + LA + HbA1c + UACR vs Clinical Model + HbA1 + UACR

Stroke		Clinical Model + LA + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model +	<5%	20 (74.1%)	7 (25.9%)	0 (0%)	27	25.9
HbA1c + UACR	5-9.99%	8 (18.2%)	27 (61.4%0	9 (20.5%)	44	38.7
N (%)	≥10%	0 (0%)	7 (11.7%)	53 (88.3%)	60	11.7
Total					131	

No Stroke		Clinical M				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model +	<5%	897 (90.5%)	94 (9.5%)	0 (0%)	991	9.5
HbA1c + UACR	5-9.99%	143 (20.9%)	462 (67.5%)	79 (11.5%)	684	32.4
N (%)	≥10%	0 (0%)	83 (17.3%)	397 (82.7%)	480	17.3
Total					2,155	

Supplemental Table 2M. Reclassification Tables for Clinical Model + LA + MAC + HbA1c vs Clinical Model

Stroke		Clinical Model + LA + MAC + HbA1c				
		<5% 5-9.99% ≥10%			Total	% Reclassified
Clinical Model	<5%	19 (73.1%)	6 (23.1%)	1 (3.8%)	26	26.9
N (%)	5-9.99%	10 (19.2%)	21 (40.4%)	21 (40.4%)	52	59.6
	≥10%	0 (0%)	6 (11.3%)	47 (88.7%)	53	11.3
Total					131	

No Stroke		Clinical Model + LA + MAC + HbA1c				
		<5%	Total	% Reclassified		
Clinical Model	<5%	796 (81.6%)	170 (17.4%)	10 (1.0%)	976	18.4
N (%)	5-9.99%	169 (23.0%)	405 (55.1%)	161 (21.9%)	735	44.9
	≥10%	9 (2.0%)	99 (22.3%)	336 (75.7%)	444	24.3
Total					2,155	

Supplemental Table 2N. Reclassification Tables for Clinical Model + LA + MAC + HbA1c vs Clinical Model + HbA1c

Stroke		Clinical Model + LA + MAC + HbA1c				
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model +	<5%	24 (72.7%)	9 (27.3%)	0 (0%)	33	27.3
HbA1c	5-9.99%	5 (10.9%)	21 (45.7%)	20 (43.5%)	46	54.4
N (%)	≥10%	0 (0%)	3 (5.8%)	49 (94.2%)	52	5.8
Total		·			131	

No Stroke		Clinical Model + LA + MAC + HbA1c				
		<5%	Total	% Reclassified		
Clinical Model +	<5%	900 (82.2%)	192 (17.5%)	3 (0.3%)	1095	17.8
HbA1c	5-9.99%	74 (11.1%)	436 (65.7%)	154 (23.2%)	664	34.3
N (%)	≥10%	0 (0%)	46 (11.6%)	350 (88.4%)	396	11.6
Total					2,155	

Supplemental Table 2O. Reclassification Tables for Clinical Model + LA + MAC + HbA1c + UACR vs Clinical Model

Stroke		Clinical Model + LA + MAC + HbA1c + UACR				
		<5% 5-9.99% ≥10%				% Reclassified
Clinical Model	<5%	17 (65.4%)	8 (30.8%)	1 (3.8%)	26	34.6
N (%)	5-9.99%	10 (19.2%)	21 (40.4%)	21 (40.4%)	52	59.6
	≥10%	0 (0%)	8 (15.1%)	45 (84.9%)	53	15.1
Total					131	

No Stroke		Clinical Model + LA + MAC + HbA1c + UACR				
		<5%	<5% 5-9.99% ≥10%			
Clinical Model	<5%	800 (82.0%)	162 (16.6%)	14 (1.4%)	976	18.0
N (%)	5-9.99%	179 (24.4%)	393 (53.5%)	163 (22.2%)	735	46.6
	≥10%	11 (2.5%)	95 (21.4%)	338 (76.1%)	444	23.9
Total					2,155	

Supplemental Table 2P. Reclassification Tables for Clinical Model + LA + MAC + HbA1c + UACR vs Clinical Model + HbA1c + UACR

Stroke		Clinical Model + LA + MAC + HbA1c + UACR				
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model +	<5%	20 (74.1%)	7 (25.9%)	0 (0%)	27	25.9
HbA1c + UACR	5-9.99%	7 (15.9%)	24 (54.5%)	13 (39.5%)	44	45.4
N (%)	≥10%	0 (0%)	6 (10.0%)	90.0%)	60	10.0
Total					131	

No Stroke		Clinical Model + LA + MAC + HbA1c + UACR				
		<5%	Total	% Reclassified		
Clinical Model +	<5%	872 (88.0%)	118 (11.9%)	1 (0.1%)	991	12.0
HbA1c + UACR	5-9.99%	118 (17.3%)	464 (67.8%)	102 (14.9%)	684	32.2
N (%)	≥10%	0 (0%)	68 (14.2%)	412 (85.8%)	480	14.2
Total					2,155	

Table 2Q. Reclassification Tables for Clinical Model + LA + MAC + HbA1c + UACR vs Clinical Model + LA + HbA1c

Stroke		Clinical Mode				
		<5% 5-9.99% ≥10%			Total	% Reclassified
Clinical Model + LA +	<5%	24 (88.9%)	3 (11.1%)	0 (0%)	27	11.1
HbA1c	5-9.99%	3 (8.1%)	29 (78.4%)	5 (13.5%)	37	21.6
N (%)	≥10%	0 (0%)	5 (7.5%)	62 (92.5%)	67	7.5
Total					131	

No Stroke		Clinical Model + LA + MAC + HbA1c + UACR				
		<5% 5-9.99% ≥10%				% Reclassified
Clinical Model + LA +	<5%	900 (94.9%)	48 (5.1%)	0 (0%)	948	5.1
HbA1c	5-9.99%	90 (13.2%)	535 (78.6%)	56 (8.2%)	681	21.4
N (%)	≥10%	0 (0%)	67 (12.7%)	459 (87.3%)	526	12.7
Total		·			2,155	

Table 2R. Reclassification Tables for Clinical Model + LA +MAC + HbA1c + UACR vs Clinical Model + LA + MAC + HbA1c

Stroke		Clinical Model + LA + MAC + HbA1c + UACR				
		<5%	5-9.99%	≥10%	Total	% Reclassified
Clinical Model	<5%	25 (86.2%)	4 (13.8%)	0 (0%)	29	13.8
LA + MAC + HbA1c	5-9.99%	2 (6.1%)	29 (87.9%)	2 (6.1%)	33	12.2
N (%)	≥10%	0 (0%)	4 (5.8%)	65 (94.2%)	69	5.8
Total					131	

No Stroke		Clinical Model +				
		<5%	Total	% Reclassified		
Clinical Model	<5%	919 (94.4%)	55 (5.6%)	0 (0%)	974	5.6
LA + MAC + HbA1c	5-9.99%	71 (10.5%)	554 (82.2%)	49 (7.3%)	674	17.7
N (%)	≥10%	0 (0%)	41 (8.1%)	466 (91.9%)	507	7.9
Total					2,155	