

Supplemental Table 1. Q-PCR primers and probes used in this study.

Target Gene	Forward Primer (5' → 3')	Probe (5' → 3')	Reverse Primer (5' → 3')	Source
<i>rpoB</i>	CCGGACGTCACGGTAACAA	TTATCTCCCGTATTTTACC	CAGGTGTTCCGTCTGGCATA	(1)
<i>betL</i>	GCACCTGATGCGCCTGAT	TTGACTCATGGACTATTT	GAAAGCCACCAAGCCCAAT	(1)
<i>bsh</i>	GGCCTTAGTATGGCAGGACTCA	CCTTGTAATCCGCATTTTCTGAGAA	CTCATTGTCCTTACCTTCTGCAAA	(8)
<i>ctc</i>	CAGTTTCCGTTGATTCTTTTGGAAA	CGCCGTATTTTCA	AATTTCTTGCCGTCCACAGTAAC	(7)
<i>dnaK</i>	CCTTACTCGTGCTAAATTTGATGAAC	TAACACATGACTTAGTTGAAC	TGACGAGTTGGCGCAATAGTAC	(4)
<i>gbuA</i>	GAAAAAGCATGAGTATGGTCTTCCA	AACTTTGGTTTATTTCCG	TTTGTCATTCTTGAATTTCAAG	(1)
<i>groES</i>	CGTTTCAGGTCGTGTGCTAGA	CGGAACAAAAGAACCAC	TGTGTCACCTTCTGCAACTTCA	(4)
<i>hly</i>	ATGGCACACCAGCATCTC	CCTGCAAGTCTTAAGAC	ATCCGCGTGTTCCTTTTCGAT	(3)
<i>inlA</i>	GACAAATGCTCAGGCAGCTACA	CAGCTCTAGCGGAAAA	TTTGCAGACCGTGTCTGTTA	(6)
<i>kat</i>	CATCGTGTAGGGCCAAACTACTT	AACTCCCAATTAACAGTCC	TCGCGACAGGCGCTTT	This work
<i>liaH</i>	GAAGAAGTGGCTCAATATGAAGAA	TTAGAAGTACAAAAGCAC	TGTTCCGACGCTTGATCGTA	This work
<i>liaR</i>	GGCGGAGACTCTGTTCTTGAA	CAGAAGTAACTGGC	TAGCGTTAGCCGTT	This work
<i>oppA</i>	GAAGATGCAAAATGGTCAAACG	CCTGTAAGTCAAAATGACTAT	TGCACGACGCCATGAGTAA	(2)
<i>opuCA</i>	ACATCGATAAAGGAGAATTTGTTTGT	TCGTTTTCCACAACCA	GCCGGTTAATCATCTTCATTGTT	(1)
<i>plcA</i>	GATTTATTTACGACGCACATTAGTTT	CCCATTAGGCGGAAAAGCATATTCGC	GAGTTCCTTATTGGCTTATTCCAGTTATT	(5)
LMOh7858_1689	GCTTGGTCTGGTACATGTTATCGA	CGCGCGTTTTTCAT	CATGCTTGTATCGTAATTGGCTACA	This work
LMOh7858_3012	GGCAAAGAATTACCAAAGA	AGTACGTAGAAACCC	CGTAACCAATTAACCGGATACATGT	This work
lmo2754	CCCTTGAGTAAGTGGCTTCTT	CCGAAAAGCAGCACAAAG	TGCGTTTGCATTTAGATTTGG	This work
lmo2812	GGGATGAAAAGCTTGATCTTGTTT	TTGGATGATCCTTCCGC	TGTAATCGCGTAAAGGGAAACC	This work

References:

1. **Bergholz, T. M., H. C. den Bakker, E. D. Fortes, K. J. Boor, and M. Wiedmann.** 2010. Salt stress phenotypes in *Listeria monocytogenes* vary by genetic lineage and temperature. *Foodborne Pathog Dis* **7**:1537-49.
2. **Chan, Y. C., S. Raengpradub, K. J. Boor, and M. Wiedmann.** 2007. Microarray-based characterization of the *Listeria monocytogenes* cold regulon in log- and stationary-phase cells. *Appl Environ Microbiol* **73**:6484-98.
3. **Chaturongakul, S., and K. J. Boor.** 2006. SigmaB activation under environmental and energy stress conditions in *Listeria monocytogenes*. *Appl Environ Microbiol* **72**:5197-203.
4. **Hu, Y., H. F. Oliver, S. Raengpradub, M. E. Palmer, R. H. Orsi, M. Wiedmann, and K. J. Boor.** 2007. Transcriptomic and phenotypic analyses suggest a network between the transcriptional regulators HrcA and sigmaB in *Listeria monocytogenes*. *Appl Environ Microbiol* **73**:7981-91.
5. **Kazmierczak, M. J., M. Wiedmann, and K. J. Boor.** 2006. Contributions of *Listeria monocytogenes* sigmaB and PrfA to expression of virulence and stress response genes during extra- and intracellular growth. *Microbiology* **152**:1827-38.

6. **Oliver, H. F., R. H. Orsi, M. Wiedmann, and K. J. Boor.** 2010. *Listeria monocytogenes* {sigma}B has a small core regulon and a conserved role in virulence but makes differential contributions to stress tolerance across a diverse collection of strains. *Appl Environ Microbiol* **76**:4216-32.
7. **Raengpradub, S., M. Wiedmann, and K. J. Boor.** 2008. Comparative analysis of the sigma B-dependent stress responses in *Listeria monocytogenes* and *Listeria innocua* strains exposed to selected stress conditions. *Appl Environ Microbiol* **74**:158-71.
8. **Sue, D., D. Fink, M. Wiedmann, and K. J. Boor.** 2004. sigmaB-dependent gene induction and expression in *Listeria monocytogenes* during osmotic and acid stress conditions simulating the intestinal environment. *Microbiology* **150**:3843-55.