

eTable 1: Coffee consumption and time to pregnancy by age at study entry, smoking status, menstrual cycle regularity, number of cycle attempts before study entry, and parity^a

	None	Coffee (servings/day)			
		<1	1	2	3+
Age at baseline					
<30	1.00	1.07 (0.94-1.23)	1.19 (1.00-1.41)	1.14 (0.89-1.45)	1.36 (0.99-1.88)
$30+$	1.00	0.96 (0.79-1.16)	0.97 (0.79-1.21)	1.07 (0.84-1.36)	0.84 (0.60-1.18)
Smoking status					
<i>Never</i>	1.00	1.01 (0.88-1.16)	1.12 (0.94-1.34)	1.08 (0.84-1.38)	0.83 (0.55-1.23)
<i>Ever</i>	1.00	1.06 (0.88-1.28)	1.04 (0.84-1.28)	1.12 (0.88-1.42)	1.16 (0.86-1.57)
Regular cycles					
<i>Yes</i>	1.00	1.04 (0.91-1.17)	1.02 (0.87-1.19)	1.02 (0.84-1.23)	1.04 (0.80-1.34)
<i>No</i>	1.00	0.98 (0.77-1.25)	1.30 (0.97-1.72)	1.48 (1.02-2.14)	1.18 (0.67-2.09)
Cycle attempts before study entry					
≤ 2	1.00	1.05 (0.93-1.20)	1.08 (0.93-1.27)	1.15 (0.95-1.40)	1.12 (0.85-1.46)
$3-6$	1.00	0.93 (0.75-1.16)	1.10 (0.84-1.44)	0.98 (0.70-1.38)	0.87 (0.54-1.40)
Parity					
<i>Nulliparous</i>	1.00	1.05 (0.92-1.20)	1.11 (0.94-1.32)	1.07 (0.84-1.35)	1.23 (0.88-1.72)
<i>Parous</i>	1.00	1.00 (0.82-1.22)	1.05 (0.83-1.31)	1.04 (0.81-1.33)	0.85 (0.61-1.18)
Alcohol consumption					
<i><3 drinks/week</i>	1.00	1.03 (0.90-1.19)	1.11 (0.93-1.33)	1.16 (0.92-1.46)	0.94 (0.67-1.31)
<i>3+ drinks/week</i>	1.00	1.02 (0.84-1.27)	1.07 (0.86-1.32)	1.09 (0.84-1.41)	1.19 (0.85-1.66)

a Adjusted for age, partner age, body mass index, pack years of smoking, alcohol (drinks/week), frequency of intercourse, physical activity (mets/week) and consumption of tea and colas.

eTable 2: Tea consumption and time to pregnancy by age at study entry, smoking status, menstrual cycle regularity, number of cycle attempts before study entry, and parity^a

Characteristic	Tea (servings/day)			
	None	<1	1	2+
Age at baseline				
<30	1.00	1.04 (0.92-1.17)	1.11 (0.88-1.40)	1.23 (0.83-1.83)
$30+$	1.00	1.27 (1.08-1.50)	0.98 (0.75-1.29)	1.27 (0.90-1.78)
Smoking status				
Never	1.00	1.09 (0.97-1.23)	1.09 (0.88-1.36)	1.18 (0.84-1.66)
Ever	1.00	1.14 (0.98-1.33)	0.97 (0.72-1.30)	1.40 (0.95-2.07)
Regular cycles				
Yes	1.00	1.13 (1.02-1.26)	1.00 (0.82-1.22)	1.18 (0.88-1.57)
No	1.00	1.03 (0.84-1.27)	1.12 (0.77-1.64)	1.81 (1.02-3.21)
Cycle attempts before study entry				
≤ 2	1.00	1.10 (0.98-1.23)	0.93 (0.76-1.14)	1.22 (0.91-1.64)
3-6	1.00	1.12 (0.93-1.36)	1.40 (0.98-2.00)	1.48 (0.89-2.46)
Parity				
Nulliparous	1.00	1.13 (1.01-1.27)	1.03 (0.82-1.30)	1.41 (1.00-1.99)
Parous	1.00	1.17 (0.99-1.38)	1.03 (0.77-1.37)	1.10 (0.75-1.62)
Alcohol consumption				
<3 drinks/week	1.00	1.12 (0.99-1.26)	1.02 (0.80-1.29)	1.17 (0.83-1.67)
3+ drinks/week	1.00	1.09 (0.94-1.28)	1.07 (0.82-1.40)	1.42 (0.97-2.07)

a Adjusted for age, partner age, body mass index, pack years of smoking, alcohol (drinks/week), frequency of intercourse, physical activity (mets/week) and consumption of coffee and colas.

eTable 3: Total soda consumption and time to pregnancy by age at study entry, smoking status, menstrual cycle regularity, number of cycle attempts before study entry, and parity^a

Characteristic	Sodas (servings/day)			
	None	<1	1	2+
Age at baseline				
<30	1.00	0.91 (0.80-1.04)	0.83 (0.66-1.06)	0.81 (0.52-1.25)
30+	1.00	0.86 (0.73-1.01)	0.86 (0.64-1.16)	0.54 (0.28-1.04)
Smoking status				
Never	1.00	0.93 (0.82-1.06)	0.87 (0.68-1.11)	0.72 (0.43-1.22)
Ever	1.00	0.83 (0.70-0.98)	0.83 (0.63-1.09)	0.75 (0.46-1.25)
Regular cycles				
Yes	1.00	0.90 (0.80-1.01)	0.84 (0.68-1.03)	0.71 (0.47-1.08)
No	1.00	0.90 (0.72-1.12)	0.92 (0.62-1.37)	0.92 (0.42-2.02)
Cycle attempts before study entry				
≤2	1.00	0.90 (0.80-1.01)	0.87 (0.70-1.07)	0.64 (0.41-1.00)
3-6	1.00	0.88 (0.72-1.08)	0.81 (0.56-1.19)	0.97 (0.52-1.83)
Parity				
Nulliparous	1.00	0.91 (0.80-1.03)	0.94 (0.75-1.19)	0.62 (0.38-1.00)
Parous	1.00	0.83 (0.70-0.99)	0.65 (0.48-0.88)	0.93 (0.53-1.61)
Alcohol consumption				
<3 drinks/week	1.00	0.93 (0.82-1.06)	0.88 (0.70-1.11)	0.80 (0.51-1.24)
3+ drinks/week	1.00	0.83 (0.69-0.99)	0.81 (0.59-1.11)	0.61 (0.32-1.17)

a Adjusted for age, partner age, body mass index, pack years of smoking, alcohol (drinks/week), frequency of intercourse, physical activity (mets/week), and caffeine consumption.