

Appendix 1 (as supplied by authors): Participation rates in cervical cancer screening for Aboriginal and non-Aboriginal women in Canada

Source*	year	Population / locality*	At least one Pap test in:	Participation rates (%)	
				Aboriginal	Non-Aboriginal
Hislop et al(2)	1990-92	FN / BC	3 years	62	84
Young et al(1)	1993-95	FN / Manitoba	3 years	42.6	60.4
Statistic Canada(6)	2000, 03	Aboriginal / Canada	3 years	85.3	86.5
Statistic Canada(3)	2000, 04	Inuit / Nunaat	3 years	75.4	80.1
Statistics Canada(4,5)	2005	Inuit / Nunavut	3 years	79.3	72.8 [†]
AFN(7)	2002	FN / Canada	3 years	75-80	N/A
Louchini et al(8)	2004	Inuit / Nunavik	2 years	77.1	N/A
Hamlin-Douglas et al(9)	2002-07	Inuit / Nunavik	3 years	71	N/A
Martens et al(10)	2004-06	Métis / MB	3 years	69.0	67.8
NAHO(13)	2002	FN / Canada	1 year	50.0	N/A
NWT Social Services(11)	2000-01	Canadian / NWT	3 years		79 [‡]
McDermott(12)	1997-00	Aboriginal / NWT	4 years	72.8	89.8

* AFN: Assembly of First Nations; FN: First Nations; NAHO: National Aboriginal Health Organization; NWT: Northwest Territories; BC: British Columbia

[†] Canadian population including Aboriginal groups

[‡] Entire Northwest Territories population