

Supplement Table II. Characteristics of cases with acute pancreatitis and frequency matched population control subjects in Sweden during the period 2006-2008. Generic name for antipsychotic drugs used time before index date.

	<i>Cases</i>		<i>Controls</i>	
	<i>Number</i>	<i>%</i>	<i>Number</i>	<i>%</i>
<i>Total</i>	6161	100.0	61637	100.0
<i>Use of atypicals</i>				
<i>No use</i>	6051	98.2	60877	98.8
<i>Current (0-114 days)</i>	81	1.3	565	0.9
<i>Recent (115-180 days)</i>	4	0.1	32	0.1
<i>Past (6-12 months)</i>	12	0.2	66	0.1
<i>Former (>12 months)</i>	13	0.2	97	0.2
<i>Use of high potency typicals</i>				
<i>No use</i>	6107	99.1	61303	99.5
<i>Current (0-114 days)</i>	29	0.5	199	0.3
<i>Recent (115-180 days)</i>	6	0.1	28	0.0
<i>Past (6-12 months)</i>	8	0.1	50	0.1
<i>Former (>12 months)</i>	11	0.2	57	0.1
<i>Use of medium potency typicals</i>				
<i>No use</i>	6067	98.5	61159	99.2
<i>Current (0-114 days)</i>	33	0.5	219	0.4
<i>Recent (115-180 days)</i>	6	0.1	44	0.1
<i>Past (6-12 months)</i>	12	0.2	73	0.1
<i>Former (>12 months)</i>	43	0.7	142	0.2
<i>Use of low potency typicals</i>				
<i>No use</i>	6096	98.9	61326	99.5
<i>Current (0-114 days)</i>	31	0.5	205	0.3
<i>Recent (115-180 days)</i>	8	0.1	17	0.0
<i>Past (6-12 months)</i>	11	0.2	37	0.1
<i>Former (>12 months)</i>	15	0.2	52	0.1