

Table S1. Raw means (main table entries) and standard errors (in parentheses) of dry body mass of mosquitoes measured in this study

Sample location	Sex and female gonotrophic state	Sampling period				
		Oct-Nov	Dec	Jan-Feb	Apr-May	Jul-Aug
Sahelian (MP)	Male	0.194 (0.016)	0.196 (0.004)	0.200 (0.005)	0.201 (0.014)	0.174 (0.006)
	Unfed Female	0.288 (0.020)	0.287 (0.040)	0.290 (0.017)	0.218 (0.014)	0.230 (0.015)
	Fed Female	0.743 (0.056)	0.994 (n/a)	0.965 (0.359)	0.760 (0.065)	n/a
	Gravid Female	0.481 (0.041)	0.437 (0.247)	0.438 (0.077)	0.523 (0.059)	0.269 (0.023)
Riparian (NG)	Male	0.200 (0.011)	0.180 (0.008)	0.203 (0.006)	0.160 (0.008)	0.191 (0.017)
	Unfed Female	0.241 (0.013)	0.230 (0.016)	0.220 (0.022)	0.200 (0.011)	0.237 (0.017)
	Fed Female	0.850 (0.056)	1.059 (0.352)	0.890 (n/a)	0.386 (0.056)	n/a
	Gravid Female	0.548 (0.030)	0.371 (0.081)	0.364 (0.043)	0.415 (0.069)	0.301 (0.024)

Sampling periods are defined as in all other tables and figures. For the mean; n/a indicates that no individuals of that gonotrophic state were collected during that sampling period; while for the standard error; n/a indicates that only 1 individual was measured.