

Supplementary information

CDK targeting of NBS1 promotes DNA-end resection, replication restart and homologous recombination

Jacob Falck, Josep V. Forment, Julia Coates, Martin Mistrik, Jiri Lukas, Jiri Bartek and Stephen P. Jackson

Supplementary methods

CDK kinase assays

CDK1/Cyclin B was from Millipore and purified MRN was a gift from T. Paull (University of Texas, Austin). Assays were performed in CDK1/Cyclin B buffer for 10 min at 37 °C in the presence of ³²P-γ-ATP.

Western blots and immunoprecipitations

Anti-NBS1 pSer-432 was generated as a rabbit polyclonal antibody (now available from Abcam). Other antibodies used: NBS1 (rabbit; Merck), MRE11 (rabbit; Novus), Cyclin A (rabbit; Santa Cruz), Histone H3 pSer-10 (rabbit; Millipore), ATM pSer-1981 (rabbit; Rockland), ATM (mouse; gift from Y. Shiloh, Tel Aviv), NBS1 pSer-343 (rabbit; Cell Signalling), CHK1 pSer-317 (rabbit; Bethyl), CHK1 (mouse; Santa Cruz), GFP (mouse; Roche), CDK1 (mouse; Oncogene), CDK2 (rabbit, Santa Cruz). NBS1 was immunoprecipitated with a mouse monoclonal antibody (Abcam; 2 μl/mg protein extract). GFP immunoprecipitations were performed with GFP-Trap® beads (Chromotek).

Transfections

GFP-CtIP (Sartori et al, 2007) was transfected in NBS-ILB1 cells using FuGENE HD (Roche) and following manufacturer's instructions. siRNA transfection was performed in U2OS cells using Lipofectamine RNAiMax (Invitrogen) following manufacturer's instructions. Sequences of the siRNAs used: luciferase 5'-cguacgcggaaacuucga-tt-3', CDK1 5'-gggguuccuaguacugcaa-tt-3', CDK2 5'-gccagaaacaaguugacgg-tt-3'.

Immunofluorescence

Primary antibody staining was for 1 h in 5% FBS in 1xPBS (NBS1, rabbit; Merck), γ H2AX (mouse or rabbit; Upstate and Cell Signalling, respectively), RPA2 (mouse; Abcam). Secondary antibody staining was done with Alexa Fluor 488 or 594 (Molecular Probes) for 30 min. All incubations were at room temperature. To detect ssDNA, cells were labeled with BrdU for 36 h, washed, and cultured in the absence of BrdU for 24 h before the experiment. BrdU was detected under native conditions with an anti-BrdU antibody (mouse; GE Healthcare).

EdU incorporation assay

Cells were pulsed with 10 μ M EdU for 30 min before collection. EdU detection was performed with Click-iT® EdU Flow Cytometry kit (Invitrogen) following manufacturer's instructions.


Supplementary figure legends

Figure S1: EdU incorporation is similar in all used cell lines. The *y* axis represents intensity of EdU signal (EdU incorporation). The *x* axis represents DNA content as measured by intensity of 4',6-diamidino-2-phenylindole (DAPI) signal.

Figure S2: HR assay. (A) Schematic of the HR assay system introduced in NBS-ILB1 cells. (B) Only cells positive for RFP (indicating transfection with the *I-SceI* endonuclease) and GFP (indicating recombination between the GFP fragments) are counted as HR events in the assay. (C) Representative images from one of the HR assays. Percentages indicate the amount of HR events.


FALCK et al SUPP. FIGURE S1

NBS-ILB1 cells




FALCK et al SUPP. FIGURE S2

A



B



C

