	Probability of being the best treatment choice in terms of healing (%)										
	Ahroni 1993 removed	Donaghue 1998 removed	Blackman 1994 removed	Mazzone 1993 removed	Roberts 2001 removed	Piaggesi 2001 removed	D'Hemecourt 1998 removed	Vandeputte 1997 removed	Jenson 1998 removed	Baker 1993 removed	Foster 1994 removed
Matrix hydrocolloid	71	66	67	67	72	70	35	73	72	63	75
Hydrogel	9	21	17	18	10	14	62	8	9	21	6
Foam	14	11	12	12	16	13	2	14	15	9	18
Silver fibrous hydrocolloid	6	2	3	3	2	3	1	3	3	7	1
Basic wound contact layer	0	0	0	0	0	0	0	0	0	0	0
Alginate	0	0	0	0	0	0	0	0	0	0	0
Fibrous hydrocolloid	0	0	0	0	0	0	0	0	0	0	0
Impregnated-iodine	0	0	0	0	0	0	0	0	0	0	0
Protease modulating	0	0	0	0	0	0	0	0	0	0	0

ESM Table 4 Sensitivity analyses. Table shows the probability that each treatment is the best in terms of healing diabetic foot ulcers when one trial (named in column) had been removed. Shaded cell highlights study with highest probability for each column.