

Chromophore Formation in Resorcinarene Solutions and the Visual Detection of Mono- and Oligosaccharides

*Ming He, Rolanda J. Johnson, Jorge O. Escobedo, Patricia A. Beck, Kyu Kwang Kim, Nadia N. St. Luce, Claude J. Davis, Patrick T. Lewis, Frank R. Fronczek, Bruce J. Melancon, Anthony A. Mrse, W. Dale Treleaven and Robert M. Strongin**

Contribution from the Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803

Supporting Information

Table of Crystallographic Data for Compound 7.

Compound	7•2MeOH
Formula	C ₃₂ H ₃₂ O ₈ •2 CH ₃ OH
Color/shape	colorless fragment
Formula weight	608.7
Crystal system	Orthorhombic
Space group	Pnma
Temp., K	120
Cell constants	
a, Å	14.0627(4)
b, Å	13.6940(4)
c, Å	16.0505(5)
Cell volume, Å ³	3090.92(16)
Formula units/unit cell	4
Dcalc, g cm ⁻³	1.308
μcalc, cm ⁻¹	0.96
Diffractometer / scan	Nonius KappaCCD/ ω
Radiation, graphite monochr.	MoKα ($\lambda = 0.71073 \text{ \AA}$)
Max. crystal dimensions, mm	0.27 x 0.37 x 0.42
Reflections measured	14,994
R _{int}	0.037
Independent reflections	4660

2θ range, deg	5<2θ<60
Range of h,k,l	±19, ±19, ±22
Reflections observed	3626
Criterion for observed	I>2σ(I)
Data/parameters	4660/240
R (obs)	0.064
R(all data)	0.090
GOF	1.116
Max resid. peaks (eÅ ⁻³)	0.39, -0.61